



Nijat Aliyev

BHOS

Calculus

October 11, 2023

## Rate of Change

Let  $y = f(x)$  be a given function.

## Rate of Change

Let  $y = f(x)$  be a given function.

We consider  $\frac{f(x+h)-f(x)}{h}$  as the average rate of change in  $f$  over the interval from  $x$  to  $x + h$ .

## Rate of Change

Let  $y = f(x)$  be a given function.

We consider  $\frac{f(x+h)-f(x)}{h}$  as the average rate of change in  $f$  over the interval from  $x$  to  $x + h$ .

We interpret the limit of the above fraction as  $h \rightarrow 0$  as the rate at which  $f$  is changing at  $x$ .

## Rate of Change

Let  $y = f(x)$  be a given function.

We consider  $\frac{f(x+h)-f(x)}{h}$  as the average rate of change in  $f$  over the interval from  $x$  to  $x + h$ .

We interpret the limit of the above fraction as  $h \rightarrow 0$  as the rate at which  $f$  is changing at  $x$ .

Thus, instantaneous rates are limits of average rates..

## Rate of Change

Let  $y = f(x)$  be a given function.

We consider  $\frac{f(x+h)-f(x)}{h}$  as the average rate of change in  $f$  over the interval from  $x$  to  $x + h$ .

We interpret the limit of the above fraction as  $h \rightarrow 0$  as the rate at which  $f$  is changing at  $x$ .

Thus, instantaneous rates are limits of average rates..

**DEFINITION** The **instantaneous rate of change** of  $f$  with respect to  $x$  at  $x_0$  is the derivative

$$f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h},$$

provided the limit exists.

## Rate of Change

Let  $y = f(x)$  be a given function.

We consider  $\frac{f(x+h)-f(x)}{h}$  as the average rate of change in  $f$  over the interval from  $x$  to  $x + h$ .

We interpret the limit of the above fraction as  $h \rightarrow 0$  as the rate at which  $f$  is changing at  $x$ .

Thus, instantaneous rates are limits of average rates..

**DEFINITION** The **instantaneous rate of change** of  $f$  with respect to  $x$  at  $x_0$  is the derivative

$$f'(x_0) = \lim_{h \rightarrow 0} \frac{f(x_0 + h) - f(x_0)}{h},$$

provided the limit exists.

## Rate of Change

**Example:** The area  $A$  of a circle is related to its diameter by the equation

$$A = \frac{\pi}{4}D^2$$

How fast does the area change with respect to the diameter when the diameter is 10 m?



## Rate of Change

**Example:** The area  $A$  of a circle is related to its diameter by the equation

$$A = \frac{\pi}{4}D^2$$

How fast does the area change with respect to the diameter when the diameter is 10 m?

**Solution:**

The rate of change of the area with respect to the diameter is

$$\frac{dA}{dD} = \frac{\pi}{4} \cdot 2D = \frac{\pi D}{2}.$$

## Rate of Change

**Example:** The area  $A$  of a circle is related to its diameter by the equation

$$A = \frac{\pi}{4}D^2$$

How fast does the area change with respect to the diameter when the diameter is 10 m?

**Solution:**

The rate of change of the area with respect to the diameter is

$$\frac{dA}{dD} = \frac{\pi}{4} \cdot 2D = \frac{\pi D}{2}.$$

When  $D = 10$  m, the area is changing with respect to the diameter at the rate of  $(\pi/2)10 = 5\pi \text{ m}^2/\text{m} \approx 15.71 \text{ m}^2/\text{m}$ . ■

## Velocity

Suppose the object is moving along a line and its position is given by the function  $s = f(t)$ , a function of time.

## Velocity

Suppose the object is moving along a line and its position is given by the function  $s = f(t)$ , a function of time.

The displacement of the object over the time interval from  $t$  to  $t + \Delta t$  is

$$\Delta s = f(t + \Delta t) - f(t)$$

## Velocity

Suppose the object is moving along a line and its position is given by the function  $s = f(t)$ , a function of time.

The displacement of the object over the time interval from  $t$  to  $t + \Delta t$  is

$$\Delta s = f(t + \Delta t) - f(t)$$

Its average velocity is

$$v_{av} = \frac{\text{Displacement}}{\text{Travel Time}} = \frac{\Delta s}{\Delta t} = \frac{f(t + \Delta t) - f(t)}{\Delta t}$$

To find the object's velocity at the exact instant  $t$ , we take the limit of the average velocity

## Velocity

Suppose the object is moving along a line and its position is given by the function  $s = f(t)$ , a function of time.

The displacement of the object over the time interval from  $t$  to  $t + \Delta t$  is

$$\Delta s = f(t + \Delta t) - f(t)$$

Its average velocity is

$$v_{av} = \frac{\text{Displacement}}{\text{Travel Time}} = \frac{\Delta s}{\Delta t} = \frac{f(t + \Delta t) - f(t)}{\Delta t}$$

To find the object's velocity at the exact instant  $t$ , we take the limit of the average velocity

Velocity (instantaneous velocity) is the derivative of position w.r.t time:

$$v(t) = \frac{ds}{dt} = \lim_{\Delta t \rightarrow 0} \frac{f(t + \Delta t) - f(t)}{\Delta t}$$

## Derivative of Trigonometric Functions

1. The derivative of the sine function is the cosine function:

$$\frac{d}{dx}(\sin x) = \cos x$$

## Derivative of Trigonometric Functions

1. The derivative of the sine function is the cosine function:

$$\frac{d}{dx}(\sin x) = \cos x$$

**Proof:** We use the following identity

$$\sin(x + h) = \sin x \cos h + \cos x \sin h.$$



## Derivative of Trigonometric Functions

1. The derivative of the sine function is the cosine function:

$$\frac{d}{dx}(\sin x) = \cos x$$

**Proof:** We use the following identity

$$\sin(x + h) = \sin x \cos h + \cos x \sin h.$$

If  $f(x) = \sin x$ , then

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin x}{h} && \text{Derivative definition} \\ &= \lim_{h \rightarrow 0} \frac{(\sin x \cos h + \cos x \sin h) - \sin x}{h} \\ &= \lim_{h \rightarrow 0} \frac{\sin x(\cos h - 1) + \cos x \sin h}{h} \\ &= \lim_{h \rightarrow 0} \left( \sin x \cdot \frac{\cos h - 1}{h} \right) + \lim_{h \rightarrow 0} \left( \cos x \cdot \frac{\sin h}{h} \right) \\ &= \sin x \cdot \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} + \cos x \cdot \lim_{h \rightarrow 0} \frac{\sin h}{h} = \sin x \cdot 0 + \cos x \cdot 1 = \cos x. \end{aligned}$$

**Example:** Find the following derivatives.

$$\begin{aligned} \text{(a)} \quad y = x^2 - \sin x: \quad \frac{dy}{dx} &= 2x - \frac{d}{dx}(\sin x) \\ &= 2x - \cos x \end{aligned}$$

$$\begin{aligned} \text{(b)} \quad y = x^2 \sin x: \quad \frac{dy}{dx} &= x^2 \frac{d}{dx}(\sin x) + 2x \sin x \\ &= x^2 \cos x + 2x \sin x. \end{aligned}$$

$$\begin{aligned} \text{(c)} \quad y = \frac{\sin x}{x}: \quad \frac{dy}{dx} &= \frac{x \cdot \frac{d}{dx}(\sin x) - \sin x \cdot 1}{x^2} \\ &= \frac{x \cos x - \sin x}{x^2} \end{aligned}$$

## Derivative of Trigonometric Functions

The derivative of the cosine function is the negative of the sine function:

$$\frac{d}{dx}(\cos x) = -\sin x$$

## Derivative of Trigonometric Functions

The derivative of the cosine function is the negative of the sine function:

$$\frac{d}{dx}(\cos x) = -\sin x$$

**Proof:** Using the identity  $\cos(x + h) = \cos x \cos h - \sin x \sin h$

## Derivative of Trigonometric Functions

The derivative of the cosine function is the negative of the sine function:

$$\frac{d}{dx}(\cos x) = -\sin x$$

**Proof:** Using the identity  $\cos(x + h) = \cos x \cos h - \sin x \sin h$

$$\begin{aligned}\frac{d}{dx}(\cos x) &= \lim_{h \rightarrow 0} \frac{\cos(x + h) - \cos x}{h} \\&= \lim_{h \rightarrow 0} \frac{(\cos x \cos h - \sin x \sin h) - \cos x}{h} \\&= \lim_{h \rightarrow 0} \frac{\cos x(\cos h - 1) - \sin x \sin h}{h} \\&= \lim_{h \rightarrow 0} \cos x \cdot \frac{\cos h - 1}{h} - \lim_{h \rightarrow 0} \sin x \cdot \frac{\sin h}{h} \\&= \cos x \cdot \lim_{h \rightarrow 0} \frac{\cos h - 1}{h} - \sin x \cdot \lim_{h \rightarrow 0} \frac{\sin h}{h} \\&= \cos x \cdot 0 - \sin x \cdot 1 \\&= -\sin x.\end{aligned}$$

**Example:** Find the following derivatives

(a)  $y = 5x + \cos x$ :

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(5x) + \frac{d}{dx}(\cos x) \\ &= 5 - \sin x.\end{aligned}$$

**Example:** Find the following derivatives

(a)  $y = 5x + \cos x$ :

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(5x) + \frac{d}{dx}(\cos x) \\ &= 5 - \sin x.\end{aligned}$$

(b)  $y = \sin x \cos x$ :

$$\begin{aligned}\frac{dy}{dx} &= \sin x \frac{d}{dx}(\cos x) + \cos x \frac{d}{dx}(\sin x) \\ &= \sin x(-\sin x) + \cos x(\cos x) \\ &= \cos^2 x - \sin^2 x\end{aligned}$$

$$(c) \quad y = \frac{\cos x}{1 - \sin x};$$

$$\begin{aligned}\frac{dy}{dx} &= \frac{(1 - \sin x) \frac{d}{dx}(\cos x) - \cos x \frac{d}{dx}(1 - \sin x)}{(1 - \sin x)^2} \\&= \frac{(1 - \sin x)(-\sin x) - \cos x(0 - \cos x)}{(1 - \sin x)^2} \\&= \frac{1 - \sin x}{(1 - \sin x)^2} \\&= \frac{1}{1 - \sin x}\end{aligned}$$



Using the identities:

$$\tan x = \frac{\sin x}{\cos x}, \quad \cot x = \frac{\cos x}{\sin x}, \quad \sec x = \frac{1}{\cos x}, \quad \text{and} \quad \csc x = \frac{1}{\sin x}$$

Using the identities:

$$\tan x = \frac{\sin x}{\cos x}, \quad \cot x = \frac{\cos x}{\sin x}, \quad \sec x = \frac{1}{\cos x}, \quad \text{and} \quad \csc x = \frac{1}{\sin x}$$

we can derive the derivatives of the other trigonometric functions

Using the identities:

$$\tan x = \frac{\sin x}{\cos x}, \quad \cot x = \frac{\cos x}{\sin x}, \quad \sec x = \frac{1}{\cos x}, \quad \text{and} \quad \csc x = \frac{1}{\sin x}$$

we can derive the derivatives of the other trigonometric functions

**The derivatives of the other trigonometric functions:**

$$\frac{d}{dx}(\tan x) = \sec^2 x$$

$$\frac{d}{dx}(\cot x) = -\csc^2 x$$

$$\frac{d}{dx}(\sec x) = \sec x \tan x$$

$$\frac{d}{dx}(\csc x) = -\csc x \cot x$$

**Example:** Find  $\frac{d}{dx}(\tan x)$ .

**Example:** Find  $\frac{d}{dx}(\tan x)$ .

**Solution :** We use the Derivative Quotient Rule to calculate the derivative:

$$\begin{aligned}\frac{d}{dx}(\tan x) &= \frac{d}{dx}\left(\frac{\sin x}{\cos x}\right) = \frac{\cos x \frac{d}{dx}(\sin x) - \sin x \frac{d}{dx}(\cos x)}{\cos^2 x} \\&= \frac{\cos x \cos x - \sin x(-\sin x)}{\cos^2 x} \\&= \frac{\cos^2 x + \sin^2 x}{\cos^2 x} \\&= \frac{1}{\cos^2 x} = \sec^2 x.\end{aligned}$$

**Example:** Find  $\frac{d}{dx}(\tan x)$ .

**Solution :** We use the Derivative Quotient Rule to calculate the derivative:

$$\begin{aligned}\frac{d}{dx}(\tan x) &= \frac{d}{dx}\left(\frac{\sin x}{\cos x}\right) = \frac{\cos x \frac{d}{dx}(\sin x) - \sin x \frac{d}{dx}(\cos x)}{\cos^2 x} \\&= \frac{\cos x \cos x - \sin x(-\sin x)}{\cos^2 x} \\&= \frac{\cos^2 x + \sin^2 x}{\cos^2 x} \\&= \frac{1}{\cos^2 x} = \sec^2 x.\end{aligned}$$

**Example:** Find  $y''$  if  $y = \sec x$ .

**Example:** Find  $y''$  if  $y = \sec x$ .

**Solution :**  $y = \sec x \implies y' = \sec x \tan x \implies y'' = \frac{d}{dx}(\sec x \tan x)$ .

$$\begin{aligned}y'' &= \frac{d}{dx}(\sec x \tan x) \\&= \sec x \frac{d}{dx}(\tan x) + \tan x \frac{d}{dx}(\sec x) \\&= \sec x(\sec^2 x) + \tan x(\sec x \tan x) \\&= \sec^3 x + \sec x \tan^2 x\end{aligned}$$



**Example:** Find  $y''$  if  $y = \sec x$ .

**Solution :**  $y = \sec x \implies y' = \sec x \tan x \implies y'' = \frac{d}{dx}(\sec x \tan x)$ .

$$\begin{aligned}y'' &= \frac{d}{dx}(\sec x \tan x) \\&= \sec x \frac{d}{dx}(\tan x) + \tan x \frac{d}{dx}(\sec x) \\&= \sec x(\sec^2 x) + \tan x(\sec x \tan x) \\&= \sec^3 x + \sec x \tan^2 x\end{aligned}$$