Dr. Nijat Aliyev

BHOS

Calculus

September 19, 2023

Function

Definition

A function f from a set A to a set B is a relation (rule) that assigns to each element (point) $x \in A$ a unique element $f(x) \in B$.

Function

Definition

A function f from a set A to a set B is a relation (rule) that assigns to each element (point) $x \in A$ a unique element $f(x) \in B$.

We usually use the notation y = f(x).

Function

Definition

A function f from a set A to a set B is a relation (rule) that assigns to each element (point) $x \in A$ a unique element $f(x) \in B$.

We usually use the notation y = f(x). Here, x is an independent variable or input and y is the dependent variable or output.

Function

Definition

A function f from a set A to a set B is a relation (rule) that assigns to each element (point) $x \in A$ a unique element $f(x) \in B$.

We usually use the notation y = f(x). Here, x is an independent variable or input and y is the dependent variable or output.

Another notation for function is $f: A \rightarrow B$.

Definition

A function f from a set A to a set B is a relation (rule) that assigns to each element (point) $x \in A$ a unique element $f(x) \in B$.

We usually use the notation y = f(x). Here, x is an independent variable or input and y is the dependent variable or output.

Another notation for function is $f: A \rightarrow B$.

The set A of all possible input values is called the domain of f.

Definition

A function f from a set A to a set B is a relation (rule) that assigns to each element (point) $x \in A$ a unique element $f(x) \in B$.

We usually use the notation y = f(x). Here, x is an independent variable or input and y is the dependent variable or output.

Another notation for function is $f: A \rightarrow B$.

The set A of all possible input values is called the domain of f.

The set B of all possible output values is called a codomain of f.

Definition

A function f from a set A to a set B is a relation (rule) that assigns to each element (point) $x \in A$ a unique element $f(x) \in B$.

We usually use the notation y = f(x). Here, x is an independent variable or input and y is the dependent variable or output.

Another notation for function is $f: A \rightarrow B$.

The set A of all possible input values is called the domain of f.

The set B of all possible output values is called a codomain of f.

The set of all actual output values of f(x) as x varies throughout the domain of f is called the range of f and is denoted by R_f .

Function

Note:

- 1. The domain and the range of a function can be any sets of objects but usually they are real numbers (in calculus course).
- 2. The range R_f of f may not include all elements in B but it is always true that $R_f \subseteq B$.

Function

Note:

- 1. The domain and the range of a function can be any sets of objects but usually they are real numbers (in calculus course).
- 2. The range R_f of f may not include all elements in B but it is always true that $R_f \subseteq B$.

Usually a function is given by a formula which determines how to calculate the output (dependent) value from the input (independent) variable.

Function

Note:

- 1. The domain and the range of a function can be any sets of objects but usually they are real numbers (in calculus course).
- 2. The range R_f of f may not include all elements in B but it is always true that $R_f \subseteq B$.

Usually a function is given by a formula which determines how to calculate the output (dependent) value from the input (independent) variable.

For instance, the equation $A=\pi r^2$ is a rule that calculates the area A of a circle with radius r.

Function

Note:

- 1. The domain and the range of a function can be any sets of objects but usually they are real numbers (in calculus course).
- 2. The range R_f of f may not include all elements in B but it is always true that $R_f \subseteq B$.

Usually a function is given by a formula which determines how to calculate the output (dependent) value from the input (independent) variable.

For instance, the equation $A=\pi r^2$ is a rule that calculates the area A of a circle with radius r.

Function

A function f can be considered as a machine that produces an output value f(x) in its range when we feed it an input value from its domain.

Function

A function f can be considered as a machine that produces an output value f(x) in its range when we feed it an input value from its domain.




Figure 1: A diagram showing a function as a kind of machine.

Function

A function can also be pictured as an arrow diagram (Figure 2).

Function

A function can also be pictured as an arrow diagram (Figure 2).




Figure 2: A function f from a set A to a set B.

Function

A function can also be pictured as an arrow diagram (Figure 2).




Figure 2: A function f from a set A to a set B.

Domain of f is A, codomain is B, the range is $\{f(x), f(a), f(b)\}$

Function

Each arrow associates an element of the domain A with a single element in B

Function

Each arrow associates an element of the domain A with a single element in B

The arrows indicate that f(a) is associated with a, f(x) is associated with x, and so on.

Function

Each arrow associates an element of the domain A with a single element in B

The arrows indicate that f(a) is associated with a, f(x) is associated with x, and so on.

Notice that a function can have the same value at two different input elements in the domain (as occurs with f(b) in Figure 2), but each input element x is assigned a single output value f(x).

Function

Domain and range of some functions

Domain and range of some functions

Function	Domain	Range
$f(x)=x^2$	$\mathbb{R}=(-\infty,\infty)$	$[0,\infty)$
$f(x)=\frac{1}{x}$	$\mathbb{R}\setminus\{0\}$	$\mathbb{R}\setminus\{0\}$
$f(x) = \sqrt{x}$	$[0,\infty)$	$[0,\infty)$
$f(x) = \sqrt{1-x}$	$[1,\infty)$	$[0,\infty)$
$f(x) = \sqrt{1 - x^2}$	$[0,\infty)$	[0, 1]

Graphs of function

The graph of a function f with domain D, is a set consists of the points in the Cartesian plane whose coordinates are the input-output pairs for f.

Graphs of function

The graph of a function f with domain D, is a set consists of the points in the Cartesian plane whose coordinates are the input-output pairs for f.

In set notation, the graph is

$$\{(x, f(x)) | x \in D\}$$

Graphs of function

The graph of a function f with domain D, is a set consists of the points in the Cartesian plane whose coordinates are the input-output pairs for f.

In set notation, the graph is

$$\big\{\big(x,f(x)\big)\,\big|\,x\in D\big\}$$




Figure 3: The graph of f(x) = x + 2 is the set of points (x, y) for which y has the value x + 2.

Graphs of function

The graph of a function f with domain D, is a set consists of the points in the Cartesian plane whose coordinates are the input-output pairs for f.

In set notation, the graph is

$$\big\{\big(x,f(x)\big)\,\big|\,x\in D\big\}$$




Figure 3: The graph of f(x) = x + 2 is the set of points (x, y) for which y has the value x + 2.

Graphs of a function

Not every curve in the coordinate plane can be the graph of a function.

Graphs of a function

Not every curve in the coordinate plane can be the graph of a function.

A function f can assign only one output value f(x) for each input value x in its domain, so any vertical line can not intersect the graph of a function more than once.

Graphs of a function

Not every curve in the coordinate plane can be the graph of a function.

A function f can assign only one output value f(x) for each input value x in its domain, so any vertical line can not intersect the graph of a function more than once.

If a is in the domain of the function f, then the vertical line x=a will intersect the graph of f at the single point (a, f(a)).

Graphs of function

Is the circle $x^2 + y^2 = 1$ the graph of any function?

Graphs of function

Is the circle $x^2 + y^2 = 1$ the graph of any function?

No. Why?

Graphs of function

Is the circle $x^2 + y^2 = 1$ the graph of any function?

No. Why?

It fails the vertical line test.

Graphs of function

Is the circle $x^2 + y^2 = 1$ the graph of any function?

No. Why?

It fails the vertical line test.




Figure 4: The circle $x^2+y^2=1$ is not the graph of a function, since it fails vertical line test.

Graphs of function

However, the upper and lower semicircles are the graphs of the functions $y=\sqrt{1-x^2}$ and $y=-\sqrt{1-x^2}$, respectively.

Graphs of function

However, the upper and lower semicircles are the graphs of the functions $y=\sqrt{1-x^2}$ and $y=-\sqrt{1-x^2}$, respectively.




Figure 5: The upper semi-circle is the graph of the function $y=\sqrt{1-x^2}$, while the lower semi-circle is the graph of the function $y=-\sqrt{1-x^2}$.

Piecewise-Defined Functions

Sometimes the domain of the function is given in pieces and the function is described by a different formula in each piece of domain.

Piecewise-Defined Functions

Sometimes the domain of the function is given in pieces and the function is described by a different formula in each piece of domain.

This kind of function is called a piecewise-defined function.

Piecewise-Defined Functions

Sometimes the domain of the function is given in pieces and the function is described by a different formula in each piece of domain.

This kind of function is called a piecewise-defined function.

Example: Absolute value function

$$|x| = \begin{cases} x, & x \ge 0 \\ -x, & x < 0 \end{cases}$$
 (First formula) (Second formula)

Piecewise-Defined Functions

Sometimes the domain of the function is given in pieces and the function is described by a different formula in each piece of domain.

This kind of function is called a piecewise-defined function.

Example: Absolute value function

$$|x| = \begin{cases} x, & x \ge 0 \\ -x, & x < 0 \end{cases}$$
 (First formula)

y

(Second formula)




Figure 6: The absolute value function has domain (∞,∞) and range $[0,\infty)$.

Piecewise-Defined Functions

Example:

 $f(x) = \lfloor x \rfloor$ —greatest integer function whose values at a given x is the greatest integer $\leq x$.

Piecewise-Defined Functions

Example:

 $f(x) = \lfloor x \rfloor$ —greatest integer function whose values at a given x is the greatest integer $\leq x$.




Figure 7: The graph of $y = \lfloor x \rfloor$ lies on or below the line y = x.

Piecewise-Defined Functions

Example:

 $f(x) = \lfloor x \rfloor$ —greatest integer function whose values at a given x is the greatest integer $\leq x$.




Figure 7: The graph of $y = \lfloor x \rfloor$ lies on or below the line y = x.

$$\lfloor 2.3 \lfloor = 2, \lfloor 1.7 \rfloor = 1, \lfloor 0 \rfloor = 0, \ \lfloor 5 \rfloor = 5, \lfloor -0.3 \rfloor = -1, \ \lfloor 0.15 \rfloor = 0, \lfloor -4 \rfloor = -4,$$

Increasing and Decreasing Functions

If the graph of a function climbs or rises as you move from left to right, we say that the function is increasing.

Increasing and Decreasing Functions

If the graph of a function climbs or rises as you move from left to right, we say that the function is increasing.

If the graph descends or falls as you move from left to right, the function is said to be decreasing.

If the graph of a function climbs or rises as you move from left to right, we say that the function is increasing.

If the graph descends or falls as you move from left to right, the function is said to be decreasing.

Formally,

Definition

Let f be a function defined on an interval I and let x_1 and x_2 be any two points in I.

- 1. If $f(x_2) > f(x_1)$ whenever $x_2 > x_1$, then f is said to be increasing on I.
- 2. If $f(x_2) < f(x_1)$ whenever $x_2 > x_1$, then f is said to be decreasing on I.

Increasing and Decreasing Functions

Example:

$$f(x) = \begin{cases} -x, & x < 0 \\ x^2, & 0 \le x \le 1 \\ 1, & x > 1 \end{cases}$$

Example:

$$f(x) = \begin{cases} -x, & x < 0 \\ x^2, & 0 \le x \le 1 \\ 1, & x > 1 \end{cases}$$

f(x) is decreasing on $[\infty,0)$, increasing on [0,1]. The function neither increasing nor decreasing (constant) on $[1,\infty)$.

Example:

$$f(x) = \begin{cases} -x, & x < 0 \\ x^2, & 0 \le x \le 1 \\ 1, & x > 1 \end{cases}$$

f(x) is decreasing on $[\infty,0)$, increasing on [0,1]. The function neither increasing nor decreasing (constant) on $[1,\infty)$.




Figure 8: Piecewise-defined function increasing, decreasing and stays constant.

Example:

$$f(x) = \begin{cases} -x, & x < 0 \\ x^2, & 0 \le x \le 1 \\ 1, & x > 1 \end{cases}$$

f(x) is decreasing on $[\infty,0)$, increasing on [0,1]. The function neither increasing nor decreasing (constant) on $[1,\infty)$.




Figure 8: Piecewise-defined function increasing, decreasing and stays constant.

Definition

A function y = f(x) is said to be an

- 1. even function of x if f(-x) = f(x),
- 2. odd function of x if f(-x) = -f(x),

for every \boldsymbol{x} in the function's domain.

Definition

A function y = f(x) is said to be an

- 1. even function of x if f(-x) = f(x),
- 2. odd function of x if f(-x) = -f(x),

for every x in the function's domain.

Note that the definitions imply that both x and -x must be in the domain of f.

Definition

A function y = f(x) is said to be an

- 1. even function of x if f(-x) = f(x),
- 2. odd function of x if f(-x) = -f(x),

for every x in the function's domain.

Note that the definitions imply that both x and -x must be in the domain of f.

Example:

If y is an even power of x, as in $y = x^2$ or $y = x^4$, it is an even function of x because $(-x)^2 = x^2$ and $(-x)^4 = x^4$.

Definition

A function y = f(x) is said to be an

- 1. even function of x if f(-x) = f(x),
- 2. odd function of x if f(-x) = -f(x),

for every x in the function's domain.

Note that the definitions imply that both x and -x must be in the domain of f.

Example:

If y is an even power of x, as in $y = x^2$ or $y = x^4$, it is an even function of x because $(-x)^2 = x^2$ and $(-x)^4 = x^4$.

If y is an odd power of x, as in y = x or $y = x^3$, it is an odd function of x because $(-x)^1 = -x^1$ and $(-x)^3 = -x^3$.



Symmetry: Even and Odd Functions

The graph of an even function is symmetric about the y-axis.

Symmetry: Even and Odd Functions

The graph of an even function is symmetric about the y-axis.

Since f(-x) = f(x), a point (x, y) lies on the graph of f if and only if the point (-x, y) lies on the graph.

The graph of an even function is symmetric about the y-axis.

Since f(-x) = f(x), a point (x, y) lies on the graph of f if and only if the point (-x, y) lies on the graph.




Figure 9: The graph of $y=x^2$ (an even function) is symmetric about the y-axis.

Symmetry: Even and Odd Functions

The graph of an odd function is symmetric about the the origin.

Symmetry: Even and Odd Functions

The graph of an odd function is symmetric about the the origin.

Since f(-x) = -f(x), a point (x, y) lies on the graph of f if and only if the point (-x, -y) lies on the graph.

The graph of an odd function is symmetric about the the origin.

Since f(-x) = -f(x), a point (x, y) lies on the graph of f if and only if the point (-x, -y) lies on the graph.




Figure 10: The graph of $y = x^3$ (an odd function) is symmetric about the origin.

Composite Functions

Definition

If f and g are functions, the composite function $f \circ g$ (" f composed with g") is defined by $(f \circ g)(x) = f(g(x))$. The domain of $f \circ g$ consists of the numbers x in the domain of g for which g(x) lies in the domain of f:

$$D_{f \circ g} := \{x: \ x \in D_g \ \text{and} \ g(x) \in D_f\}.$$

Composite Functions

Definition

If f and g are functions, the composite function $f \circ g$ (" f composed with g") is defined by $(f \circ g)(x) = f(g(x))$. The domain of $f \circ g$ consists of the numbers x in the domain of g for which g(x) lies in the domain of f:

$$D_{f \circ g} := \{x : x \in D_g \text{ and } g(x) \in D_f\}.$$




Figure 11: Arrow diagram for $f \circ g$

Composite Functions

EXAMPLE If
$$f(x) = \sqrt{x}$$
 and $g(x) = x + 1$, find

- (a) $(f \circ g)(x)$ (b) $(g \circ f)(x)$ (c) $(f \circ f)(x)$ (d) $(g \circ g)(x)$.

Composite Functions

EXAMPLE If
$$f(x) = \sqrt{x}$$
 and $g(x) = x + 1$, find

(a)
$$(f \circ g)(x)$$
 (b) $(g \circ f)(x)$ (c) $(f \circ f)(x)$ (d) $(g \circ g)(x)$.

Solution

	Composite	Domain
(a)	$(f \circ g)(x) = f(g(x)) = \sqrt{g(x)} = \sqrt{x+1}$	$[-1,\infty)$
(b)	$(g \circ f)(x) = g(f(x)) = f(x) + 1 = \sqrt{x} + 1$	$[0, \infty)$
(c)	$(f \circ f)(x) = f(f(x)) = \sqrt{f(x)} = \sqrt{\sqrt{x}} = x^{1/4}$	$[0,\infty)$
(d)	$(g \circ g)(x) = g(g(x)) = g(x) + 1 = (x + 1) + 1 = x + 2$	$(-\infty, \infty)$

Trigonometric Functions

Angles: angles are measured in degrees or radians.

Trigonometric Functions

Angles: angles are measured in degrees or radians.

The number of radians in the central angle A'CB' within a circle of radius r is defined as the number of "radius units" contained in the arc s subtended by that central angle.

Trigonometric Functions

Angles: angles are measured in degrees or radians.

The number of radians in the central angle A'CB' within a circle of radius r is defined as the number of "radius units" contained in the arc s subtended by that central angle.

In other words, one radian is a measure of angle for which the arc length \boldsymbol{s} is equal to the radius \boldsymbol{r} :

Trigonometric Functions

Angles: angles are measured in degrees or radians.

The number of radians in the central angle A'CB' within a circle of radius r is defined as the number of "radius units" contained in the arc s subtended by that central angle.

In other words, one radian is a measure of angle for which the arc length \boldsymbol{s} is equal to the radius \boldsymbol{r} :

$$\theta = s/r \text{ or } s = \theta r$$



FIGURE The radian measure of the central angle A'CB' is the number $\theta = s/r$. For a unit circle of radius r = 1, θ is the length of arc AB that central angle ACB cuts from the unit circle.

Trigonometric Functions

Angles: angles are measured in degrees or radians.

The number of radians in the central angle A'CB' within a circle of radius r is defined as the number of "radius units" contained in the arc s subtended by that central angle.

In other words, one radian is a measure of angle for which the arc length \boldsymbol{s} is equal to the radius \boldsymbol{r} :

$$\theta = s/r \text{ or } s = \theta r$$



FIGURE The radian measure of the central angle A'CB' is the number $\theta = s/r$. For a unit circle of radius r = 1, θ is the length of arc AB that central angle ACB cuts from the unit circle.

Trigonometric Functions

Since one complete revolution of the unit circle is 360° or 2π radians, we have

$$\pi$$
 radians = 180°

Trigonometric Functions

Since one complete revolution of the unit circle is 360° or 2π radians, we have

$$\pi$$
 radians = 180°


1 radian = $180/\pi (\approx 57.3)$ degrees or 1 degree = $\pi/180 (\approx 0.017)$ radians.

Trigonometric Functions

Since one complete revolution of the unit circle is 360° or 2π radians, we have

$$\pi$$
 radians = 180°

1 radian = $180/\pi (pprox 57.3)$ degrees or 1 degree = $\pi/180 (pprox 0.017)$ radians.



Trigonometric Functions

DEFINITION A function f(x) is **periodic** if there is a positive number p such that f(x + p) = f(x) for every value of x. The smallest such value of p is the **period** of f.

Trigonometric Functions

DEFINITION A function f(x) is **periodic** if there is a positive number p such that f(x + p) = f(x) for every value of x. The smallest such value of p is the **period** of f.

Periods of Trigonometric Functions

Period
$$\pi$$
: $\tan(x + \pi) = \tan x$

$$\cot(x + \pi) = \cot x$$

Period 2
$$\pi$$
: $\sin(x + 2\pi) = \sin x$

$$cos(x + 2\pi) = cos x$$

$$sec(x + 2\pi) = sec x$$

$$\csc(x + 2\pi) = \csc x$$

Trigonometric Functions

Even and odd trigonometric functions:

Trigonometric Functions

Even and odd trigonometric functions:

Even

$$\cos(-x) = \cos x$$

$$\sec(-x) = \sec x$$

Odd

$$\sin(-x) = -\sin x$$


$$\tan(-x) = -\tan x$$

$$\csc(-x) = -\csc x$$

$$\cot(-x) = -\cot x$$

Trigonometric Functions


Graphs of six basic trigonometric functions:



Domain: $-\infty < x < \infty$ Range: $-1 \le y \le 1$

Period: 2π


(a)



Domain: $x \neq \pm \frac{\pi}{2}, \pm \frac{3\pi}{2}, \dots$


Range: $y \le -1$ or $y \ge 1$ Period: 2π

(d)



Domain: $-\infty < x < \infty$ Range: $-1 \le y \le 1$


Period: 2π (b)



Domain: $x \neq 0, \pm \pi, \pm 2\pi, \dots$ Range: $y \le -1$ or $y \ge 1$


Period: 2π

(e)



Domain: $x \neq \pm \frac{\pi}{2}, \pm \frac{3\pi}{2}, \dots$

Range: $-\infty < y < \infty$ Period: (c)



Domain: $x \neq 0, \pm \pi, \pm 2\pi, \dots$ Range: $-\infty < v < \infty$

Period:

(f)

Trigonometric Identities

$$\cos^2\theta+\sin^2\theta=1.$$

Trigonometric Identities

$$\cos^2\theta+\sin^2\theta=1.$$

$$1+\tan^2 u=\sec^2 u$$

$$1 + \cot^2 u = \csc^2 u$$

Trigonometric Identities

$$\cos^2\theta+\sin^2\theta=1.$$

$$1 + \tan^2 u = \sec^2 u$$
$$1 + \cot^2 u = \csc^2 u$$

Addition Formulas:

$$cos(A + B) = cos A cos B - sin A sin B$$

$$sin(A + B) = sin A cos B + cos A sin B$$

Double-Angle Formulas

$$\cos 2u = \cos^2 u - \sin^2 u$$
$$\sin 2u = 2 \sin u \cos u$$



Trigonometric Identities

Half-Angle Formulas

$$\cos^2 u = \frac{1 + \cos 2u}{2}$$
$$\sin^2 u = \frac{1 - \cos 2u}{2}$$