

Differentiation Rules

Nijat Aliyev

BHOS

Calculus

October 11, 2023

Differentiation Rules

Calculate the following limit

$$\lim_{x \rightarrow 0} \frac{\sqrt{3x+1} - 1}{x}$$

Differentiation Rules

2. The derivative of power functions.

Differentiation Rules

2. The derivative of power functions.

If $f(x) = x$, then

$$f'(x) = \frac{df}{dx} = \frac{d}{dx}(x) = 1$$

Differentiation Rules

2. The derivative of power functions.

If $f(x) = x$, then

$$f'(x) = \frac{df}{dx} = \frac{d}{dx}(x) = 1$$

Proof:

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{x+h-x}{h} = \lim_{h \rightarrow 0} \frac{h}{h} = 1.$$

Differentiation Rules

3. Derivative of a Positive Integer Power.

Differentiation Rules

3. Derivative of a Positive Integer Power.

If $f(x) = x^n$, and n is any positive integer, then

$$\frac{df}{dx} = \frac{d}{dx}(x^n) = nx^{n-1}.$$

Differentiation Rules

3. Derivative of a Positive Integer Power.

If $f(x) = x^n$, and n is any positive integer, then

$$\frac{df}{dx} = \frac{d}{dx}(x^n) = nx^{n-1}.$$

Proof: Note that,

$$z^n - x^n = (z - x)(z^{n-1} + z^{n-2}x + \cdots + zx^{n-2} + x^{n-1})$$

So,

$$\begin{aligned} f'(x) &= \lim_{z \rightarrow x} \frac{f(z) - f(x)}{z - x} = \lim_{z \rightarrow x} \frac{z^n - x^n}{z - x} \\ &= \lim_{z \rightarrow x} \frac{(z - x)(z^{n-1} + z^{n-2}x + \cdots + zx^{n-2} + x^{n-1})}{z - x} \\ &= \lim_{z \rightarrow x} (z^{n-1} + z^{n-2}x + \cdots + zx^{n-2} + x^{n-1}) = nx^{n-1}. \end{aligned}$$

Differentiation Rules

4. Power Rule (General Version).

Differentiation Rules

4. Power Rule (General Version).

If n is any real number, then

$$\frac{d}{dx}x^n = nx^{n-1},$$

for all x where the powers x^n and x^{n-1} are defined.

Differentiation Rules

4. Power Rule (General Version).

If n is any real number, then

$$\frac{d}{dx}x^n = nx^{n-1},$$

for all x where the powers x^n and x^{n-1} are defined.

Example: Find the derivative of the following functions

Differentiation Rules

4. Power Rule (General Version).

If n is any real number, then

$$\frac{d}{dx}x^n = nx^{n-1},$$

for all x where the powers x^n and x^{n-1} are defined.

Example: Find the derivative of the following functions

(a) x^3 (b) $x^{2/3}$ (c) $x^{\sqrt{2}}$ (d) $\frac{1}{x^4}$ (e) $x^{-4/3}$ (f) $\sqrt{x^{2+\pi}}$

Differentiation Rules

Solutions:

$$(a) \frac{d}{dx}(x^3) = 3x^{3-1} = 3x^2$$

$$(b) \frac{d}{dx}(x^{2/3}) = \frac{2}{3}x^{(2/3)-1} = \frac{2}{3}x^{-1/3}$$

$$(c) \frac{d}{dx}(x^{\sqrt{2}}) = \sqrt{2}x^{\sqrt{2}-1}$$

$$(d) \frac{d}{dx}\left(\frac{1}{x^4}\right) = \frac{d}{dx}(x^{-4}) = -4x^{-4-1} = -4x^{-5} = -\frac{4}{x^5}$$

$$(e) \frac{d}{dx}(x^{-4/3}) = -\frac{4}{3}x^{-(4/3)-1} = -\frac{4}{3}x^{-7/3}$$

$$(f) \frac{d}{dx}(\sqrt{x^{2+\pi}}) = \frac{d}{dx}(x^{1+(\pi/2)}) = \left(1 + \frac{\pi}{2}\right)x^{1+(\pi/2)-1} = \frac{1}{2}(2 + \pi)\sqrt{x^\pi}$$

Differentiation Rules

4. The Constant Multiple Rule

Differentiation Rules

4. The Constant Multiple Rule

If c is a constant and f is a differentiable function, then

$$\frac{d}{dx} [cf(x)] = c \frac{d}{dx} f(x)$$

Differentiation Rules

4. The Constant Multiple Rule

If c is a constant and f is a differentiable function, then

$$\frac{d}{dx} [cf(x)] = c \frac{d}{dx} f(x)$$

Proof:

Let $g(x) = cf(x)$. Then

$$\begin{aligned} g'(x) &= \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} = \lim_{h \rightarrow 0} \frac{cf(x+h) - cf(x)}{h} \\ &= \lim_{h \rightarrow 0} c \left[\frac{f(x+h) - f(x)}{h} \right] \\ &= c \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\ &= cf'(x) \end{aligned}$$

Differentiation Rules

Example:

Differentiation Rules

Example:

$$(a) \frac{d}{dx} (3x^4) = 3 \frac{d}{dx} (x^4) = 3(4x^3) = 12x^3$$

$$(b) \frac{d}{dx} (-x) = \frac{d}{dx} [(-1)x] = (-1) \frac{d}{dx} (x) = -1(1) = -1$$

Differentiation Rules

5. The Sum Rule

Differentiation Rules

5. The Sum Rule

If f and g are both differentiable, then

$$\frac{d}{dx}[f(x) + g(x)] = \frac{d}{dx}f(x) + \frac{d}{dx}g(x)$$

Differentiation Rules

5. The Sum Rule

If f and g are both differentiable, then

$$\frac{d}{dx} [f(x) + g(x)] = \frac{d}{dx} f(x) + \frac{d}{dx} g(x)$$

Proof:

Let $F(x) = f(x) + g(x)$. Then

$$\begin{aligned} F'(x) &= \lim_{h \rightarrow 0} \frac{F(x+h) - F(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{[f(x+h) + g(x+h)] - [f(x) + g(x)]}{h} \\ &= \lim_{h \rightarrow 0} \left[\frac{f(x+h) - f(x)}{h} + \frac{g(x+h) - g(x)}{h} \right] \\ &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} + \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &= f'(x) + g'(x) \end{aligned}$$

Differentiation Rules

The Sum Rule can be extended to the sum of any number of functions.

Differentiation Rules

The Sum Rule can be extended to the sum of any number of functions.

$$(f + g + h)' = [(f + g) + h]' = (f + g)' + h' = f' + g' + h'$$

Differentiation Rules

The Sum Rule can be extended to the sum of any number of functions.

$$(f + g + h)' = [(f + g) + h]' = (f + g)' + h' = f' + g' + h'$$

6. The Difference Rule

If f and g are both differentiable, then

$$\frac{d}{dx} [f(x) - g(x)] = \frac{d}{dx} f(x) - \frac{d}{dx} g(x)$$

Differentiation Rules

The Sum Rule can be extended to the sum of any number of functions.

$$(f + g + h)' = [(f + g) + h]' = (f + g)' + h' = f' + g' + h'$$

6. The Difference Rule

If f and g are both differentiable, then

$$\frac{d}{dx} [f(x) - g(x)] = \frac{d}{dx} f(x) - \frac{d}{dx} g(x)$$

Example:

$$\begin{aligned} & \frac{d}{dx} (x^8 + 12x^5 - 4x^4 + 10x^3 - 6x + 5) \\ &= \frac{d}{dx} (x^8) + 12 \frac{d}{dx} (x^5) - 4 \frac{d}{dx} (x^4) + 10 \frac{d}{dx} (x^3) - 6 \frac{d}{dx} (x) + \frac{d}{dx} (5) \\ &= 8x^7 + 12(5x^4) - 4(4x^3) + 10(3x^2) - 6(1) + 0 \\ &= 8x^7 + 60x^4 - 16x^3 + 30x^2 - 6 \end{aligned}$$

Differentiation Rules

Example: Find the points on the curve $y = x^4 - 6x^2 + 4$ where the tangent line is horizontal.

Differentiation Rules

Example: Find the points on the curve $y = x^4 - 6x^2 + 4$ where the tangent line is horizontal.

Solution:

Horizontal tangents occur where the derivative is zero. We have

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(x^4) - 6 \frac{d}{dx}(x^2) + \frac{d}{dx}(4) \\ &= 4x^3 - 12x + 0 = 4x(x^2 - 3)\end{aligned}$$

Differentiation Rules

Example: Find the points on the curve $y = x^4 - 6x^2 + 4$ where the tangent line is horizontal.

Solution:

Horizontal tangents occur where the derivative is zero. We have

$$\begin{aligned}\frac{dy}{dx} &= \frac{d}{dx}(x^4) - 6 \frac{d}{dx}(x^2) + \frac{d}{dx}(4) \\ &= 4x^3 - 12x + 0 = 4x(x^2 - 3)\end{aligned}$$


So the curve has horizontal tangents when $x = 0, x = \sqrt{3}, -\sqrt{3}$

Differentiation Rules

The corresponding points are $(0, 4), (\sqrt{3}, -5), (-\sqrt{3}, -5))$

Differentiation Rules

The corresponding points are $(0, 4)$, $(\sqrt{3}, -5)$, $(-\sqrt{3}, -5)$



Differentiation Rules

7. Product Rule If f and g are differentiable, then

$$\frac{d}{dx}[f(x)g(x)] = f(x)\frac{d}{dx}g(x) + g(x)\frac{d}{dx}f(x)$$

Differentiation Rules

7. Product Rule If f and g are differentiable, then

$$\frac{d}{dx}[f(x)g(x)] = f(x)\frac{d}{dx}g(x) + g(x)\frac{d}{dx}f(x)$$

Proof:

$$\begin{aligned}\frac{d}{dx}[f(x)g(x)] &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(x+h)g(x+h) - f(x)g(x+h) + f(x)g(x+h) - f(x)g(x)}{h} \\ &= \lim_{h \rightarrow 0} \frac{g(x+h)[f(x+h) - f(x)] + f(x)[g(x+h) - g(x)]}{h} \\ &= \lim_{h \rightarrow 0} g(x+h) \cdot \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} + f(x) \cdot \lim_{h \rightarrow 0} \frac{g(x+h) - g(x)}{h} \\ &= g(x)f'(x) + f(x)g'(x)\end{aligned}$$

Example:

- (a) If $f(x) = xe^x$, find $f'(x)$.
- (b) Find the n th derivative, $f^{(n)}(x)$.

Example:

- (a) If $f(x) = xe^x$, find $f'(x)$.
- (b) Find the n th derivative, $f^{(n)}(x)$.

Solution:

(a) By the Product Rule, we have

$$\begin{aligned}f'(x) &= \frac{d}{dx}(xe^x) \\&= x \frac{d}{dx}(e^x) + e^x \frac{d}{dx}(x) \\&= xe^x + e^x \cdot 1 = (x + 1)e^x\end{aligned}$$

(b) Using the Product Rule a second time, we get

$$\begin{aligned}
 f''(x) &= \frac{d}{dx} [(x+1)e^x] \\
 &= (x+1) \frac{d}{dx} (e^x) + e^x \frac{d}{dx} (x+1) \\
 &= (x+1)e^x + e^x \cdot 1 = (x+2)e^x
 \end{aligned}$$

Further applications of the Product Rule give

$$f'''(x) = (x+3)e^x \quad f^{(4)}(x) = (x+4)e^x$$

In fact, each successive differentiation adds another term e^x , so

$$f^{(n)}(x) = (x+n)e^x$$

Differentiation Rules

8. Quotient Rule: If f and g are differentiable, then

$$\frac{d}{dx} \left[\frac{f(x)}{g(x)} \right] = \frac{g(x) \frac{d}{dx} f(x) - f(x) \frac{d}{dx} g(x)}{g^2(x)}$$

provided that $g(x) \neq 0$.

Example:

$$\text{Let } y = \frac{x^2 + x - 2}{x^3 + 6}. \text{ Then}$$

Example:

$$\text{Let } y = \frac{x^2 + x - 2}{x^3 + 6}. \text{ Then}$$

$$\begin{aligned} y' &= \frac{(x^3 + 6) \frac{d}{dx}(x^2 + x - 2) - (x^2 + x - 2) \frac{d}{dx}(x^3 + 6)}{(x^3 + 6)^2} \\ &= \frac{(x^3 + 6)(2x + 1) - (x^2 + x - 2)(3x^2)}{(x^3 + 6)^2} \\ &= \frac{(2x^4 + x^3 + 12x + 6) - (3x^4 + 3x^3 - 6x^2)}{(x^3 + 6)^2} \\ &= \frac{-x^4 - 2x^3 + 6x^2 + 12x + 6}{(x^3 + 6)^2} \end{aligned}$$

Example: Find an equation of the tangent line to the curve $y = \frac{e^x}{1+x^2}$.

Example: Find an equation of the tangent line to the curve $y = \frac{e^x}{1+x^2}$.

According to the Quotient Rule, we have

$$\begin{aligned}\frac{dy}{dx} &= \frac{(1+x^2) \frac{d}{dx}(e^x) - e^x \frac{d}{dx}(1+x^2)}{(1+x^2)^2} \\ &= \frac{(1+x^2)e^x - e^x(2x)}{(1+x^2)^2} \\ &= \frac{e^x(1-x)^2}{(1+x^2)^2}\end{aligned}$$

Example: Find an equation of the tangent line to the curve $y = \frac{e^x}{1+x^2}$.

According to the Quotient Rule, we have

$$\begin{aligned}\frac{dy}{dx} &= \frac{(1+x^2) \frac{d}{dx}(e^x) - e^x \frac{d}{dx}(1+x^2)}{(1+x^2)^2} \\ &= \frac{(1+x^2)e^x - e^x(2x)}{(1+x^2)^2} \\ &= \frac{e^x(1-x)^2}{(1+x^2)^2}\end{aligned}$$

So the slope of the tangent line at $(1, \frac{1}{2}e)$ is

$$\left. \frac{dy}{dx} \right|_{x=1} = 0$$

This means that the tangent line at $(1, \frac{1}{2}e)$ is horizontal and its equation is $y = \frac{1}{2}e$.

Table of Differentiation Rules

$$\frac{d}{dx}(c) = 0$$

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

$$\frac{d}{dx}(e^x) = e^x$$

$$(cf)' = cf'$$

$$(f + g)' = f' + g'$$

$$(f - g)' = f' - g'$$

$$(fg)' = fg' + gf'$$

$$\left(\frac{f}{g}\right)' = \frac{gf' - fg'}{g^2}$$