

Continuity

Nijat Aliyev

BHOS

Calculus

September 27, 2023

Definition

A function $y = f(x)$ is continuous at $x = a$ if it satisfies

$$\lim_{x \rightarrow a} f(x) = f(a)$$

Definition

A function $y = f(x)$ is continuous at $x = a$ if it satisfies

$$\lim_{x \rightarrow a} f(x) = f(a)$$

$f(x)$ is continuous at $x = a$ if as x approaches a , $f(x)$ approaches $f(a)$.


Definition

A function $y = f(x)$ is continuous at $x = a$ if it satisfies

$$\lim_{x \rightarrow a} f(x) = f(a)$$

$f(x)$ is continuous at $x = a$ if as x approaches a , $f(x)$ approaches $f(a)$.

Geometrically, if the graph of a function $y = f(x)$ can be sketched at $x = a$ in one unbroken motion, then f is a continuous function.



Continuity

A continuous function has a property that a small change in (input) x produces only a small change in $f(x)$

Continuity

A continuous function has a property that a small change in (input) x produces only a small change in $f(x)$

In fact, the change in $f(x)$ can be kept as small as we want by keeping the change in x sufficiently small.

Continuity

A continuous function has a property that a small change in (input) x produces only a small change in $f(x)$

In fact, the change in $f(x)$ can be kept as small as we want by keeping the change in x sufficiently small.

If a function f is not continuous at a point a in its domain then we say that f is discontinuous at $x = a$.

To be continuous, f must satisfy the following conditions:

1. $f(a)$ is defined (that is, a is in the domain of f)
2. $\lim_{x \rightarrow a} f(x)$ exists
3. $\lim_{x \rightarrow a} f(x) = f(a)$

Recall: $\lim_{x \rightarrow a} f(x)$ exist if and only if $\lim_{x \rightarrow a^-} f(x)$ and $\lim_{x \rightarrow a^+} f(x)$ both exist and are equal.

To be continuous, f must satisfy the following conditions:

1. $f(a)$ is defined (that is, a is in the domain of f)
2. $\lim_{x \rightarrow a} f(x)$ exists
3. $\lim_{x \rightarrow a} f(x) = f(a)$

Recall: $\lim_{x \rightarrow a} f(x)$ exist if and only if $\lim_{x \rightarrow a^-} f(x)$ and $\lim_{x \rightarrow a^+} f(x)$ both exist and are equal.

Example: Let $f(x) = \frac{x+1}{x-1}$. At which point is f discontinuous?

To be continuous, f must satisfy the following conditions:

1. $f(a)$ is defined (that is, a is in the domain of f)
2. $\lim_{x \rightarrow a} f(x)$ exists
3. $\lim_{x \rightarrow a} f(x) = f(a)$

Recall: $\lim_{x \rightarrow a} f(x)$ exist if and only if $\lim_{x \rightarrow a^-} f(x)$ and $\lim_{x \rightarrow a^+} f(x)$ both exist and are equal.

Example: Let $f(x) = \frac{x+1}{x-1}$. At which point is f discontinuous?

Example: Let $f(x) = x^2$. Is f continuous at $x = 1$?

To be continuous, f must satisfy the following conditions:

1. $f(a)$ is defined (that is, a is in the domain of f)
2. $\lim_{x \rightarrow a} f(x)$ exists
3. $\lim_{x \rightarrow a} f(x) = f(a)$

Recall: $\lim_{x \rightarrow a} f(x)$ exist if and only if $\lim_{x \rightarrow a^-} f(x)$ and $\lim_{x \rightarrow a^+} f(x)$ both exist and are equal.

Example: Let $f(x) = \frac{x+1}{x-1}$. At which point is f discontinuous?

Example: Let $f(x) = x^2$. Is f continuous at $x = 1$?

1. $f(1)$ is defined.

To be continuous, f must satisfy the following conditions:

1. $f(a)$ is defined (that is, a is in the domain of f)
2. $\lim_{x \rightarrow a} f(x)$ exists
3. $\lim_{x \rightarrow a} f(x) = f(a)$

Recall: $\lim_{x \rightarrow a} f(x)$ exist if and only if $\lim_{x \rightarrow a^-} f(x)$ and $\lim_{x \rightarrow a^+} f(x)$ both exist and are equal.

Example: Let $f(x) = \frac{x+1}{x-1}$. At which point is f discontinuous?

Example: Let $f(x) = x^2$. Is f continuous at $x = 1$?

1. $f(1)$ is defined.
2. $\lim_{x \rightarrow 1} f(x)$ exists.

To be continuous, f must satisfy the following conditions:

1. $f(a)$ is defined (that is, a is in the domain of f)
2. $\lim_{x \rightarrow a} f(x)$ exists
3. $\lim_{x \rightarrow a} f(x) = f(a)$

Recall: $\lim_{x \rightarrow a} f(x)$ exist if and only if $\lim_{x \rightarrow a^-} f(x)$ and $\lim_{x \rightarrow a^+} f(x)$ both exist and are equal.


Example: Let $f(x) = \frac{x+1}{x-1}$. At which point is f discontinuous?

Example: Let $f(x) = x^2$. Is f continuous at $x = 1$?

1. $f(1)$ is defined.
2. $\lim_{x \rightarrow 1} f(x)$ exists.
3. $\lim_{x \rightarrow 1} f(x) = f(1)$.

Continuity


Example: The graph of a function f is given below. At which points is f discontinuous?



Solution:

1. There is a break at $x = 1$. The reason is that $f(1)$ is not defined.


Example: The graph of a function f is given below. At which points is f discontinuous?



Solution:

1. There is a break at $x = 1$. The reason is that $f(1)$ is not defined.
2. There is a break at $x = 3$. $f(3)$ is defined.


Example: The graph of a function f is given below. At which points is f discontinuous?



Solution:

1. There is a break at $x = 1$. The reason is that $f(1)$ is not defined.
2. There is a break at $x = 3$. $f(3)$ is defined. But $\lim_{x \rightarrow 3}$ does not exist.

Example: The graph of a function f is given below. At which points is f discontinuous?



Solution:

1. There is a break at $x = 1$. The reason is that $f(1)$ is not defined.
2. There is a break at $x = 3$. $f(3)$ is defined. But $\lim_{x \rightarrow 3}$ does not exist. Why?
3. f is not continuous at $x = 5$. $f(5)$ is defined and limit exists. But $\lim_{x \rightarrow 5} \neq f(5)$.

Example: At which points are the following functions discontinuous?

$$(a) f(x) = \frac{x^2 - x - 2}{x - 2}$$

$$(b) f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$

$$(c) f(x) = \begin{cases} \frac{x^2 - x - 2}{x - 2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$

$$(d) f(x) = \llbracket x \rrbracket$$

Solution:

(a) $f(2)$ is not defined. So f is discontinuous at $x = 2$.

Example: At which points are the following functions discontinuous?

$$(a) f(x) = \frac{x^2 - x - 2}{x - 2}$$

$$(b) f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$

$$(c) f(x) = \begin{cases} \frac{x^2 - x - 2}{x - 2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$

$$(d) f(x) = \llbracket x \rrbracket$$

Solution:

(a) $f(2)$ is not defined. So f is discontinuous at $x = 2$.

(b) $f(0) = 1$ is defined but $\lim_{x \rightarrow 0} f(x)$ does not exist.

Example: At which points are the following functions discontinuous?

$$(a) f(x) = \frac{x^2 - x - 2}{x - 2}$$

$$(b) f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$

$$(c) f(x) = \begin{cases} \frac{x^2 - x - 2}{x - 2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$

$$(d) f(x) = \llbracket x \rrbracket$$

Solution:

(a) $f(2)$ is not defined. So f is discontinuous at $x = 2$.

(b) $f(0) = 1$ is defined but $\lim_{x \rightarrow 0} f(x)$ does not exist.

(c) $f(2)$ is defined

$$\lim_{x \rightarrow 2} f(x) = \lim_{x \rightarrow 2} \frac{(x-2)(x+1)}{x-2} = \lim_{x \rightarrow 2} (x+1) = 3$$

exists. But $\lim_{x \rightarrow 2} f(x) \neq f(2)$.

Example: At which points are the following functions discontinuous?

$$(a) f(x) = \frac{x^2 - x - 2}{x - 2}$$

$$(b) f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$

$$(c) f(x) = \begin{cases} \frac{x^2 - x - 2}{x - 2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$

$$(d) f(x) = \llbracket x \rrbracket$$

Solution:

(a) $f(2)$ is not defined. So f is discontinuous at $x = 2$.

(b) $f(0) = 1$ is defined but $\lim_{x \rightarrow 0} f(x)$ does not exist.


(c) $f(2)$ is defined

$$\lim_{x \rightarrow 2} f(x) = \lim_{x \rightarrow 2} \frac{(x-2)(x+1)}{x-2} = \lim_{x \rightarrow 2} (x+1) = 3$$


exists. But $\lim_{x \rightarrow 2} f(x) \neq f(2)$.

(d) $\lim_{x \rightarrow n} \llbracket x \rrbracket$ does not exist for $n \in \mathbb{Z}$


Continuity




$$(a) f(x) = \frac{x^2 - x - 2}{x - 2}$$




$$(b) f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$




$$(c) f(x) = \begin{cases} \frac{x^2 - x - 2}{x - 2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$




$$(d) f(x) = \lfloor x \rfloor$$




$$(a) f(x) = \frac{x^2 - x - 2}{x - 2}$$



$$(b) f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$




$$(c) f(x) = \begin{cases} \frac{x^2 - x - 2}{x - 2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$




$$(d) f(x) = \lfloor x \rfloor$$


The kind of discontinuity in (a) and (c) is called **removable discontinuity**.




$$(a) f(x) = \frac{x^2 - x - 2}{x - 2}$$



$$(b) f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$




$$(c) f(x) = \begin{cases} \frac{x^2 - x - 2}{x - 2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$




$$(d) f(x) = \lfloor x \rfloor$$

The kind of discontinuity in (a) and (c) is called **removable discontinuity**.


The discontinuity in part (b) is called an infinite discontinuity.




$$(a) f(x) = \frac{x^2 - x - 2}{x - 2}$$



$$(b) f(x) = \begin{cases} \frac{1}{x^2} & \text{if } x \neq 0 \\ 1 & \text{if } x = 0 \end{cases}$$



$$(c) f(x) = \begin{cases} \frac{x^2 - x - 2}{x - 2} & \text{if } x \neq 2 \\ 1 & \text{if } x = 2 \end{cases}$$



$$(d) f(x) = \lfloor x \rfloor$$

The kind of discontinuity in (a) and (c) is called **removable discontinuity**.

The discontinuity in part (b) is called an infinite discontinuity.

The discontinuities in part (d) are called jump discontinuities because the function “jumps” from one value to another.

Definition

A function f is called right continuous (or continuous from the right) at a point $x = a$ if

$$\lim_{x \rightarrow a^+} f(x) = f(a)$$

and f is continuous from the left at a if

$$\lim_{x \rightarrow a^-} f(x) = f(a)$$

Definition

A function f is called right continuous (or continuous from the right) at a point $x = a$ if

$$\lim_{x \rightarrow a^+} f(x) = f(a)$$

and f is continuous from the left at a if

$$\lim_{x \rightarrow a^-} f(x) = f(a)$$

Example: At each integer n the function $\lim_{x \rightarrow n} [[x]]$ is continuous from the right but discontinuous from the left because

Definition

A function f is called right continuous (or continuous from the right) at a point $x = a$ if

$$\lim_{x \rightarrow a^+} f(x) = f(a)$$

and f is continuous from the left at a if

$$\lim_{x \rightarrow a^-} f(x) = f(a)$$

Example: At each integer n the function $\lim_{x \rightarrow n} [[x]]$ is continuous from the right but discontinuous from the left because

$$\lim_{x \rightarrow n^+} [[x]] = n = f(n)$$

but

$$\lim_{x \rightarrow n^-} [[x]] = n - 1 = f(n)$$

Definition

A function f is continuous on an interval if it is continuous at every number in the interval.

Definition

A function f is continuous on an interval if it is continuous at every number in the interval.

Definition

A function f is continuous on an interval if it is continuous at every number in the interval.


Remark: If f is defined only on one side of an endpoint of the interval, continuity at the endpoint means continuous from the right or continuous from the left.

Definition


A function f is continuous on an interval if it is continuous at every number in the interval.

Remark: If f is defined only on one side of an endpoint of the interval, continuity at the endpoint means continuous from the right or continuous from the left.

Example: Show that $f(x) = 1 - \sqrt{1 - x^2}$ continuous on the interval $[-1, 1]$.



Example: Show that $f(x) = 1 - \sqrt{1 - x^2}$ continuous on the interval $[-1, 1]$.




Solution:

Using limit laws for $-1 < a < 1$ we have

$$\lim_{x \rightarrow a} f(x) = 1 - \sqrt{1 - x^2} = f(a)$$

Example: Show that $f(x) = 1 - \sqrt{1 - x^2}$ continuous on the interval $[-1, 1]$.



Solution:


Using limit laws for $-1 < a < 1$ we have

$$\lim_{x \rightarrow a} f(x) = 1 - \sqrt{1 - x^2} = f(a)$$

Similarly,

$$\lim_{x \rightarrow -1^+} f(x) = 1 = f(-1)$$

Example: Show that $f(x) = 1 - \sqrt{1 - x^2}$ continuous on the interval $[-1, 1]$.



Solution:

Using limit laws for $-1 < a < 1$ we have

$$\lim_{x \rightarrow a} f(x) = 1 - \sqrt{1 - x^2} = f(a)$$


Similarly,

$$\lim_{x \rightarrow -1^+} f(x) = 1 = f(-1)$$

and

$$\lim_{x \rightarrow 1^-} f(x) = 1 = f(1)$$

Example: Show that $f(x) = 1 - \sqrt{1 - x^2}$ continuous on the interval $[-1, 1]$.



Solution:

Using limit laws for $-1 < a < 1$ we have

$$\lim_{x \rightarrow a} f(x) = 1 - \sqrt{1 - x^2} = f(a)$$

Similarly,

$$\lim_{x \rightarrow -1^+} f(x) = 1 = f(-1)$$

and

$$\lim_{x \rightarrow 1^-} f(x) = 1 = f(1)$$

Theorem

If f, g are continuous at $x = a$ and c is any constant then the following functions are also continuous at $x = a$:

1. $f \pm g$
2. cf
3. fg
4. $\frac{f}{g}$ provided that $g(a) \neq 0$.

Proof: Let us prove only first one. f, g continuous at a means that $\lim_{x \rightarrow a} f(x) = f(a)$ and $\lim_{x \rightarrow a} g(x) = g(a)$. Then,

$$\lim_{x \rightarrow a} (f+g)(x) = \lim_{x \rightarrow a} [f(x)+g(x)] = \lim_{x \rightarrow a} f(x) + \lim_{x \rightarrow a} g(x) = f(a) + g(a) = (f+g)(a)$$

Theorem:

- (a) Any polynomial is continuous everywhere; that is, it is continuous on $\mathbb{R} = (-\infty, \infty)$.
- (b) Any rational function is continuous wherever it is defined; that is, it is continuous on its domain.

Theorem:

- (a) Any polynomial is continuous everywhere; that is, it is continuous on $\mathbb{R} = (-\infty, \infty)$.
- (b) Any rational function is continuous wherever it is defined; that is, it is continuous on its domain.

Example: Find $\lim_{x \rightarrow 2} \frac{x^2+1}{x^3-1}$.

Theorem:

- (a) Any polynomial is continuous everywhere; that is, it is continuous on $\mathbb{R} = (-\infty, \infty)$.
- (b) Any rational function is continuous wherever it is defined; that is, it is continuous on its domain.

Example: Find $\lim_{x \rightarrow 2} \frac{x^2+1}{x^3-1}$.

Solution:

The function $f(x) = \frac{x^2+1}{x^3-1}$ is rational function so it is continuous on its domain.

Theorem:

- (a) Any polynomial is continuous everywhere; that is, it is continuous on $\mathbb{R} = (-\infty, \infty)$.
- (b) Any rational function is continuous wherever it is defined; that is, it is continuous on its domain.

Example: Find $\lim_{x \rightarrow 2} \frac{x^2+1}{x^3-1}$.

Solution:

The function $f(x) = \frac{x^2+1}{x^3-1}$ is rational function so it is continuous on its domain.

$$\lim_{x \rightarrow 2} \frac{x^2+1}{x^3-1} = f(2) = 5/7.$$

Theorem:

The following types of functions are continuous at every number in their domains:

polynomials

rational functions

root functions

trigonometric functions

inverse trigonometric functions

exponential functions

logarithmic functions

Example:

Where is the function $f(x) = \frac{\ln x + \tan^{-1} x}{x^2 - 1}$ continuous?

Example:

Where is the function $f(x) = \frac{\ln x + \tan^{-1}x}{x^2 - 1}$ continuous?

Solution:

$\ln(x)$ is continuous for $x > 0$ and $\tan^{-1}x$ is continuous on \mathbb{R} thus $\ln(x) + \tan^{-1}x$ is also continuous.

Example:

Where is the function $f(x) = \frac{\ln x + \tan^{-1}x}{x^2 - 1}$ continuous?

Solution:

$\ln(x)$ is continuous for $x > 0$ and $\tan^{-1}x$ is continuous on \mathbb{R} thus $\ln(x) + \tan^{-1}x$ is also continuous.

$x^2 - 1$ is a polynomial so continuous.

Example:

Where is the function $f(x) = \frac{\ln x + \tan^{-1}x}{x^2 - 1}$ continuous?

Solution:

$\ln(x)$ is continuous for $x > 0$ and $\tan^{-1}x$ is continuous on \mathbb{R} thus $\ln(x) + \tan^{-1}x$ is also continuous.

$x^2 - 1$ is a polynomial so continuous.

Therefore, $f(x)$ is continuous everywhere except where denominator is zero.

Example:

Where is the function $f(x) = \frac{\ln x + \tan^{-1}x}{x^2 - 1}$ continuous?

Solution:

$\ln(x)$ is continuous for $x > 0$ and $\tan^{-1}x$ is continuous on \mathbb{R} thus $\ln(x) + \tan^{-1}x$ is also continuous.

$x^2 - 1$ is a polynomial so continuous.

Therefore, $f(x)$ is continuous everywhere except where denominator is zero.

So, $f(x)$ is continuous on $(0, \infty) \setminus \{1\} = (0, 1) \cup (1, \infty)$

Theorem

If f is continuous at b and $\lim_{x \rightarrow a} g(x) = b$, then

$$\lim_{x \rightarrow a} f(g(x)) = f(\lim_{x \rightarrow a} g(x)) = f(b)$$

Theorem

If f is continuous at b and $\lim_{x \rightarrow a} g(x) = b$, then

$$\lim_{x \rightarrow a} f(g(x)) = f(\lim_{x \rightarrow a} g(x)) = f(b)$$

Example:

Evaluate $\lim_{x \rightarrow 1} \arcsin\left(\frac{1 - \sqrt{x}}{1 - x}\right)$.

Theorem

If f is continuous at b and $\lim_{x \rightarrow a} g(x) = b$, then

$$\lim_{x \rightarrow a} f(g(x)) = f(\lim_{x \rightarrow a} g(x)) = f(b)$$

Example:

Evaluate $\lim_{x \rightarrow 1} \arcsin\left(\frac{1 - \sqrt{x}}{1 - x}\right)$.

Solution: Since \arcsin is a continuous function, we can apply the above theorem.

Theorem

If f is continuous at b and $\lim_{x \rightarrow a} g(x) = b$, then

$$\lim_{x \rightarrow a} f(g(x)) = f(\lim_{x \rightarrow a} g(x)) = f(b)$$

Example:

Evaluate $\lim_{x \rightarrow 1} \arcsin\left(\frac{1 - \sqrt{x}}{1 - x}\right)$.

Solution: Since \arcsin is a continuous function, we can apply the above theorem.

$$\begin{aligned} \lim_{x \rightarrow 1} \arcsin\left(\frac{1 - \sqrt{x}}{1 - x}\right) &= \arcsin\left(\lim_{x \rightarrow 1} \frac{1 - \sqrt{x}}{1 - x}\right) = \arcsin\left(\lim_{x \rightarrow 1} \frac{1 - \sqrt{x}}{(1 - \sqrt{x})(1 + \sqrt{x})}\right) \\ &= \arcsin\left(\lim_{x \rightarrow 1} \frac{1}{1 + \sqrt{x}}\right) = \arcsin \frac{1}{2} = \frac{\pi}{6} \end{aligned}$$

Theorem

If g is continuous at a and f is continuous at $g(a)$, then the composite function $f \circ g$ defined by $(f \circ g)(x) = f(g(x))$ is continuous at a .

Proof.

Since g is continuous at a , we have

$$\lim_{x \rightarrow a} g(x) = g(a)$$

Since f is continuous at $b = g(a)$ we can apply previous theorem to get

$$\lim_{x \rightarrow a} f(g(x)) = f(\lim_{x \rightarrow a} g(x)) = f(g(a))$$

Thus, we get

$$\lim_{x \rightarrow a} f(g(x)) = f(g(a)).$$

Example:

(a) $f(x) = \cos(x^3 + 2x)$

Example:

(a) $f(x) = \cos(x^3 + 2x)$

(b) $f(x) = \ln(1 + \sin x)$

Example:

(a) $f(x) = \cos(x^3 + 2x)$

(b) $f(x) = \ln(1 + \sin x)$

(c) $f(x) = e^{x^2}$