

# Section 52: Problem 4 Solution <sup>TM</sup> (https://dbfin.com/teachme/) ♪ (https://dbfin.com/search/)

“

Working problems is a crucial part of learning mathematics. No one can learn topology merely by poring over the definitions, theorems, and examples that are worked out in the text. One must work part of it out for oneself. To provide that opportunity is the purpose of the exercises.

— James R. Munkres

Let  $A \subset X$  ; suppose  $r : X \rightarrow A$  is a continuous map such that  $r(a) = a$  for each  $a \in A$  . (The map  $r$  is called a **retraction** of  $X$  onto  $A$  .) If  $a_0 \in A$  , show that

$$r_* : \pi_1(X, a_0) \rightarrow \pi_1(A, a_0)$$

is surjective.

By definition,  $r_*([f]) = [r \circ f]$  , so if  $[f] \in \pi_1(A, a_0)$  , i.e.  $f$  is a loop in  $A$  based at  $a_0$  , then  $r \circ f = f$  , and  $r_*([f]) = [f]$  .

0 Comments

dbFin

 Login

 Recommend

 Tweet

 Share

Sort by Oldest ▾




LOG IN WITH

OR SIGN UP WITH DISQUS 


Be the first to comment.

 Subscribe
  Add Disqus to your site
 Add DisqusAdd
  Disqus' Privacy Policy
 Privacy PolicyPrivacy PolicyPrivacy Policy

Copyright © 2007 - 2019 dbFin

 Powered by <sup>\*</sup>GetSimple (http://get-simple.info/)