



✓ **Congratulations! You passed!**

TO PASS 80% or higher

Keep Learning

GRADE
100%

Linear Regression with One Variable

LATEST SUBMISSION GRADE

100%

1. Consider the problem of predicting how well a student does in her second year of college/university, given how well she did in her first year.

1 / 1 point

Specifically, let x be equal to the number of "A" grades (including A-, A and A+ grades) that a student receives in their first year of college (freshmen year). We would like to predict the value of y , which we define as the number of "A" grades they get in their second year (sophomore year).

Here each row is one training example. Recall that in linear regression, our hypothesis is $h_{\theta}(x) = \theta_0 + \theta_1 x$, and we use m to denote the number of training examples.

x	y
5	4
3	4
0	1
4	3

For the training set given above (note that this training set may also be referenced in other questions in this quiz), what is the value of m ? In the box below, please enter your answer (which should be a number between 0 and 10).

✓ **Correct**

2. Consider the following training set of $m = 4$ training examples:

1 / 1 point

x	y
1	0.5
2	1
4	2
0	0

Consider the linear regression model $h_{\theta}(x) = \theta_0 + \theta_1 x$. What are the values of θ_0 and θ_1 that you would expect to obtain upon running gradient descent on this model? (Linear regression will be able to fit this data perfectly.)

✓ **Correct**

3. Suppose we set $\theta_0 = -1$, $\theta_1 = 0.5$. What is $h_{\theta}(4)$?

1 / 1 point

✓ **Correct**

4. Let f be some function so that

1 / 1 point

$f(\theta_0, \theta_1)$ outputs a number. For this problem,

f is some arbitrary/unknown smooth function (not necessarily the cost function of linear regression, so f may have local optima).

Suppose we use gradient descent to try to minimize $f(\theta_0, \theta_1)$

as a function of θ_0 and θ_1 . Which of the

following statements are true? (Check all that apply.)

✓ **Correct**

5. Suppose that for some linear regression problem (say, predicting housing prices as in the lecture), we have some training set, and for our training set we managed to find some θ_0, θ_1 such that $J(\theta_0, \theta_1) = 0$.

1 / 1 point

Which of the statements below must then be true? (Check all that apply.)

✓ Correct