1/1 point

1/1 point

Linear Algebra

TOTAL POINTS 5

1. Let two matrices be

 $A = \begin{bmatrix} 4 & 3 \\ 6 & 9 \end{bmatrix}, \qquad B = \begin{bmatrix} -2 & 9 \\ -5 & 2 \end{bmatrix}$

- $\bigcirc \begin{bmatrix} 2 & -6 \\ 1 & 7 \end{bmatrix}$

To subtract B from A, carry out the subtraction element-wise.

Let $x = \begin{bmatrix} 2 \\ 7 \\ 4 \end{bmatrix}$

What is 3 * x?

- $\bigcirc \begin{bmatrix} \frac{2}{3} & \frac{7}{3} & \frac{4}{3} & \frac{1}{3} \end{bmatrix}$

- $\bigcirc \ \begin{bmatrix} 6 & 21 & 12 & 3 \end{bmatrix}$

✓ Correct

To multiply the vector \mathbf{x} by 3, take each element of \mathbf{x} and multiply that element by 3.

3. Let u be a 3-dimensional vector, where specifically

What is u^{T} ?

- O [4 1 8]

✓ Correct

4. Let u and v be 3-dimensional vectors, where specifically

1 / 1 point

1x3 dimensional matrix, and v can also be seen as a 3x1

matrix. The answer you want can be obtained by taking

the matrix product of \boldsymbol{u}^T and $\boldsymbol{v}.) \,$ Do not add brackets to your answer.

✓ Correct

5. Let A and B be 3x3 (square) matrices. Which of the following

must necessarily hold true? Check all that apply.

1 / 1 point

- $\square \qquad \qquad A*B = B*A$
- $\square \qquad \qquad A*B*A=B*A*B$

✓ Correct

Even though matrix multiplication is not commutative in general ($A*B \neq B*A$ for general matrices A,B), for the special case where A=I, we have A*B=I*B=B, and also B*A=B*I=B. So, A*B=B*A.

A + B = B + A

✓ Correct

We add matrices element-wise. So, this must be true.