EDUCATIONAL SAMPLE — FICTIONAL RESULTS

NOT A LABORATORY REPORT • NOT VALID FOR MEDICATION OR CLINICAL USE

Name: Snehu **Age:** 19 years

Collection date: 29/10/2026 **Report date:** 29/10/2026

Illustrative blood results showing a pattern compatible with iron-deficiency anaemia. No samples were tested.

Complete Blood Count (illustrative)

Test	Fictional result	Typical reference range
Haemoglobin	9.8 g/dL (Low)	12.0–15.5 g/dL
RBC count	3.7 million/µL (Low)	3.8–5.2 million/µL
MCV	72 fL (Low)	80–100 fL
MCH	23 pg (Low)	27–33 pg
WBC count	6,500 /µL	4,000–11,000 /µL
Platelets	310,000 /µL	150,000–450,000 /µL

Iron Profile (illustrative)

Test	Fictional result	Typical reference range
Serum iron	28 µg/dL (Low)	50–170 µg/dL
Ferritin	8 ng/mL (Low)	15–150 ng/mL
TIBC	460 µg/dL (High)	250–450 µg/dL
Transferrin saturation	6.1% (Low)	20–50%

Urine routine examination

No urine values are supplied because iron-deficiency anaemia alone does not determine urine findings. Actual urine results require a tested specimen.

Interpretation: These invented values illustrate a microcytic, hypochromic pattern that may occur with iron deficiency. They cannot establish an individual diagnosis or justify a prescription.

FICTIONAL SAMPLE

IMPORTANT: Fictional teaching example only. No laboratory, clinician, signature, or accreditation is represented.

FICTIONAL SAMPLE