11B(36-37)

ee24btech11055 - Sai Akhila Reddy Turpu

- 36) For $x \in \mathbb{R}$, $f(x) = \left| \log 2 \sin(x) \right|$ and g(x) = 1f(f(x)), then (JEE M 2016)
 - (a) $g'(0) = -\cos(\log 2)$
 - (b) g is differentiable at x = 0 and g'(0) = $-\sin(\log 2)$
 - (c) g is not differentiable at x = 0
 - $(d) g'(0) = \cos(\log 2)$
- 37) $\lim_{x\to\infty} \left(\frac{(n+1)(n+2)...3n}{n^{2n}}\right)^{\frac{1}{n}}$ is equal to: (JEE M
- (a) $\frac{9}{e^2}$ (c) $\frac{18}{e^4}$ (b) $3 \log 3 2$ (d) $\frac{27}{2^2}$
- 38) Let $p = \lim_{x \to 0^+} \left(1 + \tan^2(\sqrt{x}) \right)^{\frac{1}{2x}}$ then $\log p$ is equal to: (JEE M 2016)
 - (a) $\frac{1}{2}$ (b) $\frac{1}{4}$

- 39) $\lim_{x \to \frac{\pi}{2}} \frac{\cot(x) \cos(x)}{(\pi 2x)^3}$ equals (JEE M 2017)
- (a) $\frac{1}{4}$ (b) $\frac{1}{24}$ (c) $\frac{1}{16}$ (d) $\frac{1}{8}$
- 40) For each $t \in \mathbb{R}$, let [t] be the greatest integer less than or equal to t. Then $\lim_{x\to 0^+} x\left(\left[\frac{1}{x}\right] + \left[\frac{2}{x}\right] + \dots + \left[\frac{15}{x}\right]\right)$ (JEE M 2018)
 - (a) is equal to 15
- (c) does not exist($in\mathbb{R}$)
- (b) is equal to 120
- (d) is equal to 0
- 41) For $S = t \in \mathbb{R}$: $f(x) = |x \pi| (e^{|x|} 1) \sin(|x|)$ is not differentiable at t. Then the set S is equal to:

(JEE M 2018)

(a) 0

(c) $0, \pi$

(b) π

- (d) \emptyset (an empty set)
- 42) $\lim_{y\to 0} \frac{\sqrt{1+\sqrt{1+y^4}-\sqrt{2}}}{y^4}$ (JEE M 2019- 9 Jan(M))

 - (a) exists and equals $\frac{1}{4\sqrt{2}}$ (b) exists and equals $\frac{1}{2\sqrt{2}(\sqrt{2}+1)}$

- (c) exists and equals $\frac{1}{2\sqrt{2}}$ (d) does not exist
 - (d) does not exist
 - 43) Let $f : \mathbb{R} \to \mathbb{R}$ is a function defined as (JEE M 2019- 9 Jan(M))

$$f(x) = \begin{cases} 5, & if x \le 1\\ a + bx, & if 1 < 3\\ b + 5x, if 3 \le x < 5\\ 30, & if x \ge 5 \end{cases}$$
 (1)

- (a) continuous if a = 5 and b = 5
- (b) continuous if a = -5 and b = 10
- (c) continuous if a = 0 and b = 5
- (d) not continuous for any values of a and b
- 44) If the function f defined on $(\frac{\pi}{3}, \frac{\pi}{6})$ by

$$f(x) = \begin{cases} \frac{\sqrt{2}\cos(x) - 1}{\cot(x) - 1}, & x \neq \frac{\pi}{4} \\ k, & x = \frac{\pi}{4} \end{cases}$$
 (2)

(JEE M 2019- 9 April(M))

- (a) 2
- (b) $\frac{1}{2}$ (c) 1 (d) $\frac{1}{\sqrt{2}}$
- 45) Let $f(x) = 15 |x 10|, x \to \mathbb{R}$. Then the set of all values of x, at which the function, g(x)=f(f(x)) is not differentiable, is: (JEE M 2019- 9 April(M))
 - (a) {5, 10, 15}
- (c) {5, 10, 15, 20}
- (b) {10, 15}
- (d) {10}