## Microeconomic Theory I: A Notebook

With Jonathan Libgober

Sai Zhang

Check the Github Page of this project, or email me!

September 13, 2021

#### HERE WE GO!

This is my learning notebook of Microeconomic Theory I (Course number: ECON601 at USC Economics). As one of the core courses in an economic Ph.D. curriculum, Microeconomic Theory I is beyond important to my research. Therefore, I would love to use this notebook as a commitment mechanism, to document lecture notes, discuss session and office hour intuitions, reading summaries, my personal questions regarding the topics and more. By building a file from scratch, hopefull I could have a more systematic and sophisticated understanding on the content of this course.

I thank Prof. Jonathan Libgober at USC Economics for leading the discussion of the course and providing intuitive ways to understand microeconomic theory. Please check his webpage here, he is such fun.

I also appreciate the time and effort my TA Qitong Wang put into this course, guiding me through discussing sessions and problem sets. When I have questions, he is always there to help.

Following the structure of the course, this notebook will cover three aspects of microeconomic theories: (a) individual decision making, (b) game theory, (c) mechanism design and contract theory. Apart from Jonathan's lecture notes, I will also summarize the reading materials, including: Mas-Colell et al. (1995)'s Microeconomic Theory, Mailath (2018)'s Modelling Strategic Behavior<sup>1</sup>, Fudenberg and Tirole (1991)'s Game Theory, Myerson (1991)'s Game Theory: Analysis of Conflicts, Bolton and Dewatripont (2005)'s Contract Theory, Mailath and Samuelson (2006)'s Repeated Games and Reputation and Osborne and Rubinstein (1994)'s A Course in Game Theory. Other materials will also be referred to along the way.

Building this notebook is truly a memorable journey for me. I would love to share this review and all the related materials to anyone that finds them useful. And unavoidably, I would make some typos and other minor mistakes (hopefully not big ones). So I'd really appreciate any correction. If you find any mistakes, please send the mistakes to this email address saizhang.econ@gmail.com or start a branch on Github. BIG thanks in advance!

<sup>&</sup>lt;sup>1</sup>Latest version (May 2021) available here.

## **Contents**

I	Individual Decision Making	4
1	Preferences and Choices, Utilities	5
	1.1 Preference Relations	6 9 10 14 17 18 18 18 18 19
2	Fundamentals of Consumer Theory	20
3	Lagrange Maximization and Duality	21
4	Monotone Comparative Statics	22
5	Expected Utility and Decisionmaking under Uncertainty	23
6	Aggregation and the Existence of a Representative Consumer	24
7	Producer Theory	25
8	Stochastic Choice	26
II	Game Theory	27
9	Nash Equilibrium and Bayesian Nash Equilibrium	28
10	Rationalizability and DOminant Strategies	29
11	Correlated Equilibrium	30

12 Dynamic Games and Refinements	31
13 Repeated Games/Folk Theorem	32
14 Recursive Methods in Repeated Games	33
III Mechanism Design and Contract Theory	34
15 Arrow's Theorem and Social Choice	35
16 Boundaries of the Firm and Coase's Theorem	36
17 Implementation Concepts	37
18 The Revelation Principle	38
19 Auctions and Optimal Auctions	39
20 Efficient Implementation	40
21 Moral Hazard	41
22 Full Implementation	42
Bibliography	43

# Part I Individual Decision Making

#### PREFERENCES AND CHOICES, UTILITIES

#### **Contents**

6		 	 	Preference Relations	1.1	
9		 	 	Choice Rules	1.2	
10	. 1	 	 3	Linking Preferences with	1.3	
14	. 1	 	 	ntroducing Utility	1.4	
17	. 1	 	 	Commentary	1.5	
17	. 1	 	 scriptive model .	.5.1 Preference model		
18	. 1	 	 	.5.2 Empirical limits		
18	. 1	 	 	.5.3 Framing		
18	. 1	 	 	.5.4 Indecision		
18	. 1	 	 oilistic choice	.5.5 Inconsistency and		
19	. 1	 	 ce	.5.6 Determinants of p		
19	. 1	 	 ıe	.5.7 Range of choices a		

The first chapter summarizes the basic setting of individual decision making: preferences, choices and utilities. The main reference is Chapter 1 of Mas-Colell et al. (1995).

In this chapter, we will focus on 3 domains:

choice	given a set $A$ , what choice from $A$ is made
preference	given alternatives $x$ , $y$ , which does the decision maker prefers
utility	given an object <i>X</i> , how much does the DM likes <i>X</i> (as a number)

The starting point of individual decision problem is a *set of possible (mutually exclusive) alternatives* from which the individual must choose. To model decision making process on this set of alternatives, one can:

- either start from the tastes, i.e., *preference relations* of individuals, and set up the patterns of decision making with preferences
- or, start from the actual actions of individuals, i.e. *choices*, to deduct a pattern of decision making

With this two major approaches in mind, we know what's coming: the *rationality* of preferences and the central assumption of choices, the *Weak Axiom of Revealed Preference (WARP)*. And of course, the two approaches and two basic assumptions are parallel, so we need to figure out how link the (underlying) preferences and (observed) choices.

#### 1.1 Preference Relations

We start from the basic: weak preference relation,  $\geq$ .

**Definition 1.1.1.** A weak preference relation  $\geq$  on a set X is a subset of  $X \times X$ . If  $(x, y) \in \geq x$  is at least as good as y, written as  $x \geq y$ 

A weak preference relation will induce two other types of relations on *X*:

**Definition 1.1.2.** With  $\geq$  defined by Def. 1.1.1, we have

- the *strict preference relation*, > can be induced from  $\gtrsim$  as:  $x > y \Leftrightarrow x \gtrsim y \land y \not\gtrsim x$ , or in words, x if preferred to y.
- the *indifference relation*,  $\sim$  can be induced from  $\gtrsim$  as:  $x \sim y \Leftrightarrow x \gtrsim y \land y \gtrsim x$ , or in words, x is indifferent to y.

With the definition of these relations, we now define the central assumption of relations: *rationality*.

**Definition 1.1.3.** A weak preference relation  $\geq$  is *rational* if it is:

- Complete:  $\forall x, y \in X, x \gtrsim y \text{ or } y \gtrsim x \text{ or both}$
- Transitive:  $\forall x, y, z \in X, x \geq y \land y \geq z \Rightarrow x \geq z$

How to understand them? They are both strong assumptions:

- Completeness of ≥ means it is well-defined between any two possible alternatives. From the perspective of an individual, completeness means that she will make choices, and only meditated choices.
- Transitivity of ≥ implies that the decision maker will not have a preference cycle, since whoever has a preference cycle would suffer economically for it¹.

With the definition of rational  $\geq$  in Def. 1.1.3, we can prove the following properites

<sup>&</sup>lt;sup>1</sup>There are 2 types of violations of transitivity: irrational and mechanical. Irrational violations are easy to understand: decision makers simply do not follow transivity assumption, many reasons have been raised, including mental account, framing, menu effect, attraction effect, etc. Mechanical violations means that decision makers are "forced" to violate transitivity. One example of this type of violation is aggregation of considerations: decision makers may aggregate several sub-preferences as together to make the choice, leading to violation of transitivity. Another example is when the preference is only defined for differences above a certain level (problem of perceptible differences). See Mas-Colell et al. (1995, Page 7-8), Rubinstein (2012, Page 4-5) for details

of > and  $\sim$  *induced* by  $\gtrsim$ :

#### **Theorem 1.1.1.** If $\geq$ is rational, then:

- i. > is irreflexive (x > x never holds) and transitive ( $x > y \land y > z \Rightarrow x > z$ ) Proof:
  - irreflexive: by Def. 1.1.2,  $x > x \Rightarrow x \gtrsim x \land x \ngeq x$ , self contracdiction.
  - transitive:  $x > y \Rightarrow x \gtrsim y \land y \not\gtrsim x$ ,  $y > z \Rightarrow y \gtrsim z \land z \not\gtrsim y$ . By transitivity of  $\gtrsim$ ,  $x \gtrsim y \land y \gtrsim z \Rightarrow x \gtrsim z$ . If  $z \gtrsim x$ , by transitivity of  $\gtrsim$  and  $x \gtrsim y$ , we would have  $z \gtrsim y$ , contradicting y > z. Therefore  $x \gtrsim z \land z \not\gtrsim x \Rightarrow x > z$ .
- ii. ~ is reflexive  $(x \sim x, \forall x)$ , transitive  $(x \sim y \land y \sim z \Rightarrow x \sim z)$  and symmetric  $(x \sim y \Rightarrow y \sim x)$

#### Proof:

- reflexive: by completeness of  $\gtrsim$ ,  $\forall x, x \gtrsim x \Rightarrow x \sim x$
- transitive:  $x \sim y \Rightarrow x \gtrsim y \land y \gtrsim x$ ,  $y \sim z \Rightarrow y \gtrsim z$ ,  $z \gtrsim y$ , by the transitivity of  $\gtrsim$ , we have  $x \gtrsim z \land z \gtrsim x$ , hen  $x \sim z$
- symmetric:  $x \sim y \Rightarrow x \gtrsim y \land y \gtrsim x \Leftrightarrow y \gtrsim x \land x \gtrsim y \Rightarrow y \sim x$
- iii.  $x > y \gtrsim z \Rightarrow x > z$

<u>Proof</u>:  $x > y \Rightarrow x \gtrsim y \land y \not\gtrsim x$ , hence  $x > y \gtrsim z \Rightarrow x \gtrsim z$ . If  $z \gtrsim x$ , by transitivity of  $\gtrsim$ ,  $y \gtrsim x$ , contradicting x > y. Therefore,  $z \not\gtrsim x$ 

We can also directly define a *rational* > (see Kreps (1990, Page 19-21)):

**Definition 1.1.4.** A strict preference ralation > is rational if it is:

- asymmetric:  $\nexists x, y \in X$  s.t.  $x > y \land y > x$
- negatively transitive:  $x > y \Rightarrow \forall z \in X \setminus \{x, y\}, x > z \lor z > y \lor \text{ both.}$

With Def. 1.1.4 and Def. 1.1.3, we can prove that  $\geq$  is rational iff > is rational:

**Theorem 1.1.2.**  $\geq$  is rational  $\Leftrightarrow$  > is rational, specifically:

- $\geq$  is complete  $\Leftrightarrow$  > is asymmetric
- $\geq$  is transitive  $\Leftrightarrow$  > is negatively transitive

Now we prove this theorem:

**Step 1** proof  $\geq$  is rational  $\Rightarrow$  > is rational

#### - asymmetric

if  $\exists x, y \text{ s.t. } x > y \text{ and } y > x$ , then by the definition of induced strict preference, the pair x, y must satisfy

$$\begin{cases} x \gtrsim y \text{ and } y \not\gtrsim x & (x > y) \\ y \gtrsim x \text{ and } x \not\gtrsim y & (y > x) \end{cases}$$

which is, by completeness of rational  $\gtrsim$ , impossible. Therefore, such pair x, y don't exist. > is proved to be asymetric.

#### - negatively transitive

First,  $\forall z \notin \{x, y\}$ , by completeness of rational  $\geq$ , the relation between x and z is either  $x \geq z$  or  $z \geq x$ . Similarly, the relation between y and z is either  $y \geq z$  or  $z \geq y$ .

Second, given x > y, x, y satisfies  $x \gtrsim y$  and  $y \not\gtrsim x$ .

Also, it is easy to prove that:  $x > y \land y \gtrsim z \Rightarrow x > z, x > y \land z \gtrsim x \Rightarrow z > y$ ; and  $x > y \land z \sim x \Rightarrow z > y, x > y \land y \sim z \Rightarrow x > z$ 

Now we have the following scenarios:

- 1. if  $z \gtrsim x$  and  $y \gtrsim z$ , by transitivity of rational  $\gtrsim$ ,  $y \gtrsim x$ , contradicting the definition of x > y. This scenario doesn't exist.
- 2. if  $x \gtrsim z$  and  $y \gtrsim z$ , since x > y, with the auxiliary result proved above, we have x > z
- 3. if  $z \gtrsim x$  and  $z \gtrsim y$ , since x > y, with the auxiliary result proved above, we have z > y
- 4. if  $x \gtrsim z$  and  $z \gtrsim y$ , since x > y, suppose:
  - (a)  $z \gtrsim x$  as well, then  $x \sim z$ , in this case z > y;
  - (b)  $z \not\gtrsim x$ , then x > z
  - (c)  $y \gtrsim z$  as well, then  $y \sim z$ , in this case x > z
  - (d)  $y \not\gtrsim z$ , then z > y

therefore, a complete summary of (a) to (d) would give:

	$z \gtrsim x$	$z \not \succeq x$
$y \gtrsim z$	z > y & x > z	x > z
$y \not\gtrsim z$	z > y	x > z & z > y

Combining all above, we have proved negative transitivity of >.

With asymmetry and negative transitivity proved, we've proved that  $\gtrsim$  is rational  $\Rightarrow$  is rational

**Step 2** proof > is rational ⇒  $\gtrsim$  is rational.

- Complete: with a rational x > y, we know  $\nexists x, y$  s.t. x > y and y > x by asymmetry. Therefore,  $\forall x, y$ , we have two possibilities.
  - x > y and  $y \ne x$ , which would naturally induce a weak preference  $x \ge y$
  - y > x and x ≠ y, which would naturally induce a weak preference y ≳ x therefore,  $\forall x, y$ , either x ≳ y or y ≳ x completeness of ≳ is proven.
- Transitive: with a rational x > y, negative transivity gives  $\forall z \notin \{x, y\}$ , either x > z, z > y, or both. By negative transitivity, we have:
  - x > z: following same procedure, we know  $x \gtrsim z$ . If:
    - \*  $y \gtrsim z$ , since  $x > z \Rightarrow z \not\gtrsim x$ , by completeness we have  $x \gtrsim z$ , thus  $x \gtrsim y \land y \gtrsim z \Rightarrow x \gtrsim z$
    - \*  $z \gtrsim y$ , since  $x > y \Rightarrow x \not\gtrsim y$ , by completeness we have  $x \gtrsim y$ , thus  $x \gtrsim z \land z \gtrsim y \Rightarrow x \gtrsim y$
  - z > y: again, we know z ≿ y. If:
    - \*  $x \gtrsim z$ , since  $x > y \Rightarrow y \not\gtrsim x$ , by completeness we have  $x \gtrsim y$ , thus  $z \gtrsim y \land x \gtrsim z \Rightarrow x \gtrsim y$
    - \*  $z \gtrsim x$ , with  $x \gtrsim y$ , suppose  $y \gtrsim z$ , this contradicts z > y, thus  $z \gtrsim x \land x \gtrsim y \Rightarrow z \gtrsim y$
  - x > z and z > y: again we know  $x \gtrsim z$  and  $z \gtrsim y$ . Suppose  $y \gtrsim x$ , this contradicts x > y, therefore  $x \gtrsim z \land z \gtrsim y \Rightarrow x \gtrsim y$

In all three scenarios, transitivity is proved.

With completeness and transitivity proved, we've proved that  $\succ$  is rational  $\Rightarrow \gtrsim$  is rational.

Notice that negative positivity in Def. 1.1.4, is logically equivalent to its *contrapositive*:  $\exists z \in X \setminus \{x, y\}$  s.t.  $x \neq z \land z \neq y \Rightarrow x \neq y$ . This is percisely why the definition is called negative transitivity.

#### 1.2 Choice Rules

Next, we approach the theory of decision making from choice behavior itself. Formally, choice behavior is represented by means of a *choice structure* ( $\mathcal{B}$ ,  $C(\cdot)$ ). Now, we define choice structure ( $\mathcal{B}$ ,  $C(\cdot)$ ):

**Definition 1.2.1.** A choice structure  $(\mathcal{B}, C(\cdot))$  has two ingredients:

- $\mathcal{B} \subset \mathcal{P}(X) \setminus \emptyset$ , where  $\mathcal{P}(X)$  is the power set of X. This means, every element  $B \in \mathcal{B}$  is a subset of  $X^2$ .
- $C(\cdot)$  is a *choice rule correspondence* that assigns a nonempty set of chosen elements  $C(B) \subset B$ ,  $\forall B \in \mathcal{B}^3$ .

Now we discuss the CORE assumption in this section: the Weak Axiom of Revealed Preference (WARP):

**Definition 1.2.2.** A choice set  $(\mathcal{B}, C(\cdot))$  satisfies WARP if:

-  $\forall B, B'$  and  $x, y \in B \cap B', x \in C(B), y \in C(B') \Rightarrow x \in C(B')$ 

Or in words, WARP requires that if x is chosen from some alternatives where y is also available, then there can be NO budget set containing both x and y but only y is chosen.

Following WARP, define the *reveal preference relation*  $\gtrsim$ \* as:

**Definition 1.2.3.** Given a choice structure  $(\mathcal{B}, C(\cdot))$ ,  $x \gtrsim^* y \Leftrightarrow \exists B \in \mathcal{B}$  s.t.  $x, y \in B \land x \in C(B)$ 

In words, *x* is revealed at least as good as *y*.

With revealed preference defiend, we can rephrase WARP as: *If* x *is revealed at least as good as* y, *then* y *cannot be revealed preferred to* x. Hence,  $\gtrsim^*$  is not symmetric.

One thing to remember is that  $\geq^*$  need not be either complete or transitive. For  $\geq^*$  to be comparable, for a  $B \in \mathcal{B}$  and  $x, y \in B$ , we must have either  $x \in C(B)$ ,  $y \in C(B)$  or both.

An example is:

**Example 1.2.1.** Consider a choice structure 
$$(\mathcal{B}, C(\cdot))$$
 from  $X = \{x, y, z\}$ , where  $\mathcal{B} = \{\{x, y\}, \{x, y, z\}\}$ . Under WARP,  $C\{x, y\} = \{x\} \Rightarrow y \notin C\{x, y, z\}$ . BUT,

<sup>&</sup>lt;sup>2</sup>The elements  $B \in \mathcal{B}$  are so-called *budget sets*. The budget sets in  $\mathcal{B}$  should be thought of as an exhaustive listing of all the choice experiments that can be achieved, but it is possible that some subsets of X are not achievable.

<sup>&</sup>lt;sup>3</sup>The choice set C(B) can contain a single element, which is the choice among the alternatives in B. BUT, C(B) can contain multiple elements, then elements of C(B) are the *acceptable alternatives* in B.

we can have  $z \in C(\{x, y, z\})$ .

This is why the induced preference is called *revealed*: you don't know what else is going on.

There are 3 interesting properties of choice rules to keep in mind:

- Property  $\alpha$ :  $x \in A \subseteq B$ ,  $x \in C(B) \Rightarrow x \in C(A)$ .
- Property  $\beta$ :  $x, y \in C(A), A \subseteq B, y \in C(B) \Rightarrow x \in C(B)$
- Property  $\gamma$ :  $x \in C(B_1), x \in C(B_2) \cdots x \in C(B_n) \Rightarrow x \in C(B_1 \cup B_2 \cup \cdots \cup B_n)$

For a rational preference relation  $\gtrsim$ , its induced choice function  $C(\cdot, \gtrsim)$  satisfies the three properties:

- $\alpha$ : For  $x \in A \subseteq B$ ,  $x \in C(B, \geq)$ ,  $x \geq y$ ,  $\forall y \in B$ , and  $A \subseteq B \Leftrightarrow \forall z \in A, z \in B$ , hence  $\forall z \in A, x \geq z \Rightarrow x \in C(A, \geq)$
- $\beta$ : For  $x, y \in C(A, \geq)$ ,  $A \subseteq B$ ,  $y \in C(B, \geq)$ ,  $x \geq y$  and  $\forall z \in B$ ,  $y \geq z$ , by transitivity,  $\forall z \in B, x \geq z \Rightarrow x \in C(B, \geq)$

$$\gamma: x \in C(B_i, \gtrsim), i = 1, \cdots, n \Rightarrow \forall b_k^i \in B_i, x \gtrsim b_i \Rightarrow \forall b \in \{b_k^i\}, i = 1, \cdots, n, x_i \gtrsim b_i \Rightarrow x \in C(B_1 \cup B_2 \cup \cdots \cup B_n, \gtrsim)$$

With Sen's  $\alpha$  and  $\beta$  defined, we have:

**Theorem 1.2.1.** For a non-empty C, C satisfies  $\alpha$  and  $\beta \Leftrightarrow C$  satisfies WARP

Now, let's prove this.

- **Sen's**  $\alpha$  **and**  $\beta$   $\Rightarrow$  **WARP**: Suppose  $x, y \in A \cap B, x \in C(A), y \in C(B)$ . Let  $\tilde{A} = A \cap B$ , since  $\tilde{A} \subset A$  and  $\tilde{A} \subset B$ , Sen's  $\alpha$  gives  $x, y \in C(\tilde{A})$ . Now we have  $\tilde{A} \subset B$ ,  $y \in C(B)$ ,  $x, y \in C(\tilde{A})$ , by Sen's  $\beta$ ,  $x \in C(B)$ . Thus, WARP holds.
- **WARP** ⇒ **Sen's**  $\alpha$  **and**  $\beta$ : WARP gives  $x, y \in A \cap B, x \in C(B), y \in C(A) \Rightarrow x \in C(A)$ , in the special case of  $A \subset B$ , WARP will be reduced to Sen's  $\alpha$  and  $\beta$ .<sup>4</sup>

#### 1.3 Linking Preferences with Choices

Now we have two major approaches of decision making process: preference relations in Section 1.1 and choice rules in Section 1.2, what we need to do is to link them. This linkage will emerge when we examine two central assumptions: **rationality** and **WARP**. So the major question here is:

$$\mathbf{rational} \succsim^{???} (\mathcal{B}, C(\cdot)) \ \mathbf{satisfies} \ \mathbf{WARP}$$

And the answer is: YES! but not exactly. Now let's dig in.

<sup>&</sup>lt;sup>4</sup>Another way of proving this is to introduce rationalizing preference relation (Def.1.3.2). In fact, for  $\mathcal{B}$  includes all subsets of X of up to 3 elements, WARP  $\Leftrightarrow$  C is rationalizable  $\Leftrightarrow$  Sen's  $\alpha$  and  $\beta$  are satisfied.

#### Rational $\gtrsim \Rightarrow (\mathcal{B}, \mathcal{C}(\cdot))$ satisfies WARP

First, **rational**  $\gtrsim \Rightarrow$  ( $\mathcal{B}$ ,  $C(\cdot)$ ) **satisfies WARP** is a big YES. To prove this, we need to define *induced choice correspondence*:

**Definition 1.3.1.** Given a **rational**  $\geq$  on X, if the decision maker faces a nonempty subset of alternatives  $B \subset X$ , by maximizing her preference, she would choose any one of the elements in the *induced choice correspondence*:  $C^*(B, \geq) = \{x \in B : x \geq y, \forall y \in B\}$ 

The induced choice correspondence  $C^*(B, \geq)$  has an important property:

**Theorem 1.3.1.** if *X* is finite,  $C^*(B, \geq)$  will be **nonempty**.

A brief proof of this proposition is: If X is finite, B is finite as well. We will prove by induction. Starting from |B| = 1, the only element of B is in  $C^*(B, \gtrsim)$ . Now suppose  $C^*(B, \gtrsim)$  is nonempty when  $|B_n| = n$ , let  $x^* \in C^*(B_n, \gtrsim)$ ; when  $|B_{n+1}| = n + 1$ , let the n + 1th element y ( $\{y\} = B_{n+1} \setminus B_n$ ). By the completeness of a rational  $\gtrsim$ , either  $y \gtrsim x^*$  or  $x^* \gtrsim y$ :

- i.  $y \gtrsim x^*$ : since  $x^* \in C^*(B_n, \gtrsim) \Rightarrow x^* \gtrsim x$ ,  $\forall x \in B_n$ . By transitivity of  $\gtrsim$ ,  $y \gtrsim x$ ,  $\forall \in B_n$ . By completeness,  $y \gtrsim y$  as well. Hence,  $y \in C^*(B_{n+1}, \gtrsim)$ .
- ii.  $x^* \gtrsim y$ : since  $x^* \in C^*(B_n, \gtrsim) \Rightarrow x^* \gtrsim x$ ,  $\forall x \in B_n$ , hence  $x^* \gtrsim x$ ,  $\forall x \in B_n \cup y \Rightarrow x^* \in C^*(B_{n+1}, \gtrsim)$

Notice that when B is finite, a stronger condition of  $\geq$  being acyclic and complete is equilavent to an induced choice rule  $C^*(B, \geq) \neq \emptyset$ :

**Theorem 1.3.2.** For a finite B,  $\gtrsim$  is complete and **acyclic**  $\Leftrightarrow$   $C^*(B, \gtrsim) \neq \emptyset$ 

 $\gtrsim$  is acyclic mean that:  $b_1 \gtrsim b_2, b_2 \gtrsim b_3, \cdots, b_{n-1} \gtrsim b_n \Rightarrow b_n \not\gtrsim b_1$ . An example of transitive but not *acyclic* relations is indifference  $\sim$ :  $a_1 \sim a_2 \sim \cdots \sim a_n \Rightarrow a_n \sim a_1$ . A brief proof of Theorem 1.3.2 is:

- i.  $\operatorname{acyclic} \gtrsim \Rightarrow C^*(B, \gtrsim) \neq \emptyset$ : Suppose if  $C^*(B, \gtrsim) = \emptyset$ , for  $b_1 \in B$ ,  $b_1 \notin C^*(B, \gtrsim) \Rightarrow \exists b_2$  s.t.  $b_2 \gtrsim b_1$ . Continue this process, we can generate a sequence of  $\cdots \gtrsim b_2 \gtrsim b_1$ , since B is finite, this sequence must end at  $b_n$ . If  $\gtrsim$  is acyclic,  $b_1 \not\gtrsim b_n$ , this gives  $b_n > b_1$ , which would mean  $b_n$  must be in  $C^*(B, \gtrsim)$ , contradicting.
- ii.  $C^*(B, \geq) \neq \emptyset \Rightarrow$  acyclic  $\geq$ : Suppose  $\geq$  is not acyclic, then there exists  $b_1 \geq b_2 \geq \cdots \geq b_n \geq b_1$ , then for set  $B = \{b_1, b_2, \cdots, b_n\}$ ,  $\not\exists b^*$  s.t.  $b^* \geq b_i \forall b_i \in B$ , i.e.,  $C^*(B, \geq) = \emptyset$ .

Of course, we want to extend this to the situation where B is infinite. However, in general, it is possible that  $C^*(B, \gtrsim)$  is empty (if you set the most preferred side of B open, it would be impossible to choose based on the preferences). However, we can add some assumptions to finite non-emptiness and choice coherence, to ensure that choice function  $C(\cdot)$  is well-behaved on infinite sets as well.<sup>5</sup>

$$x \in A \subseteq NBT_{\geq_C}(x) \Rightarrow C(A) \neq \emptyset$$

This means that if everything in A is revealed to be no better than x and x is also in A, some choice will

<sup>&</sup>lt;sup>5</sup> The simplest assumption is: For a choice function  $C(\cdot)$  that satisfies finite non-emptiness and choice coherence, let the corresponding preference relation be  $\succeq_C$ , this preference will generate a no-better-than subset for x:  $NBT_{\succeq_C}(x) = \{x' \in X : x \succeq_C x'\}$ . With this setting, we can have:

With induced choice correspondence  $C^*(B, \geq)$  defined and non-emptyness proved, we can then say:

**Theorem 1.3.3.** If  $\gtrsim$  is a rational preference relation, then the choice structure generated by  $\gtrsim$ ,  $(\mathcal{B}, C^*(\cdot, \gtrsim))$ , satisfies WARP

We can prove this theorem quite easily:  $\forall B, B'$  suppose we have  $x, y \in B \cap B'$  and  $x \in C^*(B, \geq)$ ,  $y \in C^*(B', \geq)$ , then  $x \geq a$ ,  $\forall a \in B$  and  $y \geq b$ ,  $\forall b \in B'$ . Naturally, we have  $x \geq y$  since  $y \in B$ . By rationality (transitivity) of  $\geq$ , we have  $x \geq y \geq b$ ,  $\forall b \in B'$ , which means  $x \in C^*(B', \geq)$ . This is precisely the definition of WARP

#### $(\mathcal{B}, C(\cdot))$ satisfies WARP $\Rightarrow$ Rational $\gtrsim$

The proof of this direction is more subtle, and is NOT necessarily a yes. Again, we start from a auxiliary definition:

**Definition 1.3.2.** For a choice structure  $(\mathcal{B}, C(\cdot))$ , a rational preference relation  $\succeq$  **rationalizes**  $C(\cdot)$  relative to  $\mathcal{B}$  if  $C(B) = C^*(B, \succeq)$ ,  $\forall B \in \mathcal{B}$ .

In words, if for all budget sets  $B \in \mathcal{B}$ , the choices generated by a rational  $\geq$ , is just the choice rule  $C(\cdot)$ ,  $C(\cdot)$  is rationalized by  $\geq$ . This is, in a sense, constructing an explanation of decision making behavior with preferences.

We already proved that  $C^*(B, \gtrsim)$  satisfies WARP, which means that if a rationalizing preference relation to exist, WARP must be satisfied. However, if WARP is satisfied, a rationalizing preference relation does **NOT** necessarily exist.<sup>6</sup> Intuitiviely, more budget sets  $B \in \mathcal{B}$  would mean that, to satisfy WARP, choice behavior would be restricted more, and it is easier to be self-contradicting. Therefore, to pin down a rational preference relation to rationalize  $C(\cdot)$  relative to  $\mathcal{B}$ , we need to put some **restrictions on**  $\mathcal{B}$ .

**Theorem 1.3.4.** If  $(\mathcal{B}, C(\cdot))$  is a choice structure that:

- i. WARP is satisfied
- ii.  $\mathcal{B}$  includes **all** subsets of X of **up to 3** elements

then there exists a rational preference relation  $\geq$  s.t.  $C(B) = C^*(B, \geq)$ ,  $\forall B \in \mathcal{B}$ . And this rational  $\geq$  is the **only** preference relation that can rationalize  $(\mathcal{B}, C(\cdot))^7$ .

Now let's prove it, by examing the natural candidate for a rationalizing preference

be made out of A. This assumption is equivalment to the conclusion that for the infinite set A, we also have  $C(\cdot) \equiv C(\cdot)_{\geq C}$ . And of course, if we reverse the assumption above, we would have:

$$\forall x \in A, A \nsubseteq NBT_{\geq_C}(x) \Rightarrow C(A) = \emptyset$$

. This is easy to understand: in this case, x is the least preferred one in A, since A is infinite, you can never find an alternative that is preferred over all others.

<sup>6</sup>A simple example is:  $X = \{x, y, z\}$ ,  $\mathcal{B} = \{\{x, y\}, \{y, z\}, \{x, z\}\}$ . Since  $\mathcal{B}$  contains 3 binary menus, the choice structure  $C(\{x, y\}) = \{x\}, C(\{y, z\}) = \{y\}, C(\{x, z\}) = \{z\}$  vacuously satisfy WARP. But, this choice structure cannot be rationalized since it contradicts transitivity.

<sup>7</sup>The existence of a rationalizing preference relation  $\succeq$  brings many interesting properties, one of them is *path-invariance*:  $\forall B_1, B_2 \in \mathcal{B}, B_1 \cup B_2 \in \mathcal{B} \land C(B_1) \cup C(B_2) \in \mathcal{B} \Rightarrow C(B_1 \cup B_2) = C(C(B_1) \cup C(B_2))$ , meaning that the decision problem can safely be subdivided. A proof is: for  $x \in C(B_1 \cup B_2)$  and  $y \in C(B_1) \cup C(B_2)$ , since  $C(B_1) \cup C(B_2) \subset B_1 \cup B_2$ , thus  $x \in C(B_1 \cup B_2) \Rightarrow x \in C(C(B_1) \cup C(B_2))$ ; for  $x \in C(C(B_1) \cup C(B_2))$  and  $y \in B_1 \cup B_2$ , we have  $x \succeq z, \forall z \in C(B_1), C(B_2) \Rightarrow x \succeq w, \forall w \in B_1 \cup B_2 \Rightarrow x \in C(B_1 \cup B_2)$ 

#### relation: the **revealed preference relation** $\gtrsim^*$ :

#### **Step 1** Prove that $\geq^*$ is rational

- Completeness: By (ii) of Def.1.3.2, all binary subsets of X are in  $\mathcal{B}$ . Hence,  $\{x,y\} \in \mathcal{B}$ . For this binary menu,  $C(\{x,y\})$  must contain either x or y, therefore,  $x \gtrsim^* y$  or  $y \gtrsim^* x$  or both. Completeness proved.
- Transitivity:  $\forall \{x, y, z\} \in \mathcal{B}$ ,  $C(\{x, y, z\}) \neq \emptyset$ . Suppose  $x \gtrsim^* y, y \gtrsim^* z$ , which implies that  $x \in C(\{x, y\})$ ,  $y \in C(\{y, z\})$ , we then have three cases for  $C(\{x, y, z\})$ :
  - a.  $x \in C(\{x, y, z\})$ , WARP gives that  $x \in C(\{x, z\}) \Rightarrow x \gtrsim^* z$
  - b.  $y \in C(\{x, y, z\})$ , we have  $x \in C(\{x, y\})$ . WARP gives  $x \in C(\{x, y, z\})$   $\Rightarrow x \gtrsim^* z$
  - c.  $z \in C(\{x, y, z\})$ , we have  $y \in C(\{y, z\})$ . WARP gives  $y \in C(\{x, y, z\})$ , and  $x \in C(\{x, y\})$ , WARP gives  $x \in C(\{x, y, z\}) \Rightarrow x \gtrsim^* z$

Hence,  $x \gtrsim^* y$ ,  $y \gtrsim^* z \Rightarrow x \gtrsim^* z$ 

#### **Step 2** Prove that $\geq^*$ rationalizes $C(\cdot)$ on $\mathcal{B}$

Now, we need to show  $\forall B \in \mathcal{B}, C(B) = C^*(B, \geq^*)$ . Logically, this means the revealed preference  $\geq^*$  inferred from  $C(\cdot)$  actually generates  $C(\cdot)$ . Formally, we prove it in 2 steps:

- a. Suppose  $x \in C(B)$ , which means that  $\forall y \in B, x \gtrsim^* y$  (by Def.1.2.3), hence  $x \in C^*(B, \gtrsim^*)$  (by Def.1.3.1). This proves  $C(B) \subseteq C^*(B, \gtrsim^*)$
- b. Suppose  $x \in C^*(B, \geq^*)$ , which means that  $\forall y \in B, x \geq^* y$  (by Def.1.3.1). Therefore,  $\forall y \in B$ , there must exist a set  $B_y \in \mathcal{B}$  s.t.  $x, y \in B_y \Rightarrow x \in C(B_y)$ . Since  $C(B) \neq \emptyset$ , suppose  $z \in C(B)$ , since  $x \in C(B_z)$ , WARP implies that  $x \in C(B)$ . This proves  $C^*(B, \geq^*) \subseteq C(B)$

Together, we have  $C(B) = C^*(B, \geq^*)$ .

#### **Step 3** Prove $\gtrsim^*$ is the unique choice

Since  $\mathcal{B}$  includes all two-element subsets of X, the choice behavior in  $C(\cdot)$  completely determines the pairwise preference relations over X of any rationalizing preference.

Now, it is **proved**! Notice that the main assumption(restriction) here is  $\mathcal{B}$  **includes all subsets of** X **of up to 3 elements**, this gives completeness, which is fundamental.

#### Things to keep in mind

We have proved the twoway links of preferences and choices:

- Rational  $\gtrsim \Rightarrow (\mathcal{B}, C^*(\cdot, \geq))$  satisfies WARP (see Thm.1.3.3)
- A WARP-satisfying, up-to-3-element  $(\mathcal{B}, C(\cdot))$  can be uniquely rationalized by a rational  $\gtrsim$  (see Thm.1.3.4)

However, there are still something to keep in mind.

First, for a given choice structure  $(\mathcal{B}, C(\cdot))$ , there may be **more than one** rationalizing preference relation  $\succeq$  in general. Here is the simplest example: For  $X = \{x, y\}, \mathcal{B}\{\{x\}, \{y\}\}$  and the choice structure  $C(\{x\}) = \{x\}, C(\{y\} = \{y\})$ . In this case, **ANY** relation preference relation of X can rationalize  $C(\cdot)$  This is related to both

Def.1.3.2 and (ii) of Thm.1.3.4. Thm.1.3.4 gives that if  $\mathcal{B}$  contains **ALL binary** menus of X, then there could be at most one rationalizing preference relation.

Second, the restriction for WARP $\Rightarrow$  rational  $\gtrsim$ , namely  $\mathcal B$  containing all subsets of up to 3 elements, is too strong. For many economic problems, we will not consider all possible subsets, or limit ourselves to up-to-3-element ones. A strengthened version of WARP will be introduced later for that purpose.

Finally, up till now, we define a rationalizing preference as one:  $C(B) = C^*(B, \gtrsim)$  (Def.1.3.2). A common alternative would be to require only  $C(B) \subset C^*(B, \gtrsim)$ : if C(B) is a **subset** of the most preferred choices generated by  $\gtrsim$ , i.e.,  $C^*(B, \gtrsim)$ . This will allow indifferences to be more than the situation of anything might be picked. And it is empirically intuitive in a sense that observed choices will never fully reveal decision makers' entire preferencing maximizing choice set. Naturally,  $C(B) \subset C^*(B, \gtrsim)$  is weaker than  $C(B) = C^*(B, \gtrsim)$ . But  $C(B) \subset C^*(B, \gtrsim)$  has an interesting property: the all-indifferent preference will be able to rationalize *any* choice behavior. Therefore, when  $C(B) \subset C^*(B, \gtrsim)$  is used, you would always need to put some additional restrictions on the rationalizing preference relation for the specific economic context.

#### 1.4 Introducing Utility

Now, with preferences and choices defined, and the linkage between the two established, we need to transfer these concepts into math for analytic studies. This is exactly why utility functions are introduced: to assign a number and rank the elements in *X* according to preferences.

**Definition 1.4.1.** A function  $u: X \to \mathbb{R}$  is a utility function representing relation  $\geq if \ \forall x, y \in X, x \geq y \Leftrightarrow u(x) \geq u(y)$ 

Notice that a utility function representing a preference relation  $\geq$  is NOT unique. **Rank-preserving** is the only requirement, hence, any strictly increasing function  $f: \mathbb{R} \to \mathbb{R}$ , v(x) = f(u(x)) will also represent  $\geq$  as  $u(\cdot)$ . The logic is quite straight forward: for  $x, y \in X$  and  $u(\cdot)$  represents  $\geq$ , then  $x \geq y \Leftrightarrow u(x) \geq u(y)$ , for a strictly increasing  $f(\cdot)$ ,  $u(x) \geq u(y) \Leftrightarrow f(u(x)) \geq f(u(y)) \Leftrightarrow v(x) \geq v(y)$ , hence  $v(\cdot)$  represents  $\geq$  as well. The major requirement here is **strictly increasing**  $f(\cdot)$ .

Two concepts to keep in mind:

- 1. **Ordinal** properties of utility functions: the **invariant** properties of  $u(\cdot)$  across all of its strictly increasing transformations  $f(u(\cdot))$ . Ranking (i.e. the preference represented by utility functions) is ordinal.
- 2. **Cardinal** properties of utility functions: the **variant** properties of  $u(\cdot)$  across all of its strictly increasing transformation  $f(u(\cdot))$ . Numerical values associated with the alternatives in X (i.e. the magnitude of the differences between alternatives) is cardinal.

The numerical value, or even the size of relative differences have no particular meaning. Only ranking is "real", since the "level of utility" is **unobservable** and anything required to know the "level of utility" is **untestable**.

The central theorem of utility functions is closely linked to rationality:

**Theorem 1.4.1.** A preference relation  $\gtrsim$  can be represented by a utility function  $\Rightarrow \gtrsim$  is rational

#### The proof is

- *Completeness*. Since  $u(\cdot)$  represents preference relations between alternatives, and  $u: X \to \mathbb{R}$ , thus  $\forall x, y \in X$ , either  $u(x) \ge u(y)$  or  $u(y) \ge u(x)$ . By Def.1.4.1, we have either  $x \ge y$  or  $y \ge x$ , hence  $\ge$  is complete.
- Transitivity. For  $x \gtrsim y$ ,  $y \gtrsim z$ . By Def.1.4.1,  $u(x) \geq u(y)$ ,  $u(y) \geq u(z)$ , hence  $u(x) \geq u(z) \Rightarrow x \gtrsim z$ .

What about the other way? It is true, subject to some prerequisites:

**Theorem 1.4.2.**  $\gtrsim$  is rational and X is **finite**  $\Rightarrow$  there is a utility function representing  $\gtrsim$ .

The major prerequisite here is X being **finite**. The proof is done by induction: Suppose there are N elements in X:

- When N = 1, any number could be assigned to that element as its utility.
- Suppose a rational  $\gtrsim$  on  $X = x_1, x_2, \cdots, x_{N-1}$  could be represented by a utility function  $u(\cdot)$ . Without losing generality, we can assume  $u(x_1) \le u(x_2) \le \cdots \le u(x_{N-1})$ . For the Nth element  $x_N$ , by the rationality of  $\gtrsim$ , we have three scenarios:

i 
$$\forall i \in 1, \dots, N-1, x_N \geq x_i$$
: by Def.1.4.1,  $u(x_N) \geq u(x_i)$ .

- ii  $\forall i \in 1, \dots, N-1, x_i \gtrsim x_N:, u(x_N) \leq u(x_i).$
- iii  $\exists i, j \in 1, \dots, N-1, i \neq j, x_j \gtrsim x_N \gtrsim x_i$ :  $u(x_j) \geq u(x_N) \geq u(x_i)$ . By completeness and transitivity,  $x_1, x_2, \dots, x_{N-1}$  can be "divided" by  $x_N$ , meaning that for  $I = \{i : x_N \gtrsim x_i\}$  and  $J = \{j : s_j \gtrsim x_N\}$ ,  $I \cup J = \{1, \dots, N-1\}$ . Note that we have assumed the index as the ranking, hence let  $i^* = \max I, j^* = \min J, i^*+1 = j^*$ , hence we must have  $u(x_i) \leq u(x_{i^*}) \leq u(x_j) \leq u(x_j)$

In all 3 scenarios,  $u(\cdot)$  represents  $\geq$  on  $X = \{x_1, \dots, x_{N-1}, x_N\}$  as well.

With this induction, we prove Thm.1.4.28.

Now extend Thm.1.4.2 from finite *X* to countable infinite *X*:

**Theorem 1.4.3.**  $\gtrsim$  is rational and X is **countable**  $\Rightarrow$  there is a utility function representing  $\gtrsim$ .

To prove Thm.1.4.3, we can construct a utility function: for any set S, its emuneration  $\{s_1, s_2, \dots\}$  (which exists if S is countable), define an auxiliary function  $d: S \to \mathbb{R}$  as  $d(s_i) = \left(\frac{1}{2}\right)^n$ , then for a countable set  $X = \{x_1, x_2, \dots\}$ , the utility of any element  $\tilde{x} \in X$  can be defined as

$$u(x^*) = \sum_{\tilde{x}_i \in NBT(x^*)} d(\tilde{x}_i)$$

<sup>9</sup> where  $NBT(x^*)$  is the set of all elements that are **n**ot **b**etter than  $x^*$ , i.e.  $NBT(x^*) = \{\tilde{x}_i : \tilde{x}_i \in X \land x^* \gtrsim \tilde{x}_i\}$ . It is easy to see that  $NBT(x^*)$  is a countable subset of X. Suppose

<sup>&</sup>lt;sup>8</sup>Another way of proof is: Start with  $x^{start} \in X$ , define  $W_x = \{y : y < x^{start}\}$  then  $W_x$  is either empty or not: If not empty, pick  $\tilde{x} \in W_x$ , shrink  $W_x$  to  $\{y : y < \tilde{x}\}$  and repeat this procedure till  $a \sim x^{stop}$  where u(a) = 0, then  $x^{stop}$  is the "lower bound" of the set. With this process, we can generate a utility function for any finite set X that is rational.

Notice that  $\lim_{n\to\infty} \sum_{i=1}^n \frac{1}{2^i} = 1$ , utility is bounded to [0, 1).

 $NBT(x^*)$  has k elements (k < n), we can calculate the utility  $u(x^*) = \sum_{i=1}^k \left(\frac{1}{2}\right)^i$ . After this construction, rest of the proof is trivial:  $\forall x, y \in X, x \gtrsim y \Rightarrow NBT(x) \supseteq NBT(y)$ , which means that NBT(x) has at least as many elements as NBT(y), by the constructed utility function, it is easy to see  $u(x) \ge u(y)$ ; Conversely,  $u(x) \ge u(y)$  simply means that NBT(x) contains at least as many elements as NBT(y) does, which directly leads to  $x \gtrsim y$ .

Now, let's figure out the difficult question: what about uncountable sets? Here is a very general proposition:

**Theorem 1.4.4.** For a rational preference  $\geq$  on a set X,  $\geq$  can be represented by  $u(\cdot)$  if and only if some countable set  $X^*$  of X has the property that  $\forall x, y \in X$ ,  $x > y \Rightarrow \exists x^* \in X^*$  s.t.  $x \geq x^* > y$ .

The proof of Thm.1.4.4 is not that difficult:

Step 1: such  $X^*$  exists  $\Rightarrow$  rational  $\gtrsim$  can be represented by  $u(\cdot)$ .

- Given such  $X^*$ ,  $x \gtrsim y \Rightarrow u(x) \geq u(y)$ Suppose  $X^*$  exists, let it be  $X^* = \{x_1^*, x_2^*, \cdots\}$ . Again, define  $d(x_n^*) = \frac{1}{2^n}$ , we can then construct a utility function as

$$\forall x \in X, u(x) = \sum_{\tilde{x}_i^* \in X^* \cap NBT(x)} d(\tilde{x}_i^*)$$

Since  $x \gtrsim y \Leftrightarrow NBT(x) \supseteq NBT(y)$ , hence  $NBT(x) \cap X^* \supseteq NBT(y) \cap X^*$ , which, by the construction of  $\sum \frac{1}{2^n}$ , leading to  $u(x) \ge u(y)$ .

- Given such  $X^*$ ,  $u(x) \ge u(y) \Rightarrow x \ge y$ We can prove the contrapositive:  $y \not\succeq x \Rightarrow u(y) \not\succeq u(x)$ . Given the rationality of  $\ge$ ,  $y \not\succeq x \Rightarrow x > y$ , then  $\exists x^* \in X^*$  s.t.  $x \ge x^* > y$ , hence we know NBT(x) is strictly larger then NBT(y) (NBT(x) includes  $x^*$ ), therefore, by definition,  $u(x) > u(y) \Rightarrow u(y) \not\succeq u(x)$ .
- Step 2: rational  $\geq$  can be represented by  $u(\cdot) \Rightarrow$  such  $X^*$  exists We want to prove this, but it is very difficult to prove in general, so we construct a special case: Let  $\{I_n\}$  be a set of all closed intervals with rantional endpoints, that is, each  $I_n$  is an interval of  $[\underline{q}_n, \bar{q}_n]$  where  $\bar{q}_n > \underline{q}_n$  are rational numbers. The set of rational numbers is countable, the cross product of two countable sets is also countable, hence  $\{I_n\}$  is countable as well. Let u(X) denote the set of real numbers  $\{r \in \mathbb{R} : \exists x \in X, r = u(x)\}$ , there will be 3 possible scenarios:
  - i.  $u(X) \cap I_n \neq \emptyset$ : for each  $I_n$ , pick one  $x \in X$  s.t.  $u(x) \in I_n$  and name it  $x_n$
  - ii.  $u(X) \cap I_n = \emptyset$ : let  $\bar{r}_n = \inf\{r \in u(X) : r > \bar{q}_n\}$ . If  $\exists x \in X$  s.t.  $u(x) = \bar{r}_n$ , choose one such x and name it  $x_n$
  - iii.  $u(X) \cap I_n = \emptyset \land \forall x \in X, \bar{r}_n \neq u(x)$ , no x will be defined as  $x_n$

If we define  $X^*$  as the collection of all  $x_n$  in case i) and ii). Notice there is less than one  $x_n$  for each  $I_n$  and  $I_n$  is countable, hence  $X^*$  is countable as well.

Suppose x > y for  $x, y \in X$ , we have  $u(\cdot)$  representing  $\geq$ , hence u(x) > u(y). Choose some rational number q in the open interval (u(y), u(x)) and let  $\bar{r} = \inf\{r \in u(X) : r > q\}$ . Given this setup, we have  $u(x) \geq \bar{r}$  since u(x) > q. If:

-  $u(x) > \bar{r}$ : we can always find a rational number q' s.t.  $u(x) > q' > \bar{r}$ . Let n be the index of the interval [q, q'], since  $q < \bar{r} < q'$ ,  $\bar{r} \in u(X) \cap [q, q'] \Rightarrow$ 

- $u(X) \cap [q, q'] \neq \emptyset$ . Therefore,  $\exists x^* \in X^*$ , namely  $x_n$ , s.t.  $u(x^*) \in [q, q']$ , leading to  $u(x) > u(x^*) > u(y)$ .
- $u(x) = \bar{r}$ : we can always find a rational number q' s.t. q > q' > u(y). Let n be the index of the interval [q, q']. If  $u(X) \cap [q', q] \neq \emptyset$ , then  $\exists x^* \in X^*$  s.t.  $u(x) \geq q \geq u(x) \geq q' \geq u(y)$ , then  $x \gtrsim x^* > y$ . If  $u(X) \cap [q, q'] = \emptyset$ , then [q, q'] fits into category (ii) above, and  $\exists x^* \in X^*$ , namely  $x_n$ , such that  $u(x^*) = \bar{r} = u(x)$ . With this  $x^*$ , we have  $u(x^*) = u(x) > u(y) \Rightarrow x \gtrsim x^* > y$ .

This is a very smart proof, and it is very general as well. However, you would have to make sure that the countable seubset  $X^*$  exists, which is not very practical. This problem leads to topological  $\geq$ , which will be covered later.

It is natural that if we start from >, we would have the same logic (See Kreps, 1990, Page 30):

- similar definition:  $x > y \Leftrightarrow u(x) > u(y)$
- similar theorems:
  - i  $\exists u(\cdot)$  representing  $> \Rightarrow$  rational > (asymmetric and negatively transitive)
  - ii If X is finite or at least countably infinite,  $\exists u(\cdot)$  representing  $> \Leftrightarrow$  rational >

#### 1.5 Commentary

In this section, I discuss some of common commentaries on the standard preference model presented above.

#### 1.5.1 Preference model as a descriptive model

A common complaint about the standard utility maximization/preference ranking model is that no one in reality actually calculates a number as utility before making choices. This comment has a lot of sense to it since we rarely care about utility, let alone doing some math, before grocery shopping. But this observation does NOT invalidate the usefulness of preference/utility model.

The standard model does NOT regulate agents to consciously maximize utility, instead, it assumes individuals act *as if* they maximize utility. Mathematically, we have already proven that if choice behavior satisfies finite nonemptiness and WARP, then something will be chosen, and agents' choice behavior is just *as if* it were preference driven, or the choice behavior can be linked to a preference. If the set of choices is countable, then the preference-driven choice can be indexed by numbers, hence, becomes a mathematical question.

Utility/preference/choice system is considered as a description of choice behavior. Long as people do make a choice, and that choice satisfies WARP, we can always find a numerical way to *describe* the behavioral pattern.

#### 1.5.2 Empirical limits

To verify utility maximization as a model of choices over the choice space X, we need to check every subset A of it. And we also need to know all of C(A). Of course we have already managed to verify the two-way link between preference, utility and choice for all choice menus with no more than 3 elements, but above that, it would be extremely difficult. Empirically, we will observe (at best) C(A) for finitely many subsets of X, we would most likely observe only the *one* element that is selected out of C(A) while failing to identify equally-preferred alternatives simply because they are not observed to be chosen. In these scenarios, how can we tell whether our observations are aligned with utility maximization? In later chapters, we will come back to this problem.

#### 1.5.3 Framing

The way bundles are framed/presented can affect how they are perceived, hence influce individuals' decision making process. One of the most cited economic research by Kahneman and Tversky (1979) discussed this problem in a very clear and innovative fashion. Framing will be a problem if it induces violence of WARP: a is picked when comparing to b, but when c is available as well, b will be chosen instead. This may look silly and will never happen in real life but numerous examples of violation of WARP can be raised due to the framing problem. Designers/publicists are actually trained to take advantage of this "irrationality" to influence consumers' decision making process. A hugh strand of literature in behavioral economics discuss and explore the framing problem, limited attention, heuristics, impatience are introduced to explain such phenomena.

#### 1.5.4 Indecision

Another big problem is that agents may just NOT be able to make a decision. Sometimes the differences between alternatives are trivial or too complicated to measure, the problem of indecision could rise. Rational preferences gives that for each pair of objects x and y, an agent can choose between: x is better than y, y is better than x, x and y are equal. However, if we add another option I can't tell which is better, the transitivity would be violated quite easily. Consider it this way, the choice of "I can't decide" allows  $C(A) = \emptyset$  even for a finite set of alternatives, this simply goes against the model we have built up.

#### 1.5.5 Inconsistency and probabilistic choice

It is widely documented that an agent could be inconsistent about her choices. This could be an issue of framing, anchoring, indecision, or just unjustifiable inconsistency. This brings the stochastic side of choices: agents' choices may be subject to many random factors hence not deterministic. This will be discussed more thoroughly in later chapters.

#### 1.5.6 Determinants of preference

Since the model is a description of choice behavior, it does not provide any intuition on how a decision is made and what are the determinants of preferences. Later, we will talk about dynamic choice, where an agent's experiences affect her subsequent choices. The standard model needs to be adjusted to incorporating the evolving decision making process through time. Another situation is welfare analysis. Institutional factors would need to be included in the models: preferences could be partially determined socially, different social classes, countries, religions, cultures will likely lead agents to have different preferences. This has been examined more and more by institutional economists, I will include some inspiring works later.

#### 1.5.7 Range of choices as a value

Nobel Laureate Amartya Sen was

FUNDAMENTALS OF CONSUMER THEORY

LAGRANGE MAXIMIZATION AND DUALITY

### MONOTONE COMPARATIVE STATICS

## EXPECTED UTILITY AND DECISIONMAKING UNDER UNCERTAINTY

## AGGREGATION AND THE EXISTENCE OF A REPRESENTATIVE CONSUMER

### PRODUCER THEORY

### STOCHASTIC CHOICE

# Part II Game Theory

# NASH EQUILIBRIUM AND BAYESIAN NASH EQUILIBRIUM

# RATIONALIZABILITY AND DOMINANT STRATEGIES

## CORRELATED EQUILIBRIUM

### DYNAMIC GAMES AND REFINEMENTS

REPEATED GAMES/FOLK THEOREM

RECURSIVE METHODS IN REPEATED GAMES

# Part III Mechanism Design and Contract Theory

### ARROW'S THEOREM AND SOCIAL CHOICE

## BOUNDARIES OF THE FIRM AND COASE'S THEOREM

### IMPLEMENTATION CONCEPTS

### THE REVELATION PRINCIPLE

### AUCTIONS AND OPTIMAL AUCTIONS

### EFFICIENT IMPLEMENTATION

### MORAL HAZARD

### FULL IMPLEMENTATION

#### **BIBLIOGRAPHY**

- Patrick Bolton and Mathias Dewatripont. *Contract Theory*, volume 1. The MIT Press, 2005. URL https://ideas.repec.org/b/mtp/titles/0262025760.html.
- Drew Fudenberg and Jean Tirole. Game Theory. MIT Press, 1991.
- Daniel Kahneman and Amos Tversky. Prospect theory: An analysis of decision under risk. *Econometrica*, 47(2):363–391, 1979.
- David M. Kreps. A Course in Microeconomic Theory. Princeton University Press, 1990.
- George J Mailath. *Modeling Strategic Behavior: A Graduate Introduction to Game Theory and Mechanism Design*, volume 6. World Scientific, 2018.
- George J. Mailath and Larry Samuelson. Repeated Games and Reputations: Long-Run Relationships. Oxford University Press, 2006. URL https://ideas.repec.org/b/oxp/obooks/9780195300796.html.
- Andreu Mas-Colell, Michael Dennis Whinston, et al. *Microeconomic theory*, volume 1. New York: Oxford university press, 1995.
- Roger B. Myerson. *Game Theory: Analysis of Conflict*. Harvard University Press, 1991. ISBN 9780674341166. URL http://www.jstor.org/stable/j.ctvjsf522.
- Martin J. Osborne and Ariel Rubinstein. *A Course in Game Theory*. The MIT Press, 1994. URL https://ideas.repec.org/b/mtp/titles/0262650401.html.
- Ariel Rubinstein. *Lecture Notes in Microeconomic Theory: The Economic Agent Second Edition*. Princeton University Press, rev revised, 2 edition, 2012.