

Topic 20: Random Forest

by Sai Zhang

Key points: .

Disclaimer: The note is built on Prof. *Jinchi Lv*'s lectures of the course at USC, DSO 607, High-Dimensional Statistics and Big Data Problems.

20.1 Motivation

Denote by $m(\mathbf{X})$ the measurable nonparametric regression function with p -dimensional random vector \mathbf{X} taking values in $[0, 1]^p$. The Random Forest algorithm aims to learn the regression function in a non-parametric way based on the observations $\mathbf{x}_i \in [0, 1]^p, y_i \in \mathbb{R}, i = 1, \dots, n$, from the model

$$y_i = m(\mathbf{x}_i) + \epsilon_i$$

where $\mathbf{X}, \mathbf{x}_i, \epsilon_i, i = 1, \dots, n$ are independent, and $\{\mathbf{x}_i\}$ and $\{\epsilon_i\}$ are two sequences of identically distributed random variables. \mathbf{x}_i is distributed identically as \mathbf{X} .

Why Random Forest (RF)? RF has gained significant popularity due to its

- **High accuracy:** RF consistently rank among the top performers

Chi et al. (2022)

References

Chien-Ming Chi, Patrick Vossler, Yingying Fan, and Jinchi Lv. Asymptotic properties of high-dimensional random forests. *The Annals of Statistics*, 50(6):3415–3438, 2022.