

## Topic 11: Lasso

by Sai Zhang

**Key points:****Disclaimer:**

Lasso (Least absolute Shrinkage and Selection Operator), proposed by Tibshirani (1996), aims to minimize the **SSR (sum of residual squares)** subject to the **L1-norm (sum of the absolute value)** of the coefficients being less than a constant.

**11.1 Set up**

For data  $(\mathbf{x}_i, y_i)_{i=1}^n$ , where

- $y_i$  is the outcome for individual  $i$
- $\mathbf{x}_i = (x_{i1}, \dots, x_{ip})'$  is the  $p \times 1$  vector of predictors

Then the Lasso estimator  $(\hat{\alpha}, \hat{\beta})$  is defined as

$$(\hat{\alpha}, \hat{\beta}) = \arg \min_{\alpha, \beta} \left\{ \sum_{i=1}^n \left( y_i - \alpha - \sum_{j=1}^p \beta_j x_{ij} \right)^2 \right\} \quad \text{s.t.} \quad \sum_{j=1}^p |\beta_j| \leq \lambda$$

for the  $n \times 1$  response vector  $\mathbf{y} = (y_1, \dots, y_n)'$ , the  $n \times p$  design matrix  $\mathbf{X} = (\mathbf{x}_1, \dots, \mathbf{x}_n)'$  where  $\mathbf{x}_i = (x_{i1}, \dots, x_{ip})'$  is a  $p \times 1$  vector. Here  $\hat{\alpha} = \bar{y}$ , w.l.o.g., let  $\bar{y} = 0$  and omit  $\alpha$  for simplicity.

In matrix form, we have

- constrained form:

$$\hat{\beta} = \arg \min_{\beta \in \mathbb{R}^p} \left\{ \frac{1}{2n} \|\mathbf{y} - \mathbf{X}\beta\|_2^2 \right\} \quad \text{s.t.} \quad \|\beta\|_1 \leq \lambda$$

- unconstrained form:

$$\hat{\beta}(\lambda) = \arg \min_{\beta \in \mathbb{R}^p} \left\{ \frac{1}{2n} \|\mathbf{y} - \mathbf{X}\beta\|_2^2 + \lambda \|\beta\|_1 \right\}$$

where the regularization parameter  $\lambda \geq 0$ :

- $\lambda \rightarrow \infty$ :  $\hat{\beta}_{lasso} \rightarrow \hat{\beta}_{OLS}$
- $\lambda = 0$ :  $\hat{\beta}_{lasso} = \mathbf{0}$

## 11.2 Solving Lasso

Lasso is essentially a quadratic optimization problem. Hence, the solution is given by taking the derivative (of the unconstrained question) and set it equal to 0


$$\frac{d}{d\beta} \left( \frac{1}{2n} \right)$$

## 11.3 Penalized Least Square Estimation

Lasso is one special class of Penalized Least Square (PLS) Estimation. For the linear regression model  $\mathbf{y} = \mathbf{X}\beta + \epsilon$ , if  $\epsilon \sim \mathcal{N}(\mathbf{0}, \sigma^2 \mathbf{I}_n)$ , we have PLS as

$$\min_{\beta \in \mathbb{R}^p} \left\{ \frac{1}{2n} \|\mathbf{y} - \mathbf{X}\beta\|_2^2 + \sum_{j=1}^p p_\lambda(|\beta_j|) \right\}$$

where  $p_\lambda(\cdot)$  is a penalty function indexed by the regularization parameter  $\lambda \geq 0$



## References

Robert Tibshirani. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society: Series B (Methodological)*, 58(1):267–288, 1996.