

Noise-Induced Randomization in Regression Discontinuity Designs

Dean Eckles, Nikolaos Ignatiadis, Stefan Wager, Han Wu

Presented by: Sai Zhang

November 18, 2022

Outline

1 Discussion

Discussion

Literature: Continuity-Based RD

Most popular: local linear regression (Hahn et al., 2001; G. W. Imbens and Lemieux, 2008)

$$\hat{\tau}_c = \arg \min_{\tau} \left\{ \sum_{i=1}^n \underbrace{K}_{\text{weighting}} \left(\underbrace{\frac{|Z_i - c|}{h_n}}_{\text{bandwidth}} \right) (Y_i - a - \tau W_i - \beta_- (Z_i - c)_- - \beta_+ (Z_i - c)_+)^2 \right\}$$

;

Literature: Continuity-Based RD

Most popular: local linear regression (Hahn et al., 2001; G. W. Imbens and Lemieux, 2008)

$$\hat{\tau}_c = \arg \min_{\tau} \left\{ \sum_{i=1}^n \underbrace{K}_{\text{weighting}} \left(\underbrace{\frac{|Z_i - c|}{h_n}}_{\text{bandwidth}} \right) (Y_i - a - \tau W_i - \beta_- (Z_i - c)_- - \beta_+ (Z_i - c)_+)^2 \right\}$$

;

Literature: Continuity-Based RD

Most popular: local linear regression (Hahn et al., 2001; G. W. Imbens and Lemieux, 2008)

$$\hat{\tau}_c = \arg \min_{\tau} \left\{ \sum_{i=1}^n \underbrace{K}_{\text{weighting}} \left(\underbrace{\frac{|Z_i - c|}{h_n}}_{\text{bandwidth}} \right) (Y_i - a - \tau W_i - \beta_- (Z_i - c)_- - \beta_+ (Z_i - c)_+)^2 \right\}$$

- $\mu_{(w)}(z) = \mathbb{E}[Y(w) \mid Z = z]$ is **smooth**
- h_n decays at an **appropriate** rate

;

Literature: Continuity-Based RD

Most popular: local linear regression (Hahn et al., 2001; G. W. Imbens and Lemieux, 2008)

$$\hat{\tau}_c = \arg \min_{\tau} \left\{ \sum_{i=1}^n \underbrace{K}_{\text{weighting}} \left(\underbrace{\frac{|Z_i - c|}{h_n}}_{\text{bandwidth}} \right) (Y_i - a - \tau W_i - \beta_- (Z_i - c)_- - \beta_+ (Z_i - c)_+)^2 \right\}$$

- $\mu_{(w)}(z) = \mathbb{E}[Y(w) \mid Z = z]$ is **smooth**
- h_n decays at an **appropriate** rate

Robust CIs (Armstrong and Kolesár, 2020; Calonico et al., 2014; Kolesár and Rothe, 2018);

Literature: Continuity-Based RD

Most popular: local linear regression (Hahn et al., 2001; G. W. Imbens and Lemieux, 2008)

$$\hat{\tau}_c = \arg \min_{\tau} \left\{ \sum_{i=1}^n \underbrace{K}_{\text{weighting}} \left(\underbrace{\frac{|Z_i - c|}{h_n}}_{\text{bandwidth}} \right) (Y_i - a - \tau W_i - \beta_- (Z_i - c)_- - \beta_+ (Z_i - c)_+)^2 \right\}$$

- $\mu_{(w)}(z) = \mathbb{E}[Y(w) \mid Z = z]$ is **smooth**
- h_n decays at an **appropriate** rate

Robust CIs (Armstrong and Kolesár, 2020; Calonico et al., 2014; Kolesár and Rothe, 2018); Data-adaptive bandwidths (G. Imbens and Kalyanaraman, 2012)

Literature: Continuity-Based RD extended

$$\mu_{(w)}(z) = \mathbb{E}[Y(w) \mid Z = z]$$

If further assume **convexity** of $\mu_{(w)}(z)$, e.g.:

$$\left| \mu''_{(w)}(z) \right| \leq B, \forall z \in \mathbb{R}$$

Literature: Continuity-Based RD extended

$$\mu_{(w)}(z) = \mathbb{E}[Y(w) \mid Z = z]$$

If further assume **convexity** of $\mu_{(w)}(z)$, e.g.:

$$\left| \mu''_{(w)}(z) \right| \leq B, \forall z \in \mathbb{R}$$

Optimization-based RD: the treatment effect τ_c can be estimated (minimax linear estimation) via **numerical convex optimization** (Armstrong and Kolesár, 2018; G. Imbens and Wager, 2019)

Link Noise-Induced RD and Continuity-Based RD

$$\begin{aligned}\mu_{(w)}(z) &= \mathbb{E}[Y(w) \mid Z = z] \\ &= \frac{\int \mathbb{E}[Y(w) \mid U = u] p(z \mid u) dG(u)}{f_G(z)} = \frac{\int \alpha_{(w)}(u) p(z \mid u) dG(u)}{\int p(z \mid u) dG(u)}\end{aligned}$$

Link Noise-Induced RD and Continuity-Based RD

$$\begin{aligned}\mu_{(w)}(z) &= \mathbb{E}[Y(w) \mid Z = z] \\ &= \frac{\int \mathbb{E}[Y(w) \mid U = u] p(z \mid u) dG(u)}{f_G(z)} = \frac{\int \boxed{\alpha_{(w)}(u)} p(z \mid u) dG(u)}{\int p(z \mid u) dG(u)}\end{aligned}$$

Link Noise-Induced RD and Continuity-Based RD

$$\begin{aligned}\mu_{(w)}(z) &= \mathbb{E}[Y(w) \mid Z = z] \\ &= \frac{\int \mathbb{E}[Y(w) \mid U = u] p(z \mid u) dG(u)}{f_G(z)} = \frac{\int \boxed{\alpha_{(w)}(u)} p(z \mid u) dG(u)}{\int p(z \mid u) dG(u)}\end{aligned}$$

Then the worst-case possible curvature is:

$$\text{Curv}(z, \rho, p) = \sup \left\{ \left| \frac{d^2 \mu_{(w)}(z)}{dz^2} \right| : f_G(z) = \int p(z \mid u) dG(u) \geq \rho > 0, \alpha_{(w)}(\cdot) \in [0, 1] \right\}$$

Link Noise-Induced RD and Continuity-Based RD

$$\begin{aligned}\mu_{(w)}(z) &= \mathbb{E}[Y(w) \mid Z = z] \\ &= \frac{\int \mathbb{E}[Y(w) \mid U = u] p(z \mid u) dG(u)}{f_G(z)} = \frac{\int \boxed{\alpha_{(w)}(u)} p(z \mid u) dG(u)}{\int p(z \mid u) dG(u)}\end{aligned}$$

Then the worst-case possible curvature is:

$$\text{Curv}(z, \rho, p) = \sup \left\{ \left| \frac{d^2 \mu_{(w)}(z)}{dz^2} \right| : f_G(z) = \int p(z \mid u) dG(u) \geq \rho > 0, \alpha_{(w)}(\cdot) \in [0, 1] \right\}$$

Armstrong and Kolesár (2020): fit 4th-degree polynomials to $\mu_{(0)}(z)$ and $\mu_{(1)}(z)$, and take the largest estimated curvature obtained anywhere

Literature: Randomization Inference RD

Posit a non-trivial interval \mathcal{I} with $c \in \mathcal{I}$ s.t.

$$\{Y_i(0), Y_i(1)\} \perp Z_i \mid \{Z_i \in \mathcal{I}\}$$

then focus on this interval, perform classical **randomized** study inference

Literature: Randomization Inference RD

Posit a non-trivial interval \mathcal{I} with $c \in \mathcal{I}$ s.t.

$$\{Y_i(0), Y_i(1)\} \perp Z_i \mid \{Z_i \in \mathcal{I}\}$$

then focus on this interval, perform classical **randomized** study inference

- Design-based approach (Rubin, 2008)

Literature: Randomization Inference RD

Posit a non-trivial interval \mathcal{I} with $c \in \mathcal{I}$ s.t.

$$\{Y_i(0), Y_i(1)\} \perp Z_i \mid \{Z_i \in \mathcal{I}\}$$

then focus on this interval, perform classical **randomized** study inference

- Design-based approach (Rubin, 2008)

- Strong assumption

No **data-driven way** of choosing \mathcal{I}

If the interval \mathcal{I} is known a-priori, the problem collapses to a **RCT**

Measurement Error Induced RD

Rokkanen (2015) considers a similar approach, assuming:

Measurement Error Induced RD

Rokkanen (2015) considers a similar approach, assuming:

- noisy running variables **(A2)** and exogeneity of the noise **(A3)**
- **NOT** assuming prior knowledge of the noise distribution $p(\cdot | u)$

Measurement Error Induced RD

Rokkanen (2015) considers a similar approach, assuming:

- noisy running variables **(A2)** and exogeneity of the noise **(A3)**
- **NOT** assuming prior knowledge of the noise distribution $p(\cdot | u)$
- A stronger assumption: observing at least 3 noisy measurements of the latent variable U_i , $\{Z_i, Z'_i, Z''_i\}$

Measurement Error Induced RD

Rokkanen (2015) considers a similar approach, assuming:

- noisy running variables **(A2)** and exogeneity of the noise **(A3)**
- **NOT** assuming prior knowledge of the noise distribution $p(\cdot | u)$
- A stronger assumption: observing at least 3 noisy measurements of the latent variable U_i , $\{Z_i, Z'_i, Z''_i\}$

Measurement Error Induced RD

Rokkanen (2015) considers a similar approach, assuming:

- noisy running variables **(A2)** and exogeneity of the noise **(A3)**
- **NOT** assuming prior knowledge of the noise distribution $p(\cdot | u)$
- A stronger assumption: observing at least 3 noisy measurements of the latent variable U_i , $\{Z_i, Z'_i, Z''_i\}$
 - (U_i, Z_i, Z'_i, Z''_i) is **joint normal**
 - $\alpha_{(w)}(u) = \mathbb{E}[Y_i(w) | U_i = u]$ is **linear** w.r.t. u

RD with Ordinal Running Variables

Similarly, ordinal Z_i (bond rating, custody security score, etc.) can be seen as a noisy measurement of a latent variable U_i .

Li et al. (2021) assume

$$U_i = \mathbf{X}_i\beta$$

then use **inverse-propensity weighting** with estimated propensities $e(u) = \mathbb{P}[Z_i \geq c \mid U_i = u]$ for inference.

RD with Ordinal Running Variables

Similarly, ordinal Z_i (bond rating, custody security score, etc.) can be seen as a noisy measurement of a latent variable U_i .

Li et al. (2021) assume

$$U_i = \mathbf{X}_i\beta$$

then use **inverse-propensity weighting** with estimated propensities $e(u) = \mathbb{P}[Z_i \geq c \mid U_i = u]$ for inference.

Assuming: U_i can be observed, and well predicted by \mathbf{X}_i

Measurement Errors

- The **running variable** is unobserved, only a noisy measurement is observed
Bartalotti et al. (2021), Davezies and Le Barbanchon (2017), Dong and Kolesár (2021), and Pei and Shen (2017)
- Measurement error in causal inference beyond RD
Jiang and Ding (2020), Kuroki and Pearl (2014), and Pearl (2012)

A Comparison

RD designs Assumptions

Noise-induced RD	a known distribution of the measurement error $p(\cdot \mid u)$
Noise-induced RD (Rokkanen, 2015)	multiple joint-normal noisy measurements (U_i, Z_i, Z'_i, Z''_i)
	linear $\alpha_{(w)}(u) = \mathbb{E}[Y_i(w) \mid U_i = u]$

Continuity-based RD	$\mu_{(w)} = \mathbb{E}[Y(w) \mid Z = z]$ is smooth
OPTimization-based RD	convexity of $\mu_{(w)}(z)$: $ \mu''_{(w)}(z) \leq B, \forall z \in \mathbb{R}$
Randomization inference RD	an "RCT" interval \mathcal{I} : $\{Y_i(0), Y_i(1)\} \perp Z_i \mid \{Z_i \in \mathcal{I}\}$
RD with ordinal Z_i	U_i can be observed, and well predicted by \mathbf{X}_i

References I

- Armstrong, T. B., & Kolesár, M. (2018). Optimal inference in a class of regression models. *Econometrica*, 86(2), 655–683.
- Armstrong, T. B., & Kolesár, M. (2020). Simple and honest confidence intervals in nonparametric regression. *Quantitative Economics*, 11(1), 1–39.
- Bartalotti, O., Brummet, Q., & Dieterle, S. (2021). A correction for regression discontinuity designs with group-specific mismeasurement of the running variable. *Journal of Business & Economic Statistics*, 39(3), 833–848.
- Calonico, S., Cattaneo, M. D., & Titiunik, R. (2014). Robust nonparametric confidence intervals for regression-discontinuity designs. *Econometrica*, 82(6), 2295–2326.
- Davezies, L., & Le Barbanchon, T. (2017). Regression discontinuity design with continuous measurement error in the running variable. *Journal of econometrics*, 200(2), 260–281.
- Dong, Y., & Kolesár, M. (2021). When can we ignore measurement error in the running variable? *arXiv preprint arXiv:2111.07388*.

References II

- Eckles, D., Ignatiadis, N., Wager, S., & Wu, H. (2020). Noise-induced randomization in regression discontinuity designs. *arXiv preprint arXiv:2004.09458*.
- Hahn, J., Todd, P., & Van der Klaauw, W. (2001). Identification and estimation of treatment effects with a regression-discontinuity design. *Econometrica*, 69(1), 201–209.
- Imbens, G. W., & Lemieux, T. (2008). Regression discontinuity designs: A guide to practice. *Journal of econometrics*, 142(2), 615–635.
- Imbens, G., & Kalyanaraman, K. (2012). Optimal bandwidth choice for the regression discontinuity estimator. *The Review of economic studies*, 79(3), 933–959.
- Imbens, G., & Wager, S. (2019). Optimized regression discontinuity designs. *Review of Economics and Statistics*, 101(2), 264–278.
- Jiang, Z., & Ding, P. (2020). Measurement errors in the binary instrumental variable model. *Biometrika*, 107(1), 238–245.
- Kolesár, M., & Rothe, C. (2018). Inference in regression discontinuity designs with a discrete running variable. *American Economic Review*, 108(8), 2277–2304.

References III

- Kuroki, M., & Pearl, J. (2014). Measurement bias and effect restoration in causal inference. *Biometrika*, 101(2), 423–437.
- Li, F., Mercatanti, A., Mäkinen, T., & Silvestrini, A. (2021). A regression discontinuity design for ordinal running variables: Evaluating central bank purchases of corporate bonds. *The Annals of Applied Statistics*, 15(1), 304–322.
- Pearl, J. (2012). On measurement bias in causal inference. *arXiv preprint arXiv:1203.3504*.
- Pei, Z., & Shen, Y. (2017). The devil is in the tails: Regression discontinuity design with measurement error in the assignment variable. In *Regression discontinuity designs*. Emerald Publishing Limited.
- Rokkanen, M. A. (2015). Exam schools, ability, and the effects of affirmative action: Latent factor extrapolation in the regression discontinuity design.
- Rubin, D. B. (2008). For objective causal inference, design trumps analysis. *The annals of applied statistics*, 2(3), 808–840.

Thank you!