

Spatial Attention Tunes Temporal Processing in Early Visual Cortex by Speeding and Slowing Alpha Oscillations

Poppy Sharp, Tjerk Gutteling, David Melcher, Clayton Hickey

Presented by: Sai Zhang

November 8, 2022

Outline

1 Materials and Methods

Materials and Methods

Trial Structure




Figure 1: Trial Structure

Trial Structure





Figure 1: Trial Structure

Timeline:

- pre-cue: 1000-1500ms
- cue interval: 850-1350ms (randomized)
- display: 16.67ms
- ISI: 48.3ms
- response delay: 400ms

Trial Structure




visual cue: **red** cross

- 75% (T): one of the arms turn **green** (75% valid)
- 25% (C): neutral cue

Figure 1: Trial Structure

Trial Structure




2 displays:
complementary,
non-overlapping

- neutral cue: the 2 displays **complete** each other
- empty: one left **empty** in both

Figure 1: Trial Structure

Trial Structure



task: moving a highlighted square

- **segregation:** targeting the half circle
- **integration:** targeting the empty spot

Figure 1: Trial Structure

Measures

Eye tracking

MEG recording

Measures

Eye tracking

- sampling rate: 1kHz
- rejection:
 - saccades: $7 \pm 7\%$ trials
 - blinks: $3 \pm 4\%$ trials

MEG recording

Measures

Eye tracking

- sampling rate: 1kHz
- rejection:
 - saccades: $7 \pm 7\%$ trials
 - blinks: $3 \pm 4\%$ trials

MEG recording

- estimate instantaneous α -frequency: 7- to 14-Hz frequency band
- rejection
 - nonbiological noise: 10 ± 1 channels

Analysis

Source analysis

Numerical analysis

Analysis

Source analysis

- combine head digitization data with anatomic MRI data
- regions of interest:
 - parietal cortex
 - occipital cortex

Numerical analysis

Analysis

Source analysis

- combine head digitization data with anatomic MRI data
- regions of interest:
 - parietal cortex
 - occipital cortex

Numerical analysis

- method: 2-way repeated ANOVA
- noise of raw estimates of α frequency:
center on results following a neutral-cue
 - within each of the
integration/segragation conditions
separately

Other technical details

- Participants: 29 (**normal/corrected-to-normal** vision; age 24 ± 2.7 years; 11 male, 18 female)

Other technical details

- Participants: 29 (**normal/corrected-to-normal** vision; age 24 ± 2.7 years; 11 male, 18 female)
- Stimuli: projected at 120Hz onto a translucent screen, at a viewing distance of **1m**

Other technical details

- Participants: 29 (**normal/corrected-to-normal** vision; age 24 ± 2.7 years; 11 male, 18 female)
- Stimuli: projected at 120Hz onto a translucent screen, at a viewing distance of **1m**
- Pre-MEG: 30 practice trials to achieve at least 25% accuracy

Other technical details

- Participants: 29 (**normal/corrected-to-normal** vision; age 24 ± 2.7 years; 11 male, 18 female)
- Stimuli: projected at 120Hz onto a translucent screen, at a viewing distance of **1m**
- Pre-MEG: 30 practice trials to achieve at least 25% accuracy
- Number of trials: 10 blocks \times 67 trials/block

Other technical details

- Participants: 29 (**normal/corrected-to-normal** vision; age 24 ± 2.7 years; 11 male, 18 female)
- Stimuli: projected at 120Hz onto a translucent screen, at a viewing distance of **1m**
- Pre-MEG: 30 practice trials to achieve at least 25% accuracy
- Number of trials: 10 blocks \times 67 trials/block
- Base for numerical analysis: a shift in the **neutral-cue** baseline emerges equally in **ipsilateral** and **contralateral** signals

References I

Sharp, P., Gutteling, T., Melcher, D., & Hickey, C. (2022). Spatial attention tunes temporal processing in early visual cortex by speeding and slowing alpha oscillations. *Journal of Neuroscience*, 42(41), 7824–7832.

