# Noise-Induced Randomization in Regression Discontinuity Designs

Dean Eckles, Nikolaos Ignatiadis, Stefan Wager, Han Wu


Presented by: Sai Zhang


November 18, 2022

## Outline

Key Argument


Key Argument





### Assumption 1: Sharp RD design


- **I.I.D.** samples  $\{Y_i(0), Y_i(1), Z_i\} \in \mathbb{R}^3, i = 1, \dots, n$
- lacktriangledown treatment assignment:  $W_i=1$  ( $\{Z_i\geq c\}$ ), where  $c\in\mathbb{R}$  is the **cutoff**
- lacksquare observation:  $\{Y_i,Z_i\}$  where  $Y_i=Y_i(W_i)$



### **Assumption 2: Noisy running variable**

$$Z_i \mid U_i \sim p\left(\cdot \mid U_i\right)$$


where  $p(\cdot \mid \cdot)$  is a **known** conditional density w.r.t. to a measure  $\lambda$ , the latent variable  $U_i$  has an **unknown** distribution G



### **Assumption 2: Noisy running variable**

$$Z_i \mid U_i \sim \mathcal{N}(U_i, \nu^2), \nu > 0$$


where  $p(\cdot \mid \cdot)$  is a **known** conditional density w.r.t. to a measure  $\lambda$ , the latent variable  $U_i$  has an **unknown** distribution G



### **Assumption 2: Noisy running variable**

$$Z_i \mid U_i \sim \text{Binomial}(K, U_i), K \in \mathbb{N}$$


where  $p(\cdot \mid \cdot)$  is a **known** conditional density w.r.t. to a measure  $\lambda$ , the latent variable  $U_i$  has an **unknown** distribution G



### **Assumption 3: Exogeneity**

$$[\{Y_i(0),Y_i(1)\}\perp Z_i]\mid U_i$$


which implies  $\mathbb{E}\left[Y_{i}\mid U_{i},Z_{i}\right]=lpha_{\left(W_{i}\right)}\left(u\right)$ 




### **Assumption 3: Exogeneity**

$$[\{Y_i(0),Y_i(1)\}\perp Z_i]\mid U_i$$

which implies  $\mathbb{E}\left[Y_i \mid U_i, Z_i\right] = \alpha_{(W_i)}\left(u\right)$ , where  $\alpha_{(w)}\left(u\right) = \mathbb{E}\left[Y_i\left(w\right) \mid U_i = u\right]$  is the response functions for the potential oucomes conditional on the latent variable u



- A1 **Sharp** RD
- A2 Noisy  $Z_i$ :  $Z_i \mid U_i \sim p(\cdot \mid U_i)$
- A3 Exogeneity:  $\overline{\left[\left\{Y_{i}\left(0\right),Y_{i}\left(1\right)\right\} \perp Z_{i}\right] \mid U_{i}}$



- A1 Sharp RD
- A2 Noisy  $Z_i$ :  $Z_i \mid U_i \sim p(\cdot \mid U_i)$
- A3 Exogeneity:  $[\{Y_i(0), Y_i(1)\} \perp Z_i] \mid U_i$

### **Proposition 1**

Let  $\gamma_+(\cdot), \gamma_-(\cdot)$  be measurable functions of Z, then under A1-A3:

$$\mathbb{E}\left[\gamma_{+}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(1)}\left(U\right)h\left(U,\gamma_{+}\right)\right], \qquad \qquad \mathbb{E}\left[\gamma_{-}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(0)}\left(U\right)h\left(U,\gamma_{-}\right)\right]$$

where 
$$h\left(u,\gamma\right)\coloneqq\int\gamma\left(z\right)p\left(z\mid u\right)\mathrm{d}\lambda\left(z\right)$$
,  $\alpha_{\left(w\right)}\left(u\right)=\mathbb{E}\left[Y_{i}\left(w\right)\mid U_{i}=u\right]$ 

### **Proposition 1**

Let  $\gamma_+(\cdot), \gamma_-(\cdot)$  be measurable functions of Z, then under A1-A3:

$$\mathbb{E}\left[\gamma_{+}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(1)}\left(U\right)h\left(U,\gamma_{+}\right)\right], \qquad \qquad \mathbb{E}\left[\gamma_{-}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(0)}\left(U\right)h\left(U,\gamma_{-}\right)\right]$$

where 
$$h(u, \gamma) := \int \gamma(z) p(z \mid u) d\lambda(z)$$
,  $\alpha_{(w)}(u) = \mathbb{E}[Y_i(w) \mid U_i = u]$ 

### **Proposition 1**

Let  $\gamma_+(\cdot), \gamma_-(\cdot)$  be measurable functions of Z, then under A1-A3:

$$\mathbb{E}\left[\gamma_{+}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(1)}\left(U\right)h\left(U,\gamma_{+}\right)\right], \qquad \qquad \mathbb{E}\left[\gamma_{-}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(0)}\left(U\right)h\left(U,\gamma_{-}\right)\right]$$

where 
$$h\left(u,\gamma\right)\coloneqq\int\gamma\left(z\right)p\left(z\mid u\right)\mathrm{d}\lambda\left(z\right),\ \alpha_{\left(w\right)}\left(u\right)=\mathbb{E}\left[Y_{i}\left(w\right)\mid U_{i}=u\right]$$

$$\blacksquare \mathbb{E}\left[Y^2\right], \mathbb{E}\left[\gamma_-\left(Z\right)^2\right], \mathbb{E}\left[\gamma_+\left(Z\right)^2\right] < \infty$$

### **Proposition 1**

Let  $\gamma_{+}(\cdot), \gamma_{-}(\cdot)$  be measurable functions of Z, then under A1-A3:

$$\mathbb{E}\left[\gamma_{+}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(1)}\left(U\right)h\left(U,\gamma_{+}\right)\right], \qquad \qquad \mathbb{E}\left[\gamma_{-}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(0)}\left(U\right)h\left(U,\gamma_{-}\right)\right]$$

where 
$$h\left(u,\gamma\right)\coloneqq\int\gamma\left(z\right)p\left(z\mid u\right)\mathrm{d}\lambda\left(z\right),\ \alpha_{\left(w\right)}\left(u\right)=\mathbb{E}\left[Y_{i}\left(w\right)\mid U_{i}=u\right]$$

- $\blacksquare \mathbb{E}\left[Y^2\right], \mathbb{E}\left[\gamma_-\left(Z\right)^2\right], \mathbb{E}\left[\gamma_+\left(Z\right)^2\right] < \infty$
- $> \gamma_+(\cdot), \gamma_-(\cdot)$  are weighting functions s.t.

### **Proposition 1**

Let  $\gamma_+(\cdot), \gamma_-(\cdot)$  be measurable functions of Z, then under A1-A3:

$$\mathbb{E}\left[\gamma_{+}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(1)}\left(U\right)h\left(U,\gamma_{+}\right)\right], \qquad \qquad \mathbb{E}\left[\gamma_{-}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(0)}\left(U\right)h\left(U,\gamma_{-}\right)\right]$$

where 
$$h\left(u,\gamma\right)\coloneqq\int\gamma\left(z\right)p\left(z\mid u\right)\mathrm{d}\lambda\left(z\right)$$
,  $\alpha_{\left(w\right)}\left(u\right)=\mathbb{E}\left[Y_{i}\left(w\right)\mid U_{i}=u\right]$ 

- $\blacksquare \mathbb{E}\left[Y^2\right], \mathbb{E}\left[\gamma_-\left(Z\right)^2\right], \mathbb{E}\left[\gamma_+\left(Z\right)^2\right] < \infty$
- - $-\gamma_{+}(z)=0$  for z< c: assign non-zero weights only to treated units

### **Proposition 1**

Let  $\gamma_+(\cdot), \gamma_-(\cdot)$  be measurable functions of Z, then under A1-A3:

$$\mathbb{E}\left[\gamma_{+}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(1)}\left(U\right)h\left(U,\gamma_{+}\right)\right], \qquad \qquad \mathbb{E}\left[\gamma_{-}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(0)}\left(U\right)h\left(U,\gamma_{-}\right)\right]$$

where 
$$h\left(u,\gamma\right)\coloneqq\int\gamma\left(z\right)p\left(z\mid u\right)\mathrm{d}\lambda\left(z\right),\ \alpha_{\left(w\right)}\left(u\right)=\mathbb{E}\left[Y_{i}\left(w\right)\mid U_{i}=u\right]$$

- $\blacksquare \mathbb{E}\left[Y^2\right], \mathbb{E}\left[\gamma_-\left(Z\right)^2\right], \mathbb{E}\left[\gamma_+\left(Z\right)^2\right] < \infty$
- $> \gamma_+(\cdot), \gamma_-(\cdot)$  are weighting functions s.t.
  - $-\gamma_{+}(z)=0$  for z< c: assign non-zero weights only to treated units
  - $\gamma_{-}\left(z\right)=0$  for  $z\geq c$ : assign non-zero weights only to control units

### **Proposition 1**

Let  $\gamma_{+}(\cdot), \gamma_{-}(\cdot)$  be measurable functions of Z, then under A1-A3:

$$\mathbb{E}\left[\gamma_{+}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(1)}\left(U\right)h\left(U,\gamma_{+}\right)\right], \qquad \qquad \mathbb{E}\left[\gamma_{-}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(0)}\left(U\right)h\left(U,\gamma_{-}\right)\right]$$

where 
$$h\left(u,\gamma\right)\coloneqq\int\gamma\left(z\right)p\left(z\mid u\right)\mathrm{d}\lambda\left(z\right)$$
,  $\alpha_{\left(w\right)}\left(u\right)=\mathbb{E}\left[Y_{i}\left(w\right)\mid U_{i}=u\right]$ 

$$\mathbb{E}\left[\gamma_{+}\left(Z\right)Y\mid U\right]$$

### **Proposition 1**

Let  $\gamma_{+}(\cdot), \gamma_{-}(\cdot)$  be measurable functions of Z, then under A1-A3:

$$\mathbb{E}\left[\gamma_{+}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(1)}\left(U\right)h\left(U,\gamma_{+}\right)\right], \qquad \qquad \mathbb{E}\left[\gamma_{-}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(0)}\left(U\right)h\left(U,\gamma_{-}\right)\right]$$

where 
$$h\left(u,\gamma\right)\coloneqq\int\gamma\left(z\right)p\left(z\mid u\right)\mathrm{d}\lambda\left(z\right)$$
,  $\alpha_{\left(w\right)}\left(u\right)=\mathbb{E}\left[Y_{i}\left(w\right)\mid U_{i}=u\right]$ 

$$\mathbb{E}\left[\gamma_{+}\left(Z\right)Y\mid U\right] = \mathbb{E}\left[\gamma_{+}\left(Z\right)Y\cdot1\left(\left\{Z\geq c\right\}\right)\mid U\right]$$

### **Proposition 1**

Let  $\gamma_+(\cdot), \gamma_-(\cdot)$  be measurable functions of Z, then under A1-A3:

$$\mathbb{E}\left[\gamma_{+}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(1)}\left(U\right)h\left(U,\gamma_{+}\right)\right], \qquad \qquad \mathbb{E}\left[\gamma_{-}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(0)}\left(U\right)h\left(U,\gamma_{-}\right)\right]$$

where 
$$h\left(u,\gamma\right)\coloneqq\int\gamma\left(z\right)p\left(z\mid u\right)\mathrm{d}\lambda\left(z\right)$$
,  $\alpha_{\left(w\right)}\left(u\right)=\mathbb{E}\left[Y_{i}\left(w\right)\mid U_{i}=u\right]$ 

$$\mathbb{E}\left[\gamma_{+}\left(Z\right)Y\mid U\right] = \mathbb{E}\left[\gamma_{+}\left(Z\right)Y\cdot\mathbf{1}\left(\left\{Z\geq c\right\}\right)\mid U\right]$$
$$= \mathbb{E}\left[\gamma_{+}\left(Z\right)Y\left(1\right)\cdot\mathbf{1}\left(\left\{Z\geq c\right\}\right)\mid U\right]$$

### **Proposition 1**

Let  $\gamma_{+}(\cdot), \gamma_{-}(\cdot)$  be measurable functions of Z, then under A1-A3:

$$\mathbb{E}\left[\gamma_{+}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(1)}\left(U\right)h\left(U,\gamma_{+}\right)\right], \qquad \qquad \mathbb{E}\left[\gamma_{-}\left(Z\right)Y\right] = \mathbb{E}\left[\alpha_{(0)}\left(U\right)h\left(U,\gamma_{-}\right)\right]$$


where 
$$h\left(u,\gamma\right)\coloneqq\int\gamma\left(z\right)p\left(z\mid u\right)\mathrm{d}\lambda\left(z\right)$$
,  $\alpha_{\left(w\right)}\left(u\right)=\mathbb{E}\left[Y_{i}\left(w\right)\mid U_{i}=u\right]$ 

$$\mathbb{E}\left[\gamma_{+}\left(Z\right)Y\mid U\right] = \mathbb{E}\left[\gamma_{+}\left(Z\right)Y\cdot\mathbf{1}\left(\left\{Z\geq c\right\}\right)\mid U\right]$$


$$= \mathbb{E}\left[\gamma_{+}\left(Z\right)Y\left(1\right)\cdot\mathbf{1}\left(\left\{Z\geq c\right\}\right)\mid U\right]$$

$$= \mathbb{E}\left[Y\left(1\right)\mid U\right]\cdot \qquad \mathbb{E}\left[\gamma_{+}\left(Z\right)\mathbf{1}\left(\left\{Z\geq c\right\}\right)\mid U\right]$$

$$= \mathbb{E}\left[\gamma_{+}\left(Z\right)\left[U\right] = \int \gamma_{+}\left(Z\right)p\left(z\right]U\right)\mathrm{d}\lambda\left(z\right) = h\left(U,\gamma_{+}\right)$$




- A1 Sharp RD
- A2 Noisy  $Z_i$ :  $Z_i \mid U_i \sim p(\cdot \mid U_i)$
- A3 Exogeneity:  $\frac{[\{Y_i(0), Y_i(1)\} \perp Z_i] \mid U_i}{[\{Y_i(0), Y_i(1)\} \perp Z_i] \mid U_i}$



- A1 **Sharp** RD
- A2 Noisy  $Z_i$ :  $\overline{Z_i} \mid U_i \sim p(\cdot \mid U_i)$
- A3 Exogeneity:  $\overline{\left[\left\{Y_{i}\left(0\right),Y_{i}\left(1\right)\right\} \perp Z_{i}\right] \mid U_{i}}$


- No need to know G (distribution of U)
- Need to know  $p(z \mid u)$  (conditional distribution of the noise)



- A1 **Sharp** RD
- A2 Noisy  $Z_i$ :  $Z_i \mid U_i \sim p(\cdot \mid U_i)$
- A3 Exogeneity:  $[\{Y_i(0), Y_i(1)\} \perp Z_i] \mid U_i$

$$\left[\left\{Y_{i}\left(0\right),Y_{i}\left(1\right)\right\}\perp Z_{i}\right]\mid U_{i}$$

- No need to know G (distribution of U)
- Need to know  $p(z \mid u)$  (conditional distribution of the noise)
  - test-retest data, prior modelling of responses to tests, physical model of the measurement device, biomedical knowledge, etc.



- A1 **Sharp** RD
- A2 Noisy  $Z_i$ :  $Z_i \mid U_i \sim p(\cdot \mid U_i)$
- A3 Exogeneity:  $\overline{\left[\left\{Y_{i}\left(0\right),Y_{i}\left(1\right)\right\} \perp Z_{i}\right] \mid U_{i}}$

- No need to know G (distribution of U)
- Need to know  $p(z \mid u)$  (conditional distribution of the noise)
  - test-retest data, prior modelling of responses to tests, physical model of the measurement device, biomedical knowledge, etc.
  - still valid when underestimating the true noise level

### References I

Eckles, D., Ignatiadis, N., Wager, S., & Wu, H. (2020). Noise-induced randomization in regression discontinuity designs. arXiv preprint arXiv:2004.09458.

# Thank you!