

# Matrix Project On Coordinate Geometry

---

Sai Harsha Kottapalli and Abhishek Agarwal

February 14, 2019

Indian Institute of Technology Hyderabad

# Index

Problem Statement

Steps to solve the problem

Solution

Figures

# Problem Statement

---

# Problem Statement

**P** and **Q** are two distinct points on the parabola

$$\mathbf{x}^T \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \mathbf{x} - \begin{bmatrix} 4 \\ 0 \end{bmatrix} \mathbf{x} = 0$$

with parameters  $t$  and  $t_1$  respectively. If the normal at **P** passes through **Q**, then find the minimum value of  $t_1^2$ .

# Steps to solve the problem

---

# Steps

- Find the constants of the parabola

# Steps

- Find the constants of the parabola
- Find Slope of tangent

# Steps

- Find the constants of the parabola
- Find Slope of tangent
- Find slope of Normal



# Steps

- Find the constants of the parabola
- Find Slope of tangent
- Find slope of Normal
- Write eqn of Normal using slope and parametric Point

# Steps

- Find the constants of the parabola
- Find Slope of tangent
- Find slope of Normal
- Write eqn of Normal using slope and parametric Point
- Minimise as a function of parameter of point P

# Solution

---

# Solution

- Given equation:

$$\mathbf{x}^T \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \mathbf{x} - \begin{bmatrix} 4 \\ 0 \end{bmatrix} \mathbf{x} = 0$$

# Solution

- Given equation:

$$\mathbf{x}^T \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \mathbf{x} - \begin{bmatrix} 4 \\ 0 \end{bmatrix} \mathbf{x} = 0$$

- Since, 3 entries of the 2x2 matrix is 0, the parabola is of the standard form (i.e.)

$$y^2 = 4ax$$

# Solution

- Given equation:

$$\mathbf{x}^T \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \mathbf{x} - \begin{bmatrix} 4 \\ 0 \end{bmatrix} \mathbf{x} = 0$$

- Since, 3 entries of the 2x2 matrix is 0, the parabola is of the standard form (i.e.)

$$y^2 = 4ax$$

- Coefficient of  $x = 4a$

# Solution

- Given equation:

$$\mathbf{x}^T \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \mathbf{x} - \begin{bmatrix} 4 \\ 0 \end{bmatrix} \mathbf{x} = 0$$

- Since, 3 entries of the 2x2 matrix is 0, the parabola is of the standard form (i.e.)

$$y^2 = 4ax$$

- Coefficient of  $x = 4a$
- $a = -1$

## Solution Contd..

- Parametric form of points of parabola:

$$\begin{bmatrix} t^2 \\ -2t \end{bmatrix}$$



## Solution Contd..

- Parametric form of points of parabola:

$$\begin{bmatrix} t^2 \\ -2t \end{bmatrix}$$

- Differentiate the equation and substitute  $\frac{dy}{dx} = m$  (the slope of the tangent)

## Solution Contd..

- Parametric form of points of parabola:

$$\begin{bmatrix} t^2 \\ -2t \end{bmatrix}$$

- Differentiate the equation and substitute  $\frac{dy}{dx} = m$  (the slope of the tangent)
- $\frac{dx}{dx} = \begin{bmatrix} 1 & m \end{bmatrix}^T$

## Solution Contd..

- 

$$\begin{bmatrix} 1 \\ m \end{bmatrix}^T \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \mathbf{x} + \mathbf{x}^T \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ m \end{bmatrix} + \begin{bmatrix} 4 \\ 0 \end{bmatrix}^T \begin{bmatrix} 1 \\ m \end{bmatrix} = 0$$

## Solution Contd..

- 

$$\begin{bmatrix} 1 \\ m \end{bmatrix}^T \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \mathbf{x} + \mathbf{x}^T \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ m \end{bmatrix} + \begin{bmatrix} 4 \\ 0 \end{bmatrix}^T \begin{bmatrix} 1 \\ m \end{bmatrix} = 0$$

- Now, use the parametric form of  $\mathbf{x}$  (i.e.)

$$\mathbf{x} = \begin{bmatrix} t^2 \\ -2t \end{bmatrix}$$

## Solution Contd..

- 

$$\begin{bmatrix} 1 & m \end{bmatrix} \begin{bmatrix} 0 \\ -2t \end{bmatrix} + \begin{bmatrix} t^2 & -2t \end{bmatrix} \begin{bmatrix} 0 \\ m \end{bmatrix} + 4 = 0$$

## Solution Contd..

- 

$$\begin{bmatrix} 1 & m \end{bmatrix} \begin{bmatrix} 0 \\ -2t \end{bmatrix} + \begin{bmatrix} t^2 & -2t \end{bmatrix} \begin{bmatrix} 0 \\ m \end{bmatrix} + 4 = 0$$

- $-2mt - 2mt + 4 = 0$

## Solution Contd..

- 

$$\begin{bmatrix} 1 & m \end{bmatrix} \begin{bmatrix} 0 \\ -2t \end{bmatrix} + \begin{bmatrix} t^2 & -2t \end{bmatrix} \begin{bmatrix} 0 \\ m \end{bmatrix} + 4 = 0$$

- $-2mt - 2mt + 4 = 0$

- $4mt + 4 = 0$

## Solution Contd..

- 

$$\begin{bmatrix} 1 & m \end{bmatrix} \begin{bmatrix} 0 \\ -2t \end{bmatrix} + \begin{bmatrix} t^2 & -2t \end{bmatrix} \begin{bmatrix} 0 \\ m \end{bmatrix} + 4 = 0$$

- $-2mt - 2mt + 4 = 0$

- $4mt + 4 = 0$

- $m = \frac{1}{t}$



## Solution Contd..

- We know that, Two lines are perpendicular if the product of their slopes is  $-1$ .

## Solution Contd..

- We know that, Two lines are perpendicular if the product of their slopes is -1.
- Slope of normal at point **p**:  $m_N = -t$

## Solution Contd..

- We know that, Two lines are perpendicular if the product of their slopes is -1.
- Slope of normal at point **p**:  $m_N = -t$
- We calculate the equation of the line through P with slope =  $m_N$

## Solution Contd..

- Note:  $\mathbf{p} = (x_1, y_1) = (t^2, 2t)$

## Solution Contd..

- Note:  $\mathbf{p} = (x_1, y_1) = (t^2, 2t)$

- 

$$\begin{bmatrix} -m & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -m & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ y_1 \end{bmatrix}$$

## Solution Contd..

- Note:  $\mathbf{p} = (x_1, y_1) = (t^2, 2t)$

- 

$$\begin{bmatrix} -m & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -m & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ y_1 \end{bmatrix}$$

- 

$$\begin{bmatrix} t & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} t & 1 \end{bmatrix} \begin{bmatrix} t^2 \\ 2t \end{bmatrix}$$

## Solution Contd..

- Note:  $\mathbf{p} = (x_1, y_1) = (t^2, 2t)$

- 

$$\begin{bmatrix} -m & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -m & 1 \end{bmatrix} \begin{bmatrix} x_1 \\ y_1 \end{bmatrix}$$

- 

$$\begin{bmatrix} t & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} t & 1 \end{bmatrix} \begin{bmatrix} t^2 \\ 2t \end{bmatrix}$$

- 

$$\begin{bmatrix} t & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = t^3 + 2t$$

## Solution Contd..

- Now, Let  $Q: (t_1^2, 2t_1)$



## Solution Contd..

- Now, Let Q:  $(t_1^2, 2t_1)$
- Then, We have,

$$\begin{bmatrix} t & 1 \end{bmatrix} \begin{bmatrix} t_1^2 \\ 2t_1 \end{bmatrix} = t^3 + 2t$$

## Solution Contd..

- Now, Let Q:  $(t_1^2, 2t_1)$
- Then, We have,

$$\begin{bmatrix} t & 1 \end{bmatrix} \begin{bmatrix} t_1^2 \\ 2t_1 \end{bmatrix} = t^3 + 2t$$

- 

$$2t_1 + tt_1^2 = t^3 + 2t$$

## Solution Contd..

- Now, Let Q:  $(t_1^2, 2t_1)$
- Then, We have,

$$\begin{bmatrix} t & 1 \end{bmatrix} \begin{bmatrix} t_1^2 \\ 2t_1 \end{bmatrix} = t^3 + 2t$$

- 

$$2t_1 + tt_1^2 = t^3 + 2t$$

- 

$$-2(t - t_1) = t(t_1 + t)(t - t_1)$$

## Solution Contd..

- Now, Let  $Q: (t_1^2, 2t_1)$
- Then, We have,

$$\begin{bmatrix} t & 1 \end{bmatrix} \begin{bmatrix} t_1^2 \\ 2t_1 \end{bmatrix} = t^3 + 2t$$

•

$$2t_1 + tt_1^2 = t^3 + 2t$$

•

$$-2(t - t_1) = t(t_1 + t)(t - t_1)$$

•

$$t_1 = -t - \frac{2}{t}$$

## Solution Contd..

- Squaring the previous equation, we get,

$$t_1^2 = t^2 + \frac{4}{t^2} + 4$$

## Solution Contd..

- Squaring the previous equation, we get,

$$t_1^2 = t^2 + \frac{4}{t^2} + 4$$

- To solve above equation we use the property:  
Arithmetic Mean  $\geq$  Geometric Mean on  $t^2 + \frac{4}{t^2}$

## Solution Contd..

- Squaring the previous equation, we get,

$$t_1^2 = t^2 + \frac{4}{t^2} + 4$$

- To solve above equation we use the property:  
Arithmetic Mean  $\geq$  Geometric Mean on  $t^2 + \frac{4}{t^2}$
- A.M  $\geq$  G.M.  $\implies t^2 + \frac{4}{t^2} \geq 2\sqrt{4} = 4$

## Solution Contd..

- Squaring the previous equation, we get,

$$t_1^2 = t^2 + \frac{4}{t^2} + 4$$

- To solve above equation we use the property:  
Arithmetic Mean  $\geq$  Geometric Mean on  $t^2 + \frac{4}{t^2}$
- A.M  $\geq$  G.M.  $\implies t^2 + \frac{4}{t^2} \geq 2\sqrt{4} = 4$
- $t_1^2 \geq 4 + 4 = 8$



## Solution Contd..

- Squaring the previous equation, we get,


$$t_1^2 = t^2 + \frac{4}{t^2} + 4$$

- To solve above equation we use the property:  
Arithmetic Mean  $\geq$  Geometric Mean on  $t^2 + \frac{4}{t^2}$
- A.M  $\geq$  G.M.  $\implies t^2 + \frac{4}{t^2} \geq 2\sqrt{4} = 4$
- $t_1^2 \geq 4 + 4 = 8$
- $\therefore$  The Minimum possible value of  $t_1^2$  is 8.


# Figures

---

# Bounds of $t_1$ and $t_1^2$



# Normal at minimum $t_1^2$



The End