Buy Two Chocolates (easy)

Problem statement

You are in a chocolate shop where each chocolate has a price, as listed in an array `prices`. With a certain amount of money in your pocket, you aim to purchase `exactly two` chocolates.

You would like to `minimize` the combined cost of these chocolates so that you are left with as much money as possible, without going into debt.

Calculate the amount of money you'll have left after making the purchase. If it's not possible to buy two chocolates without `overspending`, return the `original amount` of money you had.

Examples

Example 1:

- **Input**: prices = **[2, 5, 3, 9]**, money = **10**
- **Expected Output**: **5**
- **Justification**: The optimal choice is to buy chocolates priced at 2 and 3, costing a total of 5. Therefore, you're left with 10 - 5 = 5.

Example 2:

- **Input**: prices = **[10, 20, 15, 5, 25, 30, 35, 40]**, money = **50**
- **Expected Output**: **35**
- **Justification**: The cheapest two chocolates has cost 5 and 10. Purchasing these costs 15 in total. Thus, you are left with 50 - 15 = 35.

Example 3:

- **Input**: prices = **[7, 1, 5, 3]**, money = **3**
- **Expected Output**: **3**
- **Justification**: It's impossible to buy two chocolates without going into debt, so you keep your original amount of 3.

Try it yourself

Try solving this question here:

Java

```java
import java.util.Arrays;

public class Solution {
    // Calculate the leftover money after buying the two cheapest chocolates
    public int maxLeftover(int[] prices, int money) {
        // ToDo: Write Your Code Here.
        return 0;
    }
}

```

Test Case | Run Code Result | Stdout

```
[10, 20, 15, 5, 25, 30, 35, 40]
50
```

Use Example Testcases | Run | Submit