Two Sum (easy)

Problem Statement

You are given an array of integers `nums` and an integer `target`. Your task is to find two distinct indices `i` and `j` such that the sum of `nums[i]` and `nums[j]` is equal to the `target`. You can assume that each input will have exactly one solution, and you may not use the same element twice.

Examples

1. **Example 1**:
   - Input: `[3, 2, 4]`, `6`
   - Expected Output: `[1, 2]`
   - Justification: `nums[1] + nums[2]` gives `2 + 4` which equals `6`.
2. **Example 2**:
   - Input: `[-1, -2, -3, -4, -5]`, `-8`
   - Expected Output: `[2, 4]`
   - Justification: `nums[2] + nums[4]` yields `-3 + (-5)` which equals `-8`.
3. **Example 3**:
   - Input: `[10, 15, 20, 25, 30]`, `45`
   - Expected Output: `[1, 3]`
   - Justification: `nums[1] + nums[3]` gives `15 + 30` which equals `45`.

Try it yourself

Try solving this question here:

Java

```java
import java.util.HashMap;
import java.util.Map;

public class Solution {
    public int[] twoSum(int[] nums, int target) {
        // TODO: Write your code here
        return new int[]{-1, -1};
    }
}

```

Test Case | Run Code Result | Stdout

```
[3, 2, 4]
6
```

Use Example Testcases | Run | Submit