

Operationalizing Community-Based Emergency Aid in Complex Crisis Environments: Field Triage, Resource-Constrained Stabilization, Disaster Risk Reduction, Psychosocial Recovery, and Technology-Enabled Responder Coordination: An Academic Review of Disaster Preparedness, Community Response, and Humanitarian Emergency Aid

Wejdan Dhafer Alshehri¹, Rana saleh alGhamdi², Afnan Abdulrahman Altoum³, Tahani Mohammed Alshehri⁴, Dehmi Mahdi Al Yami⁵, Abdulrhman Ali Almazam⁶, Ahmed Mufleh Alenazi⁷, Abdullaziz Mohammed Alkhamali⁴, Albandari Saad Alsoaib⁸, Bashaer Dabsan Albaqami⁹, Fatimah Muidh Alshehri¹⁰, Ibrahim Abdullah Al saweed¹¹, Mounira Mohammed AlQahtani³, Bodour Nasser Alsahli⁴, Maryam Abdulrahman Al-Mutairi¹²

¹ Ambulatory Care Center

² Respiratory Therapist (R.T)

³ Nursing

⁴ Social Worker

⁵ Dental Assistant

⁶ Restorative Dentistry

⁷ C.T Scan Technologist

⁸ Health Info Tech

⁹ Consultant general Pediatric

¹⁰ Pharm D

¹¹ Social services

¹² Senior specialist

Ministry of National Guard Health Affairs, Riyadh, Saudi Arabia

Abstract

First aid in complex emergencies is not simply an extension of routine treatment performed on a larger scale. It is a coordinated system of immediate actions delivered when hazards are unstable, casualties are numerous, information is incomplete, and professional resources may be delayed or overwhelmed. This paper examines community-based emergency aid in natural disasters, technological accidents, violent incidents, epidemics, displacement settings, and other mass-casualty environments. It develops a framework that connects disaster risk reduction, scene safety, rapid operational assessment, field triage, resource-constrained stabilization, referral, documentation, and psychosocial recovery. Particular attention is given to the transition from individual-centered care to population-centered prioritization when demand exceeds capacity, while recognizing the ethical risks of rationing and the need for transparent, accountable decision-making.

The paper reviews the roles of community responders, emergency medical services, health facilities, humanitarian organizations, local authorities, and affected populations. It analyzes field approaches to hemorrhage control, airway and breathing support, crush injury, burns, contamination, dehydration, hypothermia, heat illness, infectious threats, and medication continuity. It also discusses triage as a dynamic process rather than a fixed label, the importance of patient tracking and interagency communication, and the value of low-technology solutions when electricity or connectivity fails. Psychological first aid is presented as an integrated component of emergency aid, not an optional service added after physical treatment. Finally, the paper examines digital dispatch, geographic information systems, mobile applications, artificial intelligence, remote consultation, wearable sensors, and simulation-based education. The argument is that technology can improve coordination only when it is resilient, locally adapted, ethically governed, and supported by trained people. The future of disaster first aid depends on preparedness before the event, disciplined action during the acute phase, and community-centered recovery afterward.

Medical safety note: This paper is an academic review and does not replace certified first-aid training, local emergency protocols, or professional medical assessment. In a real emergency, responders should protect themselves, activate the local emergency number, follow dispatcher instructions, and avoid entering unsafe environments without appropriate training and equipment.

1. Introduction

Disasters expose the limits of ordinary emergency-care systems. A single injured person may require careful assessment, treatment, transport, and follow-up. A disaster may generate hundreds or thousands of injured, displaced, frightened, or medically dependent people within minutes. Roads may be blocked, hospitals may be damaged, communications may fail, and the number of trained responders may be insufficient. In such circumstances, first aid becomes an operational discipline that links public action, emergency services, humanitarian coordination, and health-system resilience.

Mass-casualty incidents are commonly understood as events in which the number or severity of patients overwhelms the resources immediately available to provide usual standards of individual care. The World Health Organization emphasizes that this situation requires a fundamental shift: instead of attempting to provide the best possible outcome for each person independently,

responders must identify and treat those most likely to benefit from available interventions while preserving safety and continuity of care [1]. This shift is difficult because it challenges deeply held clinical expectations and creates ethical tension between individual advocacy and population benefit.

The first minutes of a crisis are also shaped by the behavior of ordinary people. Neighbors may remove injured people from danger, apply pressure to bleeding wounds, help children locate caregivers, share medication, guide ambulances, or communicate information when formal systems are delayed. Conversely, untrained rescuers may enter unstable buildings, crowd evacuation routes, spread rumors, or move patients unnecessarily. Preparedness therefore has two dimensions: the development of technical skills and the development of a coordinated social environment in which people know what to do, what not to do, and how to connect with professional responders.

This paper adopts a systems perspective. It does not treat first aid as a list of isolated procedures. Instead, it examines how immediate care is influenced by hazard recognition, command structure, triage, logistics, public communication, patient tracking, psychological support, technology, and recovery planning. The central question is: **How can communities and response organizations deliver safe, equitable, and effective first aid when the environment is uncertain and resources are constrained?**

2. Literature Review and Conceptual Foundations

The literature on disaster first aid brings together emergency medicine, public health, humanitarian action, nursing, disaster risk reduction, psychology, logistics, and ethics. Unlike routine clinical practice, disaster response cannot be evaluated only by treatment outcomes for individual patients. It must also consider time to recognition, operational safety, resource allocation, preventable mortality, evacuation, communication, continuity of essential services, and the psychological effects on survivors and responders.

WHO has developed mass-casualty management resources that include preparedness and activation checklists, action cards, equipment checklists, patient-tracker forms, standardized clinical documentation, and integrated triage tools [1]. These resources demonstrate an important principle: response quality depends on preparation before the event. During a crisis, teams cannot

invent command relationships, patient flow, supply arrangements, and communication pathways from the beginning. They must activate structures that have been rehearsed and adapted to local hazards.

Research on mass-casualty triage indicates that widely used systems are not equally validated in all settings. A systematic review found that the accuracy, sensitivity, and specificity of several triage systems may be below the level that users intuitively expect, reinforcing the need for training, repeated assessment, and careful interpretation [2]. Triage algorithms can support consistency, but they do not eliminate uncertainty. A person's category may change as physiology, access to treatment, transport capacity, or the number of casualties changes.

The literature also distinguishes between technical capacity and social capacity. A community may possess ambulances and hospitals yet respond poorly if the public does not know how to activate EMS, if language barriers prevent communication, or if vulnerable people are excluded from planning. Conversely, a community with modest equipment may save lives through early recognition, effective bleeding control, organized evacuation, mutual aid, and clear communication. Resilience is therefore not only a property of institutions; it is produced through relationships among people, organizations, infrastructure, and information.

Psychological first aid literature emphasizes humane, practical, and non-intrusive assistance after disasters. It focuses on safety, calm communication, connection with social support, and linkage to services rather than compulsory emotional disclosure or premature diagnosis [3] [4]. This approach is particularly important in mass-casualty settings, where survivors may experience fear, grief, separation, uncertainty, and loss of control while responders themselves face exhaustion and moral distress.

Technology is increasingly present in disaster response. Geographic information systems can map hazards, shelters, roads, and health facilities. Mobile platforms can transmit patient information and request assistance. Telemedicine can connect isolated responders with clinicians. Artificial intelligence may assist demand forecasting, image interpretation, or triage support. Yet technology may fail during power outages, cyberattacks, network congestion, or misinformation campaigns. A resilient plan therefore includes analogue alternatives and treats technology as an aid to human coordination rather than a replacement for local judgment.

3. Disaster Risk Reduction and Preparedness for First Aid

3.1. Hazard Mapping and Vulnerability Analysis

Preparedness begins by identifying the hazards most likely to affect a community. Coastal areas may face flooding and storm surge; urban regions may face transport crashes, building collapse, fire, industrial incidents, heatwaves, and infectious outbreaks; rural areas may face agricultural injuries, isolation, extreme weather, and delayed transport. Hazard mapping should be connected to vulnerability analysis. A hospital may be close to a neighborhood but inaccessible during a flood. A school may have an AED but no staff trained to use it. A shelter may have food and water but no plan for insulin storage or wheelchair access.

A useful assessment identifies who may be exposed, which resources are available, which routes may fail, and what functions must continue. It should include children, older adults, people with disabilities, pregnant individuals, people with chronic disease, people who do not speak the dominant language, and people without stable housing. Preparedness that excludes vulnerable groups can create unequal survival outcomes even when the overall response appears organized.

3.2. Community Responder Networks

Community responders may include volunteers, teachers, security staff, public-transport workers, industrial personnel, faith organizations, neighborhood leaders, and trained first-aid providers. Their role is not to replace EMS or conduct unsafe rescue. Their value lies in shortening the time between injury and basic assistance, identifying urgent needs, guiding professional responders, supporting evacuation, and maintaining communication with people who may otherwise be overlooked.

A responder network should have clear boundaries. Members need to know how to activate EMS, where to assemble, which equipment is available, how to report patient numbers, and when to withdraw. Training should include physical skills, communication, personal protective equipment, ethical decision-making, psychological first aid, and practical exercises with incomplete information. Regular drills should examine not only individual performance but also whether the network can function when roads, electricity, or mobile networks are disrupted.

3.3. Emergency Supplies and Low-Technology Readiness

Emergency supplies should reflect local hazards rather than generic assumptions. A flood-prone community may need waterproof wound dressings, flotation devices, clean-water resources, thermal protection, and medication-continuity plans. An industrial area may require eye irrigation, respiratory protection, chemical references, and decontamination procedures. A sports facility may prioritize AED access, cooling equipment, bleeding-control kits, and staff training.

Low-technology readiness is essential. Printed maps, paper triage tags, battery-powered radios, whistles, flashlights, laminated action cards, manual patient logs, and physical notice boards can remain functional when digital systems fail. Supplies should be stored where they can be reached without entering danger and checked on a defined schedule. The goal is not to accumulate equipment without a plan; it is to ensure that each resource has a known purpose, a trained user, and a maintenance process.

3.4. Preparedness Exercises and Evaluation

Exercises should move beyond ceremonial demonstrations. A realistic exercise introduces a changing scenario, conflicting information, limited supplies, vulnerable patients, and communication barriers. Participants should practice switching from routine care to mass-casualty operations, establishing a triage area, opening patient-flow routes, requesting additional resources, and transferring information to hospitals.

Evaluation should produce measurable learning. Useful indicators include time to recognize the incident, time to activate EMS, time to establish command, percentage of patients receiving appropriate initial prioritization, accuracy of patient tracking, availability of critical equipment, and responder safety events. Debriefing should be constructive and should convert observations into changes in protocols, infrastructure, and training.

4. Operational Safety and the First Fifteen Minutes

4.1. Scene Safety under Dynamic Hazards

The first responder's initial task is not to touch the patient; it is to determine whether intervention is safe. Hazards may include fire, smoke, electrical current, gas, unstable structures, moving vehicles, weapons, contaminated water, aggressive crowds, radiation, biological exposure, and

aftershocks. The scene can change while care is being delivered. Responders should maintain situational awareness, use appropriate protective equipment, and withdraw when the risk becomes unacceptable.

A responder who becomes injured increases the burden on the system. This does not mean that people should ignore victims. It means that assistance must be proportional to training and conditions. A bystander may call emergency services, warn traffic, guide people away from danger, provide verbal reassurance, or support a safe evacuation without entering a collapse zone or contaminated area.

4.2. Establishing an Operational Picture

An operational picture answers several questions: What happened? Where is the hazard? How many people are involved? Which patients are immediately visible? Are more victims likely to be found? Which routes are open? What resources have arrived? What information is confirmed and what remains uncertain? The answers should be communicated through a designated channel rather than repeated through uncontrolled rumor.

The first report should be concise and action-oriented. It should include location, mechanism, major hazards, estimated patient count, obvious life threats, access limitations, and resources needed. As new information arrives, the report should be updated. Early reports are often incomplete, but delaying communication until every detail is known can cost time.

4.3. Primary Assessment in a Crisis

In a mass-casualty environment, the primary assessment remains focused on immediate threats, but it is performed rapidly and repeatedly. The responder looks for absent normal breathing, catastrophic bleeding, airway obstruction, severe respiratory distress, altered consciousness, and signs of shock. Interventions should be limited to actions that can be performed quickly and that offer substantial benefit, such as opening an airway, applying pressure to severe bleeding, or directing a walking patient to safety.

The ABCDE framework can guide individual assessment, but disaster operations add an operational layer. A responder must also ask whether treatment is possible at the current location, whether the patient should be moved to a designated zone, whether the intervention consumes

resources needed elsewhere, and how the patient will be tracked. The clinical and operational assessments should inform each other.

4.4. Communication and Command

A clear command structure reduces duplication and conflict. Even a small incident benefits from someone coordinating safety, triage, treatment, transport, logistics, and communication. Roles should be announced and recorded. Community responders should follow the incident command system used by local authorities and should avoid creating competing instructions.

Communication should use closed-loop confirmation. A request such as “We need an ambulance at the east entrance for two critical patients” should be repeated by the receiving person and confirmed when completed. Plain language is preferable to unexplained codes, especially when multiple agencies or community volunteers are present. Interpreters and accessible communication methods should be available for people with hearing, speech, visual, or language barriers.

5. Field Triage and Prioritization

5.1. The Purpose of Triage

Triage is the process of sorting patients according to urgency, expected benefit from intervention, and available resources. In routine care, triage often aims to identify who should be seen first. In a mass-casualty incident, it also determines where patients should go, what intervention should be provided immediately, and how scarce transport or treatment capacity should be used.

Triage is not a diagnosis. It is an operational decision made under uncertainty. It should be repeated as the patient’s condition and the incident change. The person who is walking may deteriorate; the person initially classified as delayed may become urgent; and a patient who was critical may stabilize after bleeding control. Reassessment is therefore a core requirement rather than an optional refinement.

5.2. Immediate, Delayed, Minor, and Expectant Categories

Many disaster systems use categories that distinguish patients requiring immediate intervention, those who can safely wait, those with minor injuries, and those whose condition or resource

demands make immediate survival unlikely. Terminology and colors differ across jurisdictions. Responders must use the local system taught during training and activated by incident command.

Operational category	General description	First-aid priority
Immediate	Life-threatening condition with a realistic possibility of benefit from rapid intervention	Control catastrophic bleeding, support airway or breathing, expedite transport
Delayed	Serious injury but temporarily stable	Protect injuries, provide reassessment, prepare for transport
Minor	Able to walk or has limited injuries	Move to a safe area, provide basic care, use as a source of appropriate assistance only when safe
Expectant or deceased, according to local policy	Survival unlikely to immediately available resources or	Follow with ethical operational guidance, signs preserve
		local and

Operational category	General description	First-aid priority
-------------------------	------------------------	-----------------------

of death are dignity,
present document
decisions

5.3. Triage Limits and Ethical Tension

Triage decisions may involve painful trade-offs. A responder may know that a patient needs advanced surgery but also know that the operating room, blood products, or transport are unavailable. Ethical practice requires transparent criteria, consistency, documentation, respect, and supervision. Decisions should not be based on social status, disability, nationality, age alone, or the perceived worth of a person's life.

A triage system should be treated as a tool for prioritizing care, not as permission to abandon patients. Those who cannot receive curative treatment immediately may still need analgesia, warmth, airway comfort, communication, family contact, and palliative support. WHO resources explicitly recognize that mass-casualty management includes patient tracking, documentation, and ethically grounded care pathways [1].

5.4. Triage in Children and Vulnerable Populations

Children may require specialized approaches because their anatomy, physiology, communication, and dependency differ from adults. A child who appears quiet may be seriously ill, while a frightened child may be difficult to assess. Triage should account for age, caregiver separation, developmental communication, and the possibility of hidden injury.

People with disabilities may have baseline behaviors or communication patterns that are mistaken for altered mental status. Mobility devices, hearing aids, medications, service animals, and personal assistants may be essential. Triage areas should be physically accessible and should avoid separating people from necessary support without a clear reason.

6. Resource-Constrained Stabilization in the Field

6.1. Hemorrhage and Damage-Control Priorities

Uncontrolled bleeding can cause preventable death rapidly. In a disaster, the first-aid response focuses on immediate external hemorrhage control rather than wound cleaning or definitive closure. Firm direct pressure, wound packing where appropriate, and tourniquet use by trained or instructed responders may be required. The intervention should be recorded and communicated during handover.

Resources may be limited, so responders should organize bleeding-control materials before an incident. Improvised materials can be contaminated or ineffective; however, in a genuine life-threatening situation, a clean available material may be better than no pressure at all. The priority is to stop bleeding while protecting the responder and preparing the patient for transport.

6.2. Airway and Breathing Support

Patients with airway obstruction, facial trauma, smoke inhalation, chest injury, or severe respiratory distress require urgent attention. First-aid providers should use maneuvers and equipment within their training and should not delay EMS while attempting advanced procedures. Positioning, removal from smoke when safe, oxygen or ventilation support by qualified personnel, and rapid transport may be more important than a prolonged examination.

An open chest wound requires careful attention to breathing deterioration. Current first-aid guidance allows several approaches depending on the wound and available dressing, including leaving it exposed to ambient air, using a clean nonocclusive dressing, or using a specialized vented seal [5]. In a disaster, the responder must also consider contamination, monitoring capacity, and the risk of tension physiology. Worsening respiratory distress demands immediate escalation.

6.3. Crush Injury and Entrapment

Crush injuries may result from collapsed buildings, machinery, landslides, or crowd compression. The visible injury may underestimate internal damage. Entrapped people should not be released casually by untrained rescuers when doing so may cause further collapse or sudden physiological deterioration. Rescue specialists and medical teams should coordinate extraction.

First-aid priorities include scene safety, airway and breathing assessment, bleeding control, protection from cold or heat, reassurance, and communication. Once released, the patient may require urgent professional assessment even if they initially appear stable. Pain, swelling, weakness, dark urine, confusion, and reduced circulation are concerning signs.

6.4. Burns, Chemical Exposure, and Decontamination

Disasters may produce thermal burns, chemical exposure, smoke inhalation, and contaminated clothing. Responders must avoid becoming exposed themselves. The person should be moved away from the source when safe, contaminated clothing managed according to local guidance, and emergency services notified about the agent when known. Decontamination areas should be separated from clean treatment areas.

Thermal burns may be cooled with clean, cool running water when feasible, while ice and oily substances should be avoided. Large, deep, electrical, chemical, airway, facial, hand, foot, genital, or circumferential burns require urgent professional care. In mass-casualty conditions, responders must balance cooling and covering the burn with the need to prevent hypothermia and maintain patient flow.

6.5. Environmental Emergencies

Floods, extreme heat, cold exposure, smoke, and prolonged sheltering create environmental threats beyond the original disaster. Heat illness may cause confusion, collapse, or seizures; hypothermia may cause clumsiness, drowsiness, and abnormal breathing. First aid includes removing the person from the exposure, replacing wet clothing, using insulation or cooling, monitoring mental status, and activating EMS for severe symptoms.

Disaster shelters should plan for hydration, sanitation, ventilation, medication storage, and protection from infectious disease. Environmental first aid is therefore connected to public health. A response that treats injuries but ignores contaminated water, overcrowding, or heat inside shelters may create additional illness.

6.6. Medication and Chronic-Disease Continuity

People with diabetes, epilepsy, asthma, heart disease, kidney disease, mental illness, or dependence on essential medications may deteriorate when supply chains are interrupted.

Preparedness plans should encourage individuals to maintain appropriate emergency information and should establish mechanisms for prescription replacement, refrigeration, dialysis, oxygen, and accessible transport.

First aid providers should not improvise medication changes. They can identify urgent symptoms, preserve medication containers, communicate the person's treatment history, and connect the person with professional services. Medication continuity is especially important in displacement camps and prolonged disasters, where the acute event may last days or weeks.

7. Community First Aid in Specific Crisis Settings

7.1. Earthquakes and Structural Collapse

Earthquake response involves aftershocks, unstable buildings, dust, broken glass, gas leaks, power lines, and multiple casualty locations. The safest community action is to move away from immediate hazards, call emergency services, help people reach assembly points, and provide basic assistance without entering unsafe structures. Trained rescuers may establish search sectors and coordinate with fire and structural specialists.

First-aid teams should expect hidden bleeding, fractures, crush injury, head trauma, and psychological distress. Water, warmth, lighting, and patient identification may be as important as dressings. Families should receive accurate information about reunification processes, and children should be protected from uncontrolled movement through dangerous areas.

7.2. Floods and Severe Storms

Flood response requires attention to drowning, contaminated water, electrical hazards, hypothermia, trauma, and displacement. Moving water should never be underestimated. First aid begins with rescue safety, not entering water without appropriate equipment. Survivors may have wounds contaminated by sewage or debris and may require medical assessment for infection and tetanus.

Shelter operations should identify people who need urgent medical care, people who require medication, people separated from caregivers, and people with mobility or communication needs. Public information should come from verified authorities because rumors about dam failure, evacuation routes, or contaminated supplies may create secondary harm.

7.3. Wildfires and Smoke Events

Wildfires create burns, smoke inhalation, heat illness, evacuation stress, and disruption of transport. People with asthma, chronic lung disease, older adults, and children may be particularly vulnerable to smoke. Responders should follow evacuation orders, use respiratory protection appropriate to the hazard, and avoid re-entering unsafe zones to retrieve belongings.

In shelters, responders should monitor breathing difficulty, chest pain, confusion, and worsening fatigue. Eye irritation and minor burns may coexist with more serious inhalation injury. A person who is hoarse, coughing severely, confused, or struggling to breathe requires urgent professional assessment.

7.4. Epidemics and Biological Emergencies

During infectious outbreaks, first aid must include infection prevention. Responders should use appropriate personal protective equipment, maintain ventilation and separation, perform hand hygiene, and avoid unnecessary contact. Symptoms such as breathing difficulty, altered consciousness, severe dehydration, or rapidly worsening illness require escalation.

Communication should reduce stigma. People who may be infected should be treated with dignity and directed toward care, testing, isolation, or public-health guidance as appropriate. First aid cannot be separated from surveillance and community engagement because early reporting may prevent wider transmission.

7.5. Violent Incidents and Ongoing Threats

In an active violence situation, responder safety and threat awareness take priority. Untrained bystanders should follow law-enforcement instructions, move away from danger, and avoid returning for equipment or additional victims. Once the scene is declared reasonably safe, bleeding control, airway support, triage, and evacuation can begin.

The psychological impact may be severe even when physical injuries are limited. Survivors may experience panic, dissociation, guilt, or fear of recurrence. First aid should combine physical protection with calm communication and referral to psychosocial services.

8. Psychosocial Recovery and Responder Well-Being

8.1. Psychological First Aid in the Acute Phase

Psychological first aid is a humane and practical response to people in distress. It begins with observing safety and urgent needs, approaching respectfully, listening without pressure, and connecting the person with family, shelter, health care, and reliable information. It does not require the responder to conduct therapy or ask the survivor to describe every detail of the event.

The principles of “look, listen, and link” provide a useful structure. Look for physical danger, urgent medical needs, separation, and severe distress. Listen to what the person says they need, using calm and simple language. Link them with practical support, trusted people, and professional services. IFRC programs emphasize early mental-health and psychosocial support, including safe spaces, help points, community engagement, and support for vulnerable groups [4].

8.2. Families, Children, and Displaced People

Disasters often disrupt social networks. A first-aid responder may help a person contact relatives, locate a child-friendly area, replace lost documents, or access shelter. These practical interventions can reduce distress and restore a sense of control. Children should receive age-appropriate explanations and should not be exposed unnecessarily to graphic scenes or rumors.

Displaced people may experience grief, cultural dislocation, discrimination, and uncertainty about housing or legal status. Support should be respectful and should not assume that all survivors want to speak about the event. Community leaders and culturally appropriate services can help identify needs that outsiders might miss.

8.3. Responder Stress and Moral Injury

Responders may experience exhaustion, fear, guilt, sadness, anger, or moral distress when they cannot provide the care they believe each patient deserves. Organizations should provide rotation, hydration, sleep opportunities, peer support, confidential counseling, and clear supervisory communication. A responder who is injured, overwhelmed, or unable to continue safely should be supported in withdrawing.

Psychological support should not be reduced to a single mandatory debriefing. Different people recover through different combinations of rest, conversation, faith, family, professional

counseling, and practical support. Persistent nightmares, severe avoidance, depression, panic, substance misuse, or thoughts of self-harm require professional assessment.

9. Data, Technology, and Coordinated Response

9.1. Patient Tracking and Documentation

Patient tracking is essential when many people move between triage areas, treatment zones, ambulances, shelters, and hospitals. A tracker should record a unique identifier, observed condition, priority category, interventions, destination, and changes in status. WHO identifies patient-tracker forms and standardized clinical documentation as tools that support communication, continuity, and resource management [1].

Digital tracking may improve speed and reduce duplicate records, but paper and verbal backups are necessary. Systems should protect confidentiality and avoid collecting information that cannot be secured. Identification methods should work for children, people without documents, people who speak different languages, and people who are unconscious.

9.2. Geographic Information and Dispatch

Geographic information systems can display hazards, road closures, ambulance locations, shelters, hospitals, water points, and population vulnerability. Dispatch centers may use these data to route resources and anticipate demand. However, maps can be outdated or inaccurate after a disaster. Field confirmation and local knowledge remain essential.

Mobile applications can guide first-aid actions, locate AEDs, share verified alerts, and connect volunteers with official coordination. Applications should be designed for low bandwidth, low battery, multiple languages, and offline use. They should show the local emergency number and make human assistance easy to access.

9.3. Artificial Intelligence and Decision Support

Artificial intelligence may help forecast casualty volumes, identify patterns in dispatch calls, support translation, prioritize supply delivery, or detect changes in patient status from connected devices. It may also assist planners in simulating evacuation routes and shelter demand. These applications should be treated as decision support rather than autonomous command.

The risks include biased data, false reassurance, automation bias, privacy violations, cyberattacks, and failure during unusual events. An algorithm trained on one city may not work in another. A system that performs well under normal conditions may fail when communication is interrupted or when populations differ from the training data. Any deployment should include validation, human review, monitoring, clear accountability, and a non-digital fallback.

9.4. Telemedicine and Remote Consultation

Telemedicine can connect isolated responders with emergency clinicians, poison specialists, mental-health professionals, or trauma teams. It may support wound assessment, triage review, medication continuity, and hospital destination decisions. Audio guidance may be more reliable than video when bandwidth is poor.

Remote consultation cannot remove the need for scene safety or physical assessment. A clinician may not see the full hazard, and a responder may not be able to perform a recommended action. Protocols should clarify consent, data security, responsibility, documentation, and what to do when the connection fails.

10. Case Studies and Applied Scenarios

10.1. Case Study One: Earthquake with Multiple Collapse Sites

Scenario. A moderate earthquake affects a densely populated district. Several buildings are damaged, roads are obstructed, and residents report trapped people, bleeding injuries, and separated children. Mobile networks are intermittent.

Risks. The principal hazards include aftershocks, unstable structures, broken gas lines, electrical wires, dust, traffic congestion, and uncontrolled crowds. Community responders should not enter damaged buildings without authorization. They should activate emergency services, establish a safe assembly point, and provide a structured report describing locations and estimated casualties.

Assessment and priorities. Walking survivors can be directed away from hazards and toward a designated safe area. People who are unresponsive, not breathing normally, or suffering catastrophic bleeding receive immediate attention when access is safe. Minor injuries should be directed to a separate area to preserve access for critical patients. Children and vulnerable adults require identification and reunification support.

First-aid actions. Responders may apply direct pressure to severe bleeding, support airway and breathing within training, protect people from cold or dust, and provide reassurance. Trapped patients require specialist rescue. Patient identifiers can be written on paper tags when digital systems fail.

Technology. A mapping platform may mark collapse sites, but local residents may provide more accurate information about blocked routes. Radios and paper maps should supplement mobile tools.

Lessons. The scenario demonstrates that community first aid must be organized geographically. The most effective responder is not necessarily the person with the most equipment, but the person who can act safely, communicate accurately, and connect local knowledge to incident command.

10.2. Case Study Two: Flooded Community with Medication Disruption

Scenario. Floodwater isolates a rural community for two days. A shelter contains people with minor wounds, hypothermia risk, diabetes, asthma, and anxiety. One older adult has become confused because essential medication is unavailable.

Risks. Contaminated water, electrical hazards, cold exposure, dehydration, infection, medication interruption, and separation from caregivers are present. The shelter is crowded and has limited communication with the nearest hospital.

Assessment and priorities. The confused person requires assessment for hypothermia, low blood glucose, infection, stroke, and medication-related causes. A person with severe breathing difficulty or altered consciousness requires urgent evacuation. People with minor injuries can be treated in a designated area while the team maintains a list of medical needs.

First-aid actions. Wet clothing is replaced, patients are insulated, wounds are cleaned and covered when safe, and hydration is managed according to ability to swallow and medical advice. Medication containers or photographs of labels are preserved for professional review. The shelter should improve ventilation, sanitation, and access routes.

Technology. A low-bandwidth telemedicine connection can support medication reconciliation. If unavailable, a paper list transmitted by radio may be sufficient. A community message board can reduce rumors and identify people needing transport.

Lessons. Disaster first aid includes continuity of chronic care. A person may not have a visible injury but may still face life-threatening deterioration because the health system has been interrupted.

10.3. Case Study Three: Industrial Chemical Release

Scenario. A chemical release occurs near a factory. Workers report eye irritation, coughing, and skin contamination. Some people attempt to approach the site to help colleagues.

Risks. The greatest immediate risk is secondary exposure to responders and the spread of contamination into vehicles, homes, and medical facilities. The public should move upwind or follow official evacuation instructions and should not enter the area without appropriate protective equipment.

Assessment and priorities. Emergency services should be notified about the substance if known, wind direction, number of exposed people, and symptoms. Decontamination and separation of clean and contaminated zones are essential. Patients with severe breathing difficulty, collapse, or altered consciousness require priority professional care after safe removal from exposure.

First-aid actions. Responders should avoid direct contact, remove contaminated clothing only according to trained procedures, irrigate eyes or skin when instructed, and prevent contaminated materials from spreading. Improvised treatment should not substitute for hazardous-materials guidance.

Technology. Chemical databases, secure radios, and hospital alerts may support the response. Digital systems should not be trusted without confirmation because incorrect substance identification can lead to dangerous advice.

Lessons. The best first aid may be to prevent additional exposure. Scene safety, information accuracy, and decontamination planning are as important as direct treatment.

10.4. Case Study Four: Transport Crash with Mass Casualties

Scenario. A highway bus crashes at night. Several passengers have bleeding injuries, some are trapped, and others are walking along the road. Traffic continues to move around the scene.

Risks. Secondary collision, fire, fuel leakage, darkness, spinal injury, and panic are present. Responders should activate police, fire, and EMS and should not enter traffic without a protected approach.

Assessment and priorities. Walking passengers are directed to a safe location and asked to remain there. Patients who are unresponsive, not breathing normally, or bleeding severely are prioritized when safe access is possible. Trapped passengers are not pulled out without rescue coordination unless there is an immediate threat such as fire.

First-aid actions. Catastrophic bleeding is controlled with direct pressure or tourniquet use by trained responders. Patients are protected from cold, monitored, and reassured. Information about seating locations, missing passengers, and possible hazards is communicated to incident command.

Technology. Location sharing, electronic passenger lists, and ambulance-dispatch data may improve coordination. Paper lists and verbal confirmation remain necessary if the network fails.

Lessons. The scenario shows that first aid is inseparable from traffic control, rescue operations, patient tracking, and communication. Treating one casualty while ignoring the wider scene may increase overall risk.

11. Best Practices for Training and System Design

Effective disaster first-aid training should combine technical knowledge, operational behavior, and emotional preparation. Learners need to practice making decisions with incomplete information and limited supplies. A course that teaches wound dressing in a quiet classroom but never addresses crowd movement, unstable hazards, patient tracking, or communication may not prepare people for a real crisis.

Training domain	Practical focus	Evaluation question
Personal and scene safety	Hazard recognition, PPE,	Did the responder avoid becoming an

Training domain	Practical focus	Evaluation question
	withdrawal decisions	additional casualty?
Rapid assessment	Recognition of breathing problems, severe bleeding, shock, and altered consciousness	Were immediately reversible threats identified quickly?
Triage	Prioritization, reassessment, documentation	Did the responder use the local system consistently?
Stabilization	Pressure, positioning, cooling, insulation, basic airway support	Were interventions safe and within competence?
Communication	EMS activation, closed-loop communication, handover	Did the message contain location, hazards,

Training domain	Practical focus	Evaluation question
		patient count, and needs?
Psychosocial care	Calm approach, responder practical support, referral	Did the preserve dignity and autonomy?
Technology and fallback	Apps, radios, tracking, paper alternatives	Could the team continue if the network failed?
Recovery learning	Debriefing, and incident review, equipment replenishment	Were lessons converted into system improvements ?

Training should use simulation with realistic constraints. Scenarios may include an interrupted power supply, a non-English-speaking survivor, a missing child, a responder injury, conflicting reports, or a rapidly changing hazard. Debriefing should examine both clinical decisions and operational behavior. Instructors should reinforce that safe withdrawal and early communication can be correct decisions even when the responder cannot provide direct treatment.

12. Ethical, Legal, and Cultural Considerations

Disaster first aid involves ethical decisions under pressure. The principle of maximizing benefit must be balanced with respect for each person. Triage should be based on clinical urgency, likely benefit from available care, and operational feasibility, not prejudice. Patients who cannot receive

immediate definitive treatment should not be treated as disposable; comfort, communication, dignity, and documentation remain important.

Consent should be sought from conscious adults whenever possible. Emergency care for an unconscious person may be justified under implied consent, but responders should still act within training and local law. Legal protections for bystanders vary by jurisdiction. Training should provide locally relevant information without promising universal immunity from liability.

Cultural humility is essential. Families may have different expectations about gender, touch, privacy, death, disability, and communication. Interpreters and community leaders can improve trust, but they should not delay life-saving care. Responders should avoid stereotyping and should ask respectful questions when time allows.

Data ethics becomes increasingly important as patient tracking and digital coordination expand. Information should be limited to what is necessary, protected from unauthorized access, and handled with special care for children, displaced people, and survivors of violence. Images should not be taken or shared merely because an incident is newsworthy.

13. Future Directions

Future disaster first-aid systems will need to address more complex hazards, including climate-related disasters, urban density, infrastructure failure, prolonged displacement, cyber disruption, and simultaneous emergencies. Preparedness should therefore combine physical first aid with public-health surveillance, climate adaptation, communication resilience, and social protection.

Research should evaluate community programs using outcomes that matter in real emergencies: time to EMS activation, survival from out-of-hospital cardiac arrest, bleeding-control performance, triage reliability, patient tracking, protection of vulnerable groups, and responder well-being. Studies should include low-resource settings because tools that require advanced technology may not be transferable globally.

Artificial intelligence and digital coordination should be evaluated prospectively. Future systems should test whether algorithms improve decisions without increasing false reassurance, delay, bias, or privacy risk. Human-centered design should include responders, patients, disabled people, local authorities, and community organizations from the beginning.

Training may become more adaptive through virtual reality, mixed-reality evacuation exercises, and data-driven simulation. However, high-tech education should be paired with low-cost physical practice. The essential question is whether a person can perform safely in a noisy, frightening, resource-limited environment—not whether the person can complete an online module.

Psychosocial recovery should be embedded from the first operational plan. Communities need safe spaces, family-reunification systems, protection services, accessible mental-health referral, and support for responders. Recovery is not complete when roads reopen; it continues while people rebuild trust, housing, income, health, and social connections.

14. Conclusion

Community-based emergency aid in complex crises requires more than the transfer of routine first-aid skills into a disaster setting. It requires a coordinated method for protecting responders, understanding hazards, communicating uncertainty, prioritizing patients, delivering high-value stabilization, tracking movement, supporting vulnerable groups, and connecting the scene with professional care. The first-aid provider operates within a system shaped by command, logistics, public behavior, infrastructure, technology, and ethics.

Field triage is central but imperfect. It must be dynamic, transparent, and repeatedly reassessed. Resource scarcity changes the operational goal from individual optimization to population benefit, but it does not remove the obligation to preserve dignity, provide comfort, document decisions, and prevent discrimination. Technology can improve mapping, communication, patient tracking, remote consultation, and planning, yet resilient systems must continue when devices, networks, and algorithms fail.

The most effective disaster first-aid model is therefore human-centered, community-based, evidence-informed, and prepared before the crisis. It combines trained responders with informed citizens, formal emergency services with local knowledge, modern digital tools with analogue backups, physical stabilization with psychological support, and acute response with long-term recovery. When these components are integrated, first aid becomes a practical foundation of disaster resilience and a vital means of reducing preventable harm in the most difficult conditions.

References

- [1] World Health Organization. (n.d.). *Mass Casualty Management*. <https://www.who.int/teams/integrated-health-services/clinical-services-and-systems/emergency-and-critical-care/mass-casualty-management>
- [2] Bazyar, J., Farrokhi, M., Salari, A., Khankeh, H. R., & Delshad, V. (2022). *Accuracy of triage systems in disasters and mass casualty incidents: A systematic review*. Journal of Emergency Medicine, Trauma and Acute Care. <https://pmc.ncbi.nlm.nih.gov/articles/PMC9078064/>
- [3] World Health Organization, War Trauma Foundation, & World Vision International. (2011). *Psychological First Aid: Guide for Field Workers*. <https://iris.who.int/handle/10665/44615>
- [4] International Federation of Red Cross and Red Crescent Societies. (n.d.). *Mental Health and Psychosocial Support*. <https://www.ifrc.org/our-work/health-and-care/community-health/mental-health-and-psychosocial-support>
- [5] Hewett Brumberg, E. K., Douma, M. J., Alibertis, K., et al. (2024). *2024 American Heart Association and American Red Cross Guidelines for First Aid*. Circulation, 150(24), e519–e579. <https://doi.org/10.1161/CIR.0000000000001281>
- [6] World Health Organization. (2018). *Basic Emergency Care: Approach to the Acutely Ill and Injured*. <https://www.who.int/publications/i/item/9789241513086>
- [7] World Health Organization. (n.d.). *Mass Casualty Management Preparedness and Activation Checklists*. <https://www.who.int/teams/integrated-health-services/clinical-services-and-systems/emergency-and-critical-care/mass-casualty-management>
- [8] World Health Organization. (n.d.). *Mass Casualty Management Action Cards and Equipment Checklists*. <https://www.who.int/teams/integrated-health-services/clinical-services-and-systems/emergency-and-critical-care/mass-casualty-management>
- [9] International Committee of the Red Cross. (n.d.). *First Aid in Armed Conflict and Other Situations of Violence*. <https://www.icrc.org/en/document/first-aid-armed-conflict-and-other-situations-violence>

- [10] Federal Emergency Management Agency. (n.d.). *Community Emergency Response Team Program*. <https://www.ready.gov/community-emergency-response-team>
- [11] Centers for Disease Control and Prevention. (n.d.). *Emergency Preparedness and Response*. <https://emergency.cdc.gov/>
- [12] United Nations Office for Disaster Risk Reduction. (2015). *Sendai Framework for Disaster Risk Reduction 2015–2030*. <https://www.undrr.org/publication/sendai-framework-disaster-risk-reduction-2015-2030>
- [13] International Federation of Red Cross and Red Crescent Societies. (2020). *World Disasters Report 2020: Come Heat or High Water*. <https://www.ifrc.org/document/world-disasters-report-2020>
- [14] European Resuscitation Council. (2021). *European Resuscitation Council Guidelines 2021*. <https://cprguidelines.eu/>
- [15] American Red Cross. (n.d.). *First Aid Steps: Perform First Aid*. <https://www.redcross.org/take-a-class/first-aid/performing-first-aid/first-aid-steps>
- [16] American Heart Association. (n.d.). *First Aid Guidelines*. <https://cpr.heart.org/en/resuscitation-science/first-aid-guidelines>
- [17] Pan American Health Organization. (2019). *Prehospital Mass Casualty Management*. <https://iris.paho.org/handle/10665.2/51484>
- [18] National Disaster Life Support Foundation. (n.d.). *Disaster Life Support Education*. <https://www.ndlsf.org/>
- [19] International Federation of Red Cross and Red Crescent Societies. (2022). *A Guide to Psychological First Aid for Red Cross and Red Crescent Societies*. <https://mhpsshub.org/resource/a-guide-to-psychological-first-aid-for-red-cross-red-crescent-societies/>

[20] World Health Organization. (n.d.). *Patient Tracker Forms for Mass Casualty Management*. <https://www.who.int/teams/integrated-health-services/clinical-services-and-systems/emergency-and-critical-care/mass-casualty-management>

[21] World Health Organization. (n.d.). *WHO Trauma Care Checklist*. <https://www.who.int/teams/integrated-health-services/clinical-services-and-systems/emergency-and-critical-care/mass-casualty-management>

[22] Bazyar, J., Farrokhi, M., & Khankeh, H. R. (2019). *Triage systems in mass casualty incidents and disasters: A review study with a worldwide approach*. *Open Access Emergency Medicine*, 11, 159–168. <https://pmc.ncbi.nlm.nih.gov/articles/PMC6390156/>

[23] American Medical Association. (2010). *Disaster and Mass Casualty Triage*. *Journal of Ethics*, 12(6). <https://journalofethics.ama-assn.org/article/disaster-and-mass-casualty-triage/2010-06>

[24] World Health Organization. (n.d.). *Emergency Care Systems for Universal Health Coverage*. <https://www.who.int/health-topics/emergency-care>