

The Contemporary Landscape of First Aid Practice: Integrating Evidence-Based Emergency Response, Rapid Assessment Strategies, Digital Technologies, and Community Resilience: An Evidence-Based Academic Review of Emergency Recognition, Immediate Care, Technology, and Community Resilience

Mohammed Salem Saeed Alqahtani¹, Saeed Munsad Alhumaid Alohtani², Adel Fahad Hulayyil Almutairi³, Faisal Khashman Marzouq Alosaimi⁴, Khalid Saleh A Alshammari⁵, Abdulaziz Hathal Mohammed Alotaibi⁵, Ziyad Mubashir Mohammed Alamri³, Saif Rashed S Alqahtani⁴, Jazaa Dughayman Hudhayrim Alharbi⁵, Fahad Naif Mathkar Almutairi⁵, Abdulelah Munawir Bin Sharid Almutairi⁵, Mohammed Eid Ali Alotaibi⁵, Abdullah Muallam Almutairi², Abdullah Ali Fahad Alshehri⁵, Ibraheem Ahmed A Al Ghamdi⁶.

¹ Technician-Operation Rooms

² Paramedic Assistant

³ Technician-Emergency Medical Services

⁴ Paramedic Technician

⁵ Health Assistant-Emergency Medical Services

⁶ Consultant-Restorative Dentistry

Ministry of National Guard Health Affairs, Saudi Arabia.

Abstract

First aid represents the immediate, accessible layer of emergency care delivered before definitive medical treatment becomes available. Although it is often associated with simple procedures, contemporary first aid encompasses structured risk assessment, early recognition of life-threatening conditions, timely activation of emergency medical services, focused interventions, psychological support, and safe transfer of responsibility to professional responders. This paper examines the development and contemporary scope of first aid from an evidence-based and community-oriented perspective. It reviews the principles that guide safe initial response, including scene assessment, primary assessment, the ABC/ABCDE framework, communication with emergency services, and the limits of lay-provider competence. It then considers the first-aid management of cardiovascular and respiratory emergencies, trauma, burns, poisoning, bites, environmental emergencies, road traffic injuries, disasters, and mass-casualty incidents. Particular attention is given to cardiopulmonary resuscitation, automated external defibrillation, choking, life-threatening bleeding, shock, head and spinal injury, and the early management of psychological distress.

The paper further explores how mobile applications, telemedicine, wearable devices, artificial intelligence, simulation, and virtual reality may improve recognition, instruction, communication, and coordination without displacing human judgment or professional emergency care. Four applied case studies demonstrate the integration of rapid assessment, prioritization, first-aid action, emergency activation, and technology-supported decision-making in realistic settings. The discussion emphasizes that first-aid systems are most effective when they combine scientifically grounded recommendations with accessible education, cultural adaptation, community participation, responder safety, and continuous quality improvement. The paper concludes that the future of first aid will depend not only on better devices and algorithms but also on stronger public confidence, equitable training, resilient communities, and ethical systems that place patient safety and human dignity at the center of every early response.

Medical safety note: This academic paper is intended for educational use and does not replace certified first-aid/CPR training, local protocols, or emergency medical advice. In an actual emergency, responders should protect themselves, activate the local emergency number promptly, and follow instructions from trained dispatchers and clinicians.

1. Introduction

Emergencies frequently begin outside hospitals, in homes, schools, workplaces, roads, public spaces, and disaster-affected communities. During the interval between the occurrence of an injury or acute illness and the arrival of professional responders, the actions of bystanders may influence survival, disability, pain, psychological recovery, and the eventual efficiency of definitive care. First aid is therefore not an isolated set of techniques but a social and clinical bridge between the onset of an emergency and the formal health-care system. Its value lies in the ability to identify danger early, prevent deterioration, provide immediate supportive care, and communicate accurate information to emergency medical services (EMS).

The modern concept of first aid has developed from organized civilian education, military experience, public-health initiatives, and advances in resuscitation science. Historical training efforts in the late nineteenth and early twentieth centuries established the idea that ordinary members of the public could learn structured emergency skills. Contemporary guidance has expanded this concept by incorporating evidence review, patient safety, trauma-informed communication, pediatric considerations, bleeding control, opioid overdose response, environmental emergencies, and psychological first aid. The 2024 guidelines jointly developed

by the American Heart Association (AHA) and the American Red Cross represent a major comprehensive update and draw on evidence evaluations coordinated through the International Liaison Committee on Resuscitation (ILCOR) [1].

The importance of first aid is also shaped by unequal access to emergency care. Rural populations, low-resource communities, people affected by conflict or natural disasters, and individuals who experience delays in transportation may depend on community responders for longer periods. Even in well-resourced systems, congestion, severe weather, multiple simultaneous incidents, and the distance between the scene and a trauma center can create critical delays. A resilient first-aid system must therefore combine individual competence with public education, effective dispatch, clear protocols, accessible equipment, and strong links to hospitals and community organizations.

This paper explores first aid as a comprehensive field that connects public participation with emergency medicine. It examines the foundations of safe response, the initial management of major medical and traumatic emergencies, the role of first aid in accidents and disasters, psychological support, technology-assisted care, training strategies, ethical considerations, and future directions. The central argument is that first aid is most effective when it is **early, safe, evidence-informed, context-sensitive, and connected to professional care**.

2. Literature Review

The literature on first aid includes clinical guidelines, systematic evidence reviews, educational research, public-health reports, disaster-response frameworks, and studies of bystander behavior. Its breadth reflects the fact that first aid is practiced by people with very different levels of training, from untrained witnesses receiving dispatcher instructions to certified responders working within a community emergency-response program. Consequently, recommendations must distinguish between what is physiologically beneficial, what is feasible for a layperson, and what may cause harm when performed without adequate training.

A central development in the field has been the movement from tradition-based instruction toward evidence-based recommendations. The AHA, American Red Cross, ILCOR, European Resuscitation Council, World Health Organization (WHO), International Federation of Red Cross and Red Crescent Societies (IFRC), and other organizations periodically review evidence and formulate guidance. The 2024 AHA/Red Cross guideline update addresses general first aid, medical and environmental emergencies, toxicology, trauma, bleeding, chest pain, stroke, opioid overdose, seizures, open chest wounds, hypothermia, frostbite, and other topics [1]. Its structure

demonstrates the contemporary expectation that first-aid recommendations should be tied to evidence certainty, feasibility, provider safety, and the need for escalation.

Resuscitation research has particularly influenced public first aid. Survival from cardiac arrest depends on rapid recognition, emergency activation, high-quality CPR, and early defibrillation when indicated. International resuscitation guidance has therefore promoted dispatcher-assisted CPR, public-access defibrillation, and wider community training. However, research also shows that training alone does not guarantee action. Fear of causing harm, uncertainty about whether the person is in cardiac arrest, concern about infection, social norms, and lack of confidence can delay intervention. Education must consequently include realistic practice, communication skills, and repeated opportunities to rehearse decisions rather than memorizing isolated procedures.

Trauma literature has similarly shifted attention toward preventable causes of death in the first minutes after injury. Uncontrolled external hemorrhage, airway compromise, impaired breathing, and hypothermia can worsen rapidly. First-aid guidance increasingly emphasizes immediate recognition of life-threatening bleeding and direct pressure, with tourniquet use or wound packing in appropriate situations by trained or instructed responders [1]. The same principle applies to road traffic collisions, workplace injuries, explosions, and mass-casualty incidents: early action should focus on threats that can be recognized and treated before hospital arrival, while avoiding unnecessary interventions that increase risk.

The literature also recognizes psychological consequences. Disasters and emergencies may produce acute stress, grief, fear, confusion, helplessness, and social disruption. Psychological first aid is generally understood as humane, supportive, and practical assistance that protects safety, reduces immediate distress, and connects people with social and professional support. IFRC describes mental-health and psychosocial support as an important component of emergency and community response, especially when delivered early and adapted to affected populations [2]. Psychological first aid is not psychotherapy and should not be used to force disclosure or diagnosis; rather, it is a compassionate first response that respects autonomy and cultural context.

Emerging research concerns digital technologies. Mobile applications may provide step-by-step prompts, location services, emergency-call functions, and access to local protocols. Telemedicine may allow a dispatcher, nurse, physician, or poison specialist to guide a responder remotely. Wearable devices may detect falls, abnormal heart rhythms, oxygen trends, or location changes. Artificial intelligence may support risk recognition and decision assistance. Nevertheless, reviews

of artificial intelligence in emergency medicine emphasize the need for external validation, transparency, data governance, monitoring for bias, and integration with professional oversight [3]. Technology can extend human capability, but it cannot remove uncertainty, replace physical assessment, or guarantee a correct decision in every context.

Overall, the literature supports a systems approach. First aid should be evaluated not only by whether an individual can perform a technique but also by whether the surrounding system enables recognition, communication, equipment access, referral, documentation, and follow-up. This systems perspective informs the sections that follow.

3. Foundations and Principles of First Aid

3.1. Definition and Scope of First Aid

First aid may be defined as the immediate assistance provided to a person who is injured or suddenly ill until professional medical care becomes available or the person reaches definitive care. The provider may be a bystander, family member, teacher, coworker, community responder, security officer, trained volunteer, or health professional acting outside a clinical facility. The scope of first aid is determined by the provider's knowledge, local law, available equipment, the nature of the emergency, and the need to avoid worsening the situation.

The purposes of first aid can be organized into four connected objectives. The first is to preserve life by recognizing and addressing immediately fatal threats such as cardiac arrest, severe airway obstruction, uncontrolled bleeding, or absent normal breathing. The second is to prevent deterioration by protecting the injured person from additional hazards, maintaining warmth, supporting an injured limb, or avoiding unsafe movement. The third is to promote recovery through appropriate comfort, reassurance, wound protection, and timely referral. The fourth is to facilitate continuity of care by communicating the mechanism of injury, observed symptoms, interventions, and changes in condition to EMS personnel.

First aid is not a substitute for diagnosis or definitive treatment. A responder should not delay EMS activation in order to complete a full assessment or search for equipment. Nor should a responder attempt invasive procedures, administer prescription medication without authorization or an established protocol, or move a person unnecessarily when spinal or pelvic injury is possible. The safest approach is to provide the highest-value intervention that can be performed competently while requesting professional guidance.

3.2. Importance of Early Intervention

The first minutes of an emergency are characterized by rapid change. A person with an obstructed airway may deteriorate within moments; a person with uncontrolled bleeding may develop shock; a person in cardiac arrest has a time-sensitive need for CPR and defibrillation; and a person with heat illness may progress to organ injury if cooling is delayed. Early intervention does not guarantee a favorable outcome, but it can preserve physiological function until definitive care is delivered.

Early action should be prioritized according to the relationship between urgency, potential benefit, and feasibility. For example, calling EMS is often more important than gathering additional history. Beginning chest compressions is more important than looking for a pulse for an extended period. Applying direct pressure to severe external bleeding is more important than cleaning the wound. Removing a person from immediate fire, traffic, electricity, or chemical exposure may be more important than performing a detailed examination at the original scene.

3.3. Evidence-Based First Aid

Evidence-based first aid integrates three elements: the best available research, clinical and public-health expertise, and the values, needs, and practical capabilities of the people involved. Evidence may be limited for some first-aid questions because emergencies are difficult to study ethically and under controlled conditions. In such cases, guidelines may rely on expert consensus, physiological reasoning, observational data, and feasibility considerations. Recommendations should therefore communicate uncertainty rather than present every action as equally proven.

Evidence-based practice also requires updating. A traditional action may be familiar but not beneficial; a newer action may be promising but insufficiently validated. The 2024 AHA/Red Cross guideline emphasizes that existing recommendations remain relevant unless specifically revised, while identifying topics with substantial new evidence or changing practice [1]. For educators, this means that course content, posters, mobile applications, and workplace protocols should be reviewed periodically rather than treated as permanent.

3.4. Safety and Scene Assessment

Scene safety is the first responsibility of any responder. A person cannot provide effective aid if they become a second victim. The responder should pause, observe traffic, fire, smoke, electricity, unstable structures, weapons, animals, chemical exposure, weather, crowd behavior, and other

hazards. Personal protective equipment such as gloves or a barrier device should be used when available, especially where blood or body fluids are present. If the scene cannot be made reasonably safe, the responder should move to a safe location, warn others, and await trained personnel.

After identifying immediate hazards, the responder should determine the number of patients and whether additional resources are needed. Multiple patients may require a different strategy from a single casualty. In a disaster, a responder should not become absorbed in one relatively stable patient while other people have immediately reversible life threats. The responder should identify an incident leader when possible and follow the local emergency plan.

A practical scene assessment asks: What happened? Is the scene safe? How many people are involved? What mechanisms or exposures are present? Does anyone require immediate life-saving action? What resources are available? The answers guide the next step: activating EMS, using protective equipment, providing an intervention, or preparing for handover.

4. Rapid Assessment and Initial Response

4.1. Primary Assessment

Primary assessment is a rapid process designed to detect immediate threats rather than establish a complete diagnosis. The responder begins by checking responsiveness and normal breathing while observing the overall appearance of the patient. A person who is unresponsive and not breathing normally requires immediate emergency activation and resuscitation according to local protocol. A person who is conscious but severely short of breath, actively bleeding, confused, cyanotic, or collapsing also requires urgent professional help.

The primary assessment should be repeated after any intervention and whenever the patient's condition changes. A normal initial assessment does not guarantee safety, because internal bleeding, poisoning, stroke, infection, or cardiac disease may initially present with subtle signs. The responder should therefore monitor responsiveness, breathing, skin color, pain, bleeding, and ability to communicate while waiting for EMS.

4.2. ABC/ABCDE Approach

The ABC or ABCDE approach provides a memorable sequence for identifying life threats. The exact wording differs by organization and context, but the underlying logic is consistent: address

airway, breathing, circulation, neurological disability, and environmental exposure while protecting the patient from further harm.

Airway assessment considers whether the person can speak, breathe, or maintain an open passage. An unconscious person may lose airway tone, while vomit, blood, swelling, or a foreign body may obstruct airflow. A responder should use only maneuvers appropriate to training and should avoid placing fingers blindly in the mouth. If trauma is suspected, unnecessary movement should be minimized while maintaining the ability to open the airway and provide life-saving care.

Breathing assessment considers normality, effort, rate, sounds, and associated signs such as cyanosis or inability to speak in full sentences. Agonal gasps are not normal breathing. A person who is unresponsive and not breathing normally should be treated according to cardiac-arrest protocol rather than observed passively.

Circulation includes looking for severe external bleeding, signs of shock, and the overall quality of the patient's response. A first aider should control life-threatening bleeding immediately with direct pressure and should use a tourniquet or wound packing when appropriate and trained or guided to do so [1].

Disability refers to neurological status, including responsiveness, new confusion, weakness, seizure, speech difficulty, or a change in behavior. A simple assessment of responsiveness and the ability to follow commands can identify the need for urgent escalation.

Exposure and environment involve looking for hidden injuries while preserving dignity and preventing heat loss or overheating. The responder should protect the patient from rain, cold ground, smoke, sun, and unnecessary exposure. The goal is not a full clinical examination but the identification of conditions that change immediate priorities.

4.3. Recognition of Life-Threatening Conditions

Recognition is often more important than technical complexity. Warning signs include unresponsiveness, abnormal or absent breathing, severe respiratory distress, chest pain, sudden facial droop or arm weakness, uncontrolled bleeding, severe allergic reaction, persistent seizure, blue or gray skin, rapidly worsening confusion, signs of stroke, severe burns, and suspected poisoning. In children and older adults, serious illness may present atypically, so a low threshold for professional assessment is appropriate.

The responder should use a pattern of “recognize, activate, intervene, reassess.” Recognition identifies a possible emergency; activation calls EMS or the relevant service; intervention provides the immediate action within competence; reassessment determines whether the person improves, deteriorates, or requires a different priority. This cycle is more reliable than performing a fixed checklist once and then stopping.

4.4. Emergency Communication and Activation of EMS

Emergency communication should be early, concise, and specific. The caller should provide the exact location, the number of patients, the apparent condition, hazards, mechanism of injury, and actions already taken. The phone should be placed on speaker when safe so the caller can begin care while following dispatcher instructions. If the location is difficult to find, another person should meet responders or provide landmarks.

A structured handover supports continuity. The responder should report what happened, when it happened, what was observed, what interventions were provided, how the patient responded, and whether there are relevant medications, allergies, or medical conditions known to the responder. In technology-supported systems, this information may be transmitted through an emergency app or digital platform, but verbal confirmation remains important when connectivity or data accuracy is uncertain.

5. Cardiovascular and Respiratory Emergencies

5.1. Cardiopulmonary Resuscitation

Cardiac arrest is a medical emergency in which the heart is no longer producing effective circulation. The person is typically unresponsive and not breathing normally. Immediate recognition and action are essential because the brain and other organs are highly sensitive to interruption of oxygen delivery. A responder should ensure scene safety, check responsiveness, call the local emergency number, obtain an AED if available, and begin CPR according to training and dispatcher instructions.

For an adult who is unresponsive and not breathing normally, high-quality CPR consists primarily of firm, rhythmic chest compressions in the center of the chest, allowing complete recoil and minimizing interruptions. Current resuscitation guidance commonly recommends a compression rate of approximately 100–120 per minute and a depth appropriate for the patient’s age and size; trained responders may add ventilations, while untrained responders may provide compression-

only CPR when they cannot or do not wish to provide breaths [4]. The priority is to start rather than delay because of uncertainty or fear of imperfect performance.

The responder should reassess only when directed by an AED, when the person shows clear signs of life, or when professional responders take over. If another trained person is present, rescuers should rotate compressions to reduce fatigue while minimizing pauses. For children and infants, the general principles remain rapid recognition, emergency activation, CPR, and AED use, but technique and compression depth must be adapted to the patient's size and local guidance.

5.2. Automated External Defibrillation

An automated external defibrillator analyzes the heart rhythm and, when indicated, delivers an electrical shock intended to terminate a shockable rhythm. Public-access AED programs can shorten the interval between cardiac arrest and defibrillation, particularly in airports, schools, sports facilities, workplaces, and other public locations. The device provides voice or visual prompts, which means that a responder does not need to interpret the electrocardiogram independently.

When an AED arrives, the responder should turn it on, expose and dry the chest as necessary, attach the pads as shown in the diagram, and follow the device prompts. No one should touch the patient during rhythm analysis or shock delivery. CPR should resume immediately when instructed. If the person is in water, the rescuer should remove them from the water and ensure the chest is sufficiently dry before pad application. Excessive concern about shaving, moving furniture, or obtaining perfect pad placement should not create a substantial delay.

AED access is not only a device issue. Effective programs require procurement, maintenance, visible placement, battery and pad replacement, staff awareness, legal and organizational support, and linkage to the EMS system. Training should address both technical use and the emotional barriers that prevent bystanders from acting.

5.3. Choking and Airway Obstruction

Choking occurs when a foreign object partially or completely obstructs the airway. A person with a mild obstruction may cough forcefully and should be encouraged to continue coughing while being observed. A person who cannot speak, breathe, or cough effectively requires immediate assistance and emergency activation. The responder should use age-appropriate back blows,

abdominal thrusts, chest thrusts, or other maneuvers according to current local training guidance, with modifications for infants, pregnancy, and individuals with large body size.

If the person becomes unresponsive, the responder should lower them safely and begin CPR. The mouth may be checked when opening the airway, but blind finger sweeps should be avoided because they may push the object deeper. After successful relief of choking, medical assessment may still be appropriate because residual injury or aspiration can occur.

5.4. Recognition of Cardiac and Respiratory Emergencies

Acute chest pain, pressure, sweating, nausea, unexplained shortness of breath, fainting, or pain radiating to the arm, jaw, or back may indicate a cardiac emergency. EMS should be activated promptly rather than relying on transport by a private vehicle. The person should rest in a comfortable position, be monitored continuously, and be reassured. In selected alert adults with nontraumatic acute chest pain, current AHA/Red Cross guidance allows consideration of chewing and swallowing aspirin when there is no known allergy or professional advice against it; this recommendation is context-dependent and should never delay EMS activation or be applied to children without appropriate clinical direction [1].

Stroke recognition requires attention to sudden facial asymmetry, arm weakness, speech disturbance, vision change, severe imbalance, or sudden confusion. The time of symptom onset or the last time the person was known to be well should be recorded. The person should not be given food, drink, or unprescribed medication. Rapid transport is critical because some treatments are time-sensitive.

Asthma, anaphylaxis, and severe respiratory distress require prompt assessment. A responder may assist the person in using their prescribed inhaler or epinephrine auto-injector according to instructions and local protocols. A person who becomes exhausted, confused, cyanotic, or unable to speak normally requires emergency escalation even if symptoms initially appear familiar.

6. Trauma and Injury Management

6.1. Bleeding and Hemorrhage Control

Life-threatening external bleeding is one of the most important conditions that a first aider can recognize and treat. Warning signs include blood that is spurting or flowing continuously, rapidly soaking clothing or dressings, pooling on the ground, or producing pallor, weakness, confusion,

or collapse. The responder should use gloves or a barrier if available, call EMS, and apply firm direct pressure with a dressing or clean material. Pressure should be maintained without repeatedly lifting the material to inspect the wound.

If bleeding continues from an extremity and the responder is trained or instructed, a commercial tourniquet may be applied according to its design and local protocol. In wounds where a tourniquet cannot be applied, such as some junctional areas, wound packing with appropriate material and sustained pressure may be considered by trained responders. Improvised methods should not be used casually because ineffective devices may delay definitive care. The time of tourniquet application should be recorded and communicated to EMS.

The purpose of first-aid bleeding control is not to make the wound clean or close it; it is to prevent exsanguination. Embedded objects should generally not be removed. A protruding object may be stabilized with dressings around it while pressure is applied to the surrounding area. The person should be kept warm and monitored for shock.

6.2. Wounds

Minor cuts and abrasions should be assessed for depth, contamination, foreign bodies, loss of function, and signs of infection. Once significant bleeding has been controlled, the wound may be gently rinsed with clean running water and covered with a sterile or clean dressing. A wound that is deep, gaping, caused by a bite, contaminated, associated with numbness or impaired movement, or located on the face, hand, genitals, or over a joint may require professional evaluation.

Puncture wounds can appear small while extending deeply. The risk of retained material, tendon injury, infection, and tetanus should be considered. Animal and human bites need medical assessment because of tissue damage and infection risk. The responder should not apply caustic chemicals or attempt to remove deeply embedded material.

6.3. Fractures, Sprains, and Dislocations

A fracture may be suspected when there is deformity, swelling, bruising, severe pain, abnormal movement, inability to bear weight, or a grating sensation. A sprain involves ligament injury, while a dislocation occurs when a joint surface is displaced. In first aid, the distinction cannot always be established reliably, and the immediate priorities are similar: protect the area, limit movement, control bleeding, apply a cold pack wrapped in cloth when suitable, and arrange medical assessment.

A deformed joint should not be forced back into position. The injured limb may be supported in the position found, using padding or a sling if the responder is trained and circulation can be monitored. Rings, watches, and tight objects should be removed early if swelling is expected and removal does not cause additional pain. Fingers and toes should be observed for color, warmth, sensation, and movement, and changes should be reported.

6.4. Head and Spinal Injuries

Head injury may involve concussion, skull fracture, intracranial bleeding, or associated neck injury. Concerning signs include loss of consciousness, worsening headache, repeated vomiting, seizure, increasing drowsiness, confusion, unequal pupils, weakness, fluid or blood from the nose or ears, or deterioration after an apparently minor impact. EMS should be activated for serious mechanisms, neurological changes, anticoagulant use, or any uncertainty about the person's condition.

The responder should avoid unnecessary movement when a spinal injury is possible, but airway and breathing take priority. A person who is unconscious and breathing normally may need positioning that maintains the airway, even when spinal injury is suspected. If vomiting or airway obstruction occurs, the responder should follow dispatcher guidance and move the person only as necessary to preserve life. Helmets should not be removed routinely by untrained responders unless required to provide CPR or manage an airway.

6.5. Shock and Traumatic Emergencies

Shock is a state in which the circulation fails to deliver sufficient oxygen to tissues. It may result from bleeding, severe allergy, cardiac disease, infection, dehydration, burns, or spinal injury. Signs can include pale or clammy skin, rapid breathing, weakness, anxiety, thirst, confusion, fainting, or deteriorating responsiveness. The person should be placed in a safe position, kept warm, reassured, and monitored while EMS is activated. Oral food or drink should generally be avoided in a seriously injured or potentially surgical patient.

The most effective first-aid response to traumatic shock is treatment of the underlying immediate threat, especially severe bleeding, airway compromise, and breathing difficulty. A responder should not elevate the legs automatically, because this may worsen pain or injury and is not appropriate in every condition. Continuous reassessment is more important than a single positioning rule.

7. Burns, Poisoning, Bites, and Environmental Emergencies

7.1. Burns

Burns may result from heat, chemicals, electricity, radiation, or friction. The first priority is to stop the burning process while protecting the responder. The person should be removed from the source when safe, and clothing or jewelry that is not stuck to the skin may be removed. Thermal burns are commonly cooled with clean, cool running water for an appropriate period according to local guidance. Ice, butter, oils, and adhesive materials should not be applied because they may worsen tissue injury or complicate later treatment.

Large, deep, electrical, chemical, facial, airway, genital, hand, foot, or circumferential burns require urgent medical assessment. Burns in infants, older adults, and people with major medical conditions may also be serious even when the surface area appears limited. The responder should cover the burn loosely with a clean non-adherent material, prevent hypothermia, and avoid breaking blisters. Chemical burns require attention to the safety of the scene and may require prolonged irrigation or specialized decontamination instructions.

7.2. Poisoning

Poisoning can occur through ingestion, inhalation, injection, skin contact, or eye exposure. The responder should move away from contaminated air or surfaces, avoid direct contact, and call EMS or the local poison center. Important information includes the substance, amount, route, time of exposure, age and weight of the person, symptoms, and packaging. The person should not be made to vomit unless specifically instructed by a poison specialist.

An unresponsive person, one with seizures, severe breathing difficulty, altered consciousness, or suspected opioid overdose requires immediate emergency activation. In suspected opioid overdose, current guidance supports activating EMS, providing CPR when the person is unresponsive and not breathing normally, and administering naloxone when available and authorized [1]. The responder should continue monitoring because recurrent respiratory depression may occur.

7.3. Animal Bites and Insect Stings

Bites can cause tissue damage, infection, allergic reaction, and exposure to diseases such as rabies depending on the animal and location. The wound should be washed with clean water and soap

when safe, bleeding should be controlled, and medical assessment should be arranged for deep wounds, bites to the face or hands, uncontrolled bleeding, signs of infection, unknown vaccination status, or possible rabies exposure. The animal should not be chased or captured if doing so creates risk; useful information may be reported to public-health authorities.

Insect stings are often local and self-limited, but anaphylaxis can develop rapidly. Difficulty breathing, swelling of the tongue or throat, widespread hives, faintness, vomiting, or sudden collapse requires immediate EMS activation and use of a prescribed epinephrine auto-injector when available. A tick should be removed as soon as possible using steady traction close to the skin, and the site should be monitored.

7.4. Heat-Related Emergencies

Heat exhaustion may present with heavy sweating, weakness, dizziness, headache, nausea, and cool or clammy skin. The person should be moved to a cooler environment, loosened from excess clothing, cooled with water, fans, or cool compresses, and observed carefully. Heat stroke is more serious and may involve altered mental status, collapse, seizures, very high body temperature, or hot skin; it is a medical emergency requiring rapid cooling and EMS activation.

Oral fluids should not be forced in a person who is confused, vomiting, or unable to swallow safely. First aid should avoid delaying transport while attempting to achieve perfect cooling. In athletic or outdoor settings, whole-body cooling may be initiated when feasible and consistent with local protocols.

7.5. Hypothermia and Cold-Related Emergencies

Hypothermia occurs when body temperature falls due to cold exposure, wetness, wind, immersion, or impaired thermoregulation. Early signs include shivering, numbness, clumsiness, and confusion; severe hypothermia may cause drowsiness, slow breathing, collapse, or cardiac arrest. The person should be removed from the cold, wet clothing replaced with dry layers, and the body insulated with blankets. Rough handling should be avoided in severe hypothermia.

Frostbite involves freezing injury to tissue. The area should be protected from further injury and refreezing, and medical assessment should be obtained. Rubbing, breaking blisters, and direct dry heat can cause additional damage. Rewarming decisions are context-dependent and should follow professional guidance, particularly when transport conditions make refreezing possible.

8. First Aid in Accidents, Disasters, and Mass-Casualty Incidents

8.1. Road Traffic Accidents

Road traffic collisions create simultaneous hazards: moving vehicles, fire, fuel, glass, unstable structures, traumatic injury, and psychological distress. A responder should not enter traffic without warning systems or a safe plan. EMS and police should be activated with precise location information, the number of vehicles, visible hazards, and suspected casualties. The responder should approach only when the scene is reasonably protected.

At the scene, the priorities are to identify people who are unresponsive, not breathing normally, severely bleeding, trapped, or in immediate environmental danger. Unless there is fire, drowning, collapse, or another imminent threat, occupants should not be pulled from vehicles unnecessarily. A conscious person should be reassured and asked not to move. A helmeted motorcyclist should not have the helmet removed by an untrained bystander unless required to provide life-saving airway management.

8.2. Natural Disasters

Earthquakes, floods, hurricanes, wildfires, heatwaves, landslides, and storms create changing hazards and may disrupt communications, electricity, water, transportation, and health-care capacity. First aid in disasters therefore begins with personal and community safety, not immediate treatment of every injury. Responders should follow evacuation instructions, avoid contaminated water, use protective equipment, and remain alert to aftershocks, unstable structures, smoke, and electrical hazards.

Disaster first aid combines immediate care with public-health measures. Wounds should be protected from contaminated environments; dehydration and heat exposure should be monitored; medications and chronic diseases should be considered; and vulnerable people should be connected with family and shelter services. Community responders should document patients and communicate with incident managers when possible.

8.3. Mass-Casualty Incidents

A mass-casualty incident occurs when the number or severity of patients exceeds the immediately available resources. In this context, the objective changes from providing a complete response to one patient toward achieving the greatest overall benefit. Responders should establish command,

request additional resources, identify hazards, and use a disaster-triage method authorized by the local system.

Immediate actions may include directing ambulatory patients to a designated area, opening access routes, separating contaminated patients, controlling severe bleeding, supporting breathing, and marking or tracking patient movement. Triage categories and colors vary by protocol and should not be improvised. START and JumpSTART are examples of systems used in some settings, but local training and incident command instructions take precedence.

8.4. Triage and Prioritization

Triage is a dynamic process, not a permanent label. A patient initially categorized as delayed may deteriorate, while a patient who appears critical may stabilize after a simple intervention. Triage should be repeated whenever resources, patient condition, or scene hazards change. The responder should prioritize people with immediately reversible threats, while recognizing that triage decisions in resource scarcity involve ethical and operational complexity.

Disaster-triage priority	General meaning	Typical first-aid focus
Immediate	Life-threatening catastrophic problem with potential benefit from intervention	Control bleeding, open airway, rapid breathing, EMS/transport coordination
Delayed	Serious but temporarily stable injury	Protect wounds, immobilize when appropriate, reassess frequently

Disaster-triage priority	General meaning	Typical first-aid focus
Minor	Walking wounded or limited injuries	Direct to a safe area, provide basic care and information
Expectant/deceased	Resources do not command	Follow incident- policy, provide dignity and comfort when possible
sed, according to local protocol	permit immediate life-saving care	

8.5. Community Emergency Response

Community emergency response depends on trust, preparation, local leadership, and equitable access to training. Community first responders may include volunteers, teachers, drivers, security personnel, industrial workers, faith-based organizations, and neighborhood groups. Their roles should be clearly defined so that they can provide immediate support without creating parallel systems that interfere with EMS.

Community programs are strongest when they include risk mapping, first-aid kits, AED placement, emergency communication plans, multilingual education, disability inclusion, and regular drills. Training should use realistic scenarios such as road crashes, cardiac arrest, severe bleeding, flood evacuation, and missing-person events. Evaluation should measure not only knowledge but also willingness to act, time to EMS activation, equipment availability, and the quality of handover.

9. Psychological First Aid

9.1. Psychological Responses to Emergencies

People respond to emergencies in diverse ways. Some become highly anxious or agitated; others become quiet, detached, confused, or unable to make decisions. Children may regress, cling to caregivers, or express distress through play and behavior rather than words. Older adults and

people with disabilities may experience additional barriers related to communication, mobility, medication, or sensory impairment. A first aider should avoid assuming that a particular reaction indicates weakness or psychiatric illness.

Acute stress reactions are often understandable responses to danger. The initial objectives are safety, calm communication, practical assistance, connection with trusted people, and access to information. The responder should respect privacy, obtain consent where possible, and avoid forcing a person to recount the event. If there is imminent risk of suicide, violence, severe confusion, or inability to care for basic needs, professional emergency services are required.

9.2. Supporting Victims and Families

Psychological first aid begins with a calm presence. The responder can introduce themselves, explain what they can do, ask about immediate needs, and provide simple factual information. Practical actions such as locating family members, obtaining water when safe, finding shelter, protecting children, or arranging medication may reduce distress more effectively than lengthy conversation. The person should be offered choices whenever possible to restore a sense of control.

Communication should be culturally respectful and free of false promises. Statements such as “You are safe right now; we will help you contact your family” are more appropriate than guaranteeing that everything will be fine. The responder should listen without judgment, recognize grief, and refer people to mental-health or social services when distress persists or becomes dangerous. IFRC emphasizes that community-based mental-health and psychosocial support may include safe spaces, help lines, community engagement, and support for vulnerable groups [2].

9.3. Psychological First Aid Principles

A practical framework for psychological first aid is to look, listen, and link. “Look” means observing safety, urgent physical needs, and signs of severe distress. “Listen” means approaching respectfully, understanding what the person needs, and communicating calmly. “Link” means connecting the person with family, community resources, shelter, health services, or specialized mental-health care. This framework is flexible and should be adapted to the setting rather than applied as a rigid script.

Psychological first aid should never replace medical assessment when physical symptoms are present. Chest pain, shortness of breath, fainting, confusion, or agitation may be caused by

medical emergencies, intoxication, hypoxia, or head injury. Emotional support and physical assessment should therefore be integrated.

9.4. Psychological Support for First Responders

First responders may experience repeated exposure to suffering, moral distress, fatigue, sleep disruption, and fear for their own safety. Organizations should normalize appropriate support without treating every emotional response as pathology. Rest periods, peer support, confidential professional services, practical supervision, and opportunities for structured debriefing can strengthen resilience. Compulsory forced disclosure immediately after an incident is not a universal solution and may be counterproductive for some people.

Responder well-being is a safety issue. Exhausted or distressed responders may miss hazards, communicate poorly, or make unsafe decisions. Emergency plans should include hydration, rotation, protective equipment, psychological support, and clear procedures for withdrawing from a scene when personal risk becomes excessive.

10. Emerging Technologies in First Aid

10.1. Mobile First-Aid Applications

Mobile applications can make first-aid information available at the point of need. Useful functions include illustrated instructions, audio prompts, emergency-call buttons, location sharing, AED mapping, medication reminders, translation, and checklists for communicating with dispatchers. An application may also help users recognize that a situation exceeds first-aid capacity and prompt them to call EMS. The strongest designs are simple, accessible under stress, usable offline, and reviewed by recognized medical authorities.

However, an application can introduce new risks. Users may spend too long navigating menus, follow advice that is not adapted to local protocols, rely on outdated content, or assume that a symptom checker can provide a diagnosis. Applications should therefore place emergency activation prominently, use plain language, support low literacy, include disability-accessible features, and display update dates and sources. Privacy design is essential when an application collects location, health, photographs, or voice recordings.

10.2. Artificial Intelligence

Artificial intelligence may support first aid through risk classification, speech recognition, image analysis, personalized instruction, and prediction of deterioration. A conversational system could help a caller describe symptoms, identify the need for EMS, translate instructions, or remind the responder to reassess breathing and responsiveness. Computer vision might support recognition of a collapsed person or severe bleeding in a monitored public space. Predictive models could help emergency systems allocate ambulances or identify high-risk callers.

These possibilities must be separated from proven clinical effectiveness. An algorithm may perform well in a development dataset yet fail when applied to a different population, language, device, lighting condition, or health system. Bias may arise when training data underrepresent children, older adults, darker skin tones, people with disabilities, rural populations, or non-English speakers. An AI system may also produce confident but incorrect advice, especially when information is incomplete.

Responsible deployment requires transparent performance evaluation, external validation, human oversight, clear escalation pathways, cybersecurity, and continuous monitoring after implementation. AI should be designed as a decision-support layer that helps people access appropriate care, not as an authority that overrides dispatchers, clinicians, or the responder's observation. In first aid, a modest and reliable prompt to activate EMS may be more valuable than an ambitious system that attempts to diagnose every condition.

10.3. Telemedicine and Remote Guidance

Telephone and video guidance can extend professional expertise to people at the scene. EMS dispatchers already provide instructions for CPR and other immediate actions in many systems. Telemedicine may add visual assessment, language interpretation, remote clinical consultation, and support for rural or isolated communities. A remote professional may help a responder identify an AED, assess bleeding control, prepare for evacuation, or document changes while an ambulance is traveling.

Remote guidance has limitations. Connectivity may fail, video may distort color or depth, the responder may not be able to position the camera safely, and a remote clinician cannot physically control the environment. The technology should therefore supplement rather than delay local EMS

activation. Systems should define who is responsible for the decision, how consent and privacy are handled, how recordings are stored, and what happens when the connection is lost.

10.4. Wearable Devices

Wearable devices can monitor heart rate, movement, temperature, oxygen saturation, or location. Fall detection may alert a contact when a person does not respond; a smartwatch may identify a possible abnormal rhythm; and a sensor may transmit data from an athlete, worker, or older adult. In disaster response, wearable identification systems may support patient tracking and reduce duplication of records.

Measurements must be interpreted cautiously. Consumer devices may be affected by motion, poor skin contact, cold extremities, device placement, skin pigmentation, or algorithmic limitations. A displayed number should never override the person's symptoms or a direct safety assessment. Wearables are most useful when they trigger human review and rapid communication rather than when they operate as autonomous diagnostic systems.

10.5. Simulation, Virtual Reality, and Digital Training

First-aid training is often limited by the fact that serious emergencies are infrequent. Simulation allows learners to practice high-consequence decisions without endangering a real patient. Scenarios can include sudden cardiac arrest, catastrophic bleeding, choking, road traffic trauma, heat stroke, disaster triage, and psychologically distressed survivors. Effective simulation includes briefing, realistic cues, hands-on performance, communication, debriefing, and reassessment.

Virtual reality and augmented reality can add environmental complexity, multiple casualties, noise, time pressure, and changing hazards. Nevertheless, technology should not replace physical practice with compressions, bandaging, tourniquet application, recovery positioning, or AED operation. A blended model combines digital knowledge acquisition with supervised hands-on training and periodic refreshers. Competence should be assessed by observable performance and decision quality, not merely by completion of an online module.

11. Case Studies and Practical Applications

11.1. Case Study 1: Road Traffic Accident with Suspected Hemorrhage

Scenario. At dusk, a two-car collision occurs on a busy road. One driver is conscious but has a deep bleeding wound in the thigh. Another occupant is unresponsive inside a vehicle. Traffic continues to pass, and a small fuel leak is visible.

Risks and assessment. The immediate risks include secondary collision, fire, uncontrolled hemorrhage, airway compromise, and spinal injury. A bystander should stop at a safe distance, activate EMS and police, warn approaching traffic if it can be done safely, and avoid entering the vehicle until hazards are controlled. The number of patients and the fuel leak should be communicated clearly.

Priorities and first aid. The unresponsive person should be assessed for normal breathing only when the scene is safe and access is possible. If the person is not breathing normally, dispatcher-guided resuscitation becomes the priority. For the conscious driver, the responder should apply firm direct pressure to the thigh wound. If trained or guided, a commercial tourniquet may be applied to life-threatening extremity bleeding. The person should be kept warm, reassured, and monitored. The responder should not remove trapped occupants or move the neck unnecessarily unless there is an immediate threat such as fire.

EMS activation and technology. The emergency call should include the road direction, landmark, number of vehicles, entrapment, fuel leak, patient count, and bleeding severity. Location sharing may help ambulances reach the scene. A dispatcher may provide CPR or bleeding-control instructions. Photographs should not be taken or shared if they compromise privacy or distract from care.

Lessons learned. This scenario demonstrates that scene safety precedes patient contact, that catastrophic bleeding is treated immediately, and that airway and breathing decisions must be integrated with the hazards of entrapment and traffic. It also shows why first aid is a systems activity involving bystanders, dispatchers, police, fire services, and EMS.

11.2. Case Study 2: Sudden Cardiac Arrest in a Public Facility

Scenario. A middle-aged person collapses in a community sports center. The person does not respond when spoken to and is making occasional abnormal gasping sounds. A wall-mounted AED is located approximately one minute away.

Risks and assessment. The scene is safe, but the person is unresponsive and not breathing normally. A bystander should point to a specific person and instruct them to call EMS and retrieve the AED. Another bystander should begin chest compressions. The gasping should not be mistaken for normal breathing.

Priorities and first aid. CPR should begin immediately, with compressions in the center of the chest and minimal interruptions. When the AED arrives, it should be turned on, pads applied to the bare chest, and the prompts followed. Everyone should stand clear during analysis and shock delivery. CPR should resume immediately when directed. If a trained responder can provide ventilations, they may do so with a barrier device; otherwise, compression-only CPR is preferable to delay.

Technology and communication. The AED is a simple decision-support device, but its effectiveness depends on maintenance, visibility, and staff familiarity. A facility may improve outcomes through an AED registry, regular drills, and an emergency action plan. The caller should provide the facility entrance and access instructions so EMS can reach the patient quickly.

Lessons learned. The critical lesson is that recognition, emergency activation, CPR, and defibrillation form one chain. The presence of an AED is valuable only when people know where it is and feel prepared to use it. The scenario also illustrates the importance of realistic refresher training to reduce hesitation.

11.3. Case Study 3: Mass-Casualty Incident after a Building Collapse

Scenario. A partial building collapse produces dust, structural instability, multiple injured people, and a crowd of uninjured witnesses. Some casualties are walking, while others are trapped or bleeding.

Risks and assessment. The hazards include further collapse, dust inhalation, electrical lines, unstable debris, and uncontrolled movement of the crowd. A trained responder should not enter

the collapse zone without appropriate rescue authorization. EMS, fire, police, and incident command must be activated. A safe assembly point should be identified.

Priorities and first aid. Walking wounded may be directed to a safe area for assessment. Responders with appropriate training may control catastrophic bleeding and support airway or breathing problems when access is safe. Patients should be prioritized according to the disaster-triage system used by the local authority. Trapped patients should not be moved by untrained people unless an immediate life threat requires it and trained rescue personnel direct the action.

Technology and coordination. A digital patient-tracking system may support identification, triage category, location, and destination. Radios or secure communication platforms can reduce duplication, but responders should maintain a paper or verbal backup if networks fail. Drones and mapping tools may help incident command assess the site, but they do not replace structural specialists.

Lessons learned. The appropriate first-aid response is not simply “treat the most visible injury.” It is a coordinated process that protects responders, creates order, applies triage, performs high-value interventions, and preserves information for receiving hospitals.

11.4. Case Study 4: Urban Flood and Psychological Distress

Scenario. After severe flooding, residents gather in a temporary shelter. Several people have minor wounds and wet clothing. A child is separated from a caregiver, and an older adult is confused about medication. Another survivor is shaking and repeatedly states that they cannot cope.

Risks and assessment. The first-aid team should assess environmental safety, hypothermia, contaminated water, infection risk, medication interruption, dehydration, and safeguarding concerns. The child should be connected with trained safeguarding personnel rather than informally handed to an unknown adult. The confused older adult requires medical and medication assessment because confusion may indicate hypothermia, hypoxia, infection, or another acute illness.

Priorities and first aid. Wet clothing should be replaced, people insulated, wounds cleaned and covered when safe, and urgent medical symptoms escalated. The distressed survivor should be approached calmly, offered a safe space, asked about immediate needs, and connected with trusted

people and psychosocial services. The responder should not force a detailed account of the flood or promise that the consequences will disappear.

Technology. A shelter-management system may help locate family members, record medical needs, and coordinate transport. SMS alerts can provide verified information, but rumor control is essential. Telemedicine may support medication reconciliation where clinicians are not physically present.

Lessons learned. Disaster first aid includes physical, psychological, social, and information needs. A technically correct wound dressing is insufficient if people remain unsafe, separated from caregivers, unable to access medication, or excluded from reliable communication.

12. Best Practices and Training Strategies

First-aid education should be designed around decisions and actions rather than memorization alone. Learners should repeatedly practice recognizing danger, calling EMS, assessing responsiveness and breathing, controlling severe bleeding, using an AED, managing choking, positioning a patient safely, and communicating a concise handover. Scenarios should include uncertainty, distractions, language differences, bystander disagreement, and limited equipment because real emergencies rarely occur in ideal classroom conditions.

Training component	Purpose	Quality indicator
Knowledge preparation	Introduce principles, recognition, escalation	Learner can identify life threats and explain when EMS is required
Hands-on skills	Develop physical performance	Learner safe demonstrates CPR, AED use, bleeding control, and basic injury care

Training component	Purpose	Quality indicator
Scenario simulation	Integrate assessment, communication, and prioritization	Learner acts under time pressure and adapts when the situation changes
Debriefing	Convert experience into learning	Learner identifies what was done well and what should change
Refresher practice	Reduce decay hesitation	Performance skill remains acceptable at scheduled intervals
Community evaluation	Measure system outcomes	Improved response time, AED use, EMS activation, and handover quality

Training should be inclusive. Materials should be available in relevant languages and formats, and examples should represent different ages, body types, disabilities, and cultural settings. Programs should address the needs of people who may be excluded from conventional courses, including rural residents, migrants, low-income communities, and people with limited internet access. Certification should be connected to realistic competence rather than treated as permanent proof of readiness.

Organizations should maintain first-aid kits, AEDs, emergency contact lists, evacuation maps, and incident-reporting procedures. Equipment should be inspected, accessible, clearly labeled, and appropriate to the hazards of the setting. Workplaces should review incidents and near misses without a punitive focus, using findings to improve layout, training, staffing, and communication.

13. Challenges, Ethical Considerations, and Safety

A major challenge is the gap between public need and public readiness. Many people may recognize that first aid is important but still hesitate because they fear legal consequences, infection, criticism, or causing injury. Education should explain that doing nothing can also be harmful and that responders should act within their training, follow dispatcher guidance, and stop when professional help takes over. Legal protections vary by jurisdiction; therefore, local guidance should be included rather than making universal legal claims.

Equity is another concern. Access to training, AEDs, reliable mobile networks, transportation, and emergency services is not evenly distributed. A technology-centered model may unintentionally widen disparities if it assumes smartphones, data plans, literacy, or continuous electricity. Community programs should prioritize high-risk and underserved areas and provide non-digital alternatives.

Consent and autonomy matter even in emergencies. A conscious person should be told what the responder intends to do and given the opportunity to agree or decline when possible. A person who refuses care should be respected unless there is a legally defined exception or immediate incapacity. Privacy should be protected by limiting photography, social-media sharing, and unnecessary disclosure of health information.

Technology creates additional ethical questions. Who is responsible when an AI recommendation is wrong? How should data be stored? Can a model be trusted across languages and populations? What happens when an app fails or gives outdated guidance? Governance should require clear accountability, auditability, user notification, cybersecurity, and an accessible path to human help. The principle of “do no avoidable harm” applies to algorithms as much as to physical interventions.

Responder safety must remain central. A first aider is not required to enter fire, active violence, unstable structures, toxic atmospheres, or unsafe water without appropriate training and equipment. The public should be encouraged to help, but not to undertake rescue tasks that exceed their capability. Ethical first aid balances the duty to assist with the duty to avoid creating additional casualties.

14. Future Directions

The future of first aid will likely be shaped by the interaction of public education, emergency-dispatch innovation, data systems, and community resilience. Research should examine which training models produce sustained willingness to act, not only short-term knowledge gains. Studies should include diverse populations and measure outcomes such as time to recognition, time to CPR, quality of bleeding control, EMS activation, patient experience, and responder safety.

Artificial intelligence requires prospective evaluation in real settings. Future work should compare AI-supported tools with standard dispatch and educational approaches, assess false reassurance and unnecessary escalation, and test performance across languages, devices, age groups, skin tones, disabilities, and rural environments. AI should be embedded in accountable systems with human review and clear fallback procedures.

Simulation may become more adaptive through virtual and augmented reality, but research should establish whether immersive technology improves real-world performance and retention compared with lower-cost methods. Training systems should focus on transfer: can the learner act correctly in a noisy, emotionally difficult, resource-limited emergency?

Climate change and urbanization will increase the relevance of heat illness, floods, wildfire smoke, infectious threats, displacement, and mass-casualty preparedness. First-aid programs should integrate climate adaptation, public-health communication, accessible shelter, medication continuity, and psychological support. Community responders may become increasingly important when health-care facilities are overwhelmed or transport routes are disrupted.

Future first-aid systems should also become more person-centered. Patients and families should be treated as partners in information exchange, cultural interpretation, and decision-making. Disability-inclusive design, multilingual instructions, trauma-informed communication, and protection of children and vulnerable adults should be built into emergency plans rather than added after an incident.

15. Conclusion

First aid is a critical component of the emergency-care continuum. Its purpose is not to reproduce hospital medicine at the scene but to provide safe, timely, and high-value actions during the interval before professional care becomes available. Effective first aid begins with scene safety

and rapid assessment, identifies life-threatening conditions, activates EMS early, delivers appropriate interventions such as CPR, AED use, bleeding control, airway support, cooling, or protection from further injury, and communicates clearly during handover.

The field has expanded beyond isolated technical skills. Contemporary first aid includes trauma and environmental care, disaster triage, psychological first aid, community preparedness, technology-assisted communication, ethical reasoning, and responder well-being. The evidence supports interventions that are practical, teachable, and connected to local emergency systems. It also warns against overconfidence, unnecessary procedures, unsupported technologies, and the assumption that a device or algorithm can replace human judgment.

The most promising future is therefore a human-centered and evidence-informed model. Mobile tools, telemedicine, wearables, AI, and simulation may improve recognition and coordination, but they must be validated, equitable, private, and governed by professional oversight. Strong communities will combine these tools with repeated hands-on education, accessible equipment, trusted communication, inclusive planning, and respect for patient dignity. When these elements work together, first aid becomes more than an emergency reaction: it becomes a practical expression of community resilience and a dependable first link in the chain of survival.

References

- [1] Hewett Brumberg, E. K., Douma, M. J., Alibertis, K., Charlton, N. P., Goldman, M. P., Harper-Kirksey, K., Hawkins, S. C., et al. (2024). *2024 American Heart Association and American Red Cross Guidelines for First Aid*. *Circulation*, 150(24), e519–e579. <https://doi.org/10.1161/CIR.0000000000001281>
- [2] International Federation of Red Cross and Red Crescent Societies. (n.d.). *Mental health and psychosocial support*. <https://www.ifrc.org/our-work/health-and-care/community-health/mental-health-and-psychosocial-support>
- [3] Mikhail, D., et al. (2023). *Artificial Intelligence in Emergency Medicine: Viewpoint of Current Applications and Future Directions*. *Journal of Medical Internet Research* / PMC record. <https://pmc.ncbi.nlm.nih.gov/articles/PMC10245226/>
- [4] American Heart Association. (2025). *CPR and ECC Guidelines*. <https://cpr.heart.org/en/resuscitation-science/cpr-and-ecc-guidelines>

- [5] International Liaison Committee on Resuscitation. (n.d.). *ILCOR Publications*. <https://ilcor.org/publications>
- [6] American Red Cross. (n.d.). *First Aid Steps: Perform First Aid*. <https://www.redcross.org/take-a-class/first-aid/performing-first-aid/first-aid-steps>
- [7] American Red Cross. (n.d.). *How to Perform CPR: Adult CPR Steps*. <https://www.redcross.org/take-a-class/cpr/performing-cpr/cpr-steps>
- [8] American Heart Association. (n.d.). *Part 8: First Aid*. <https://cpr.heart.org/en/resuscitation-science/first-aid-guidelines/first-aid>
- [9] World Health Organization. (2011). *Psychological First Aid: Guide for Field Workers*. WHO. <https://iris.who.int/handle/10665/44615>
- [10] World Health Organization. (2018). *Basic Emergency Care: Approach to the Acutely Ill and Injured*. WHO and ICRC. <https://www.who.int/publications/i/item/9789241513086>
- [11] International Federation of Red Cross and Red Crescent Societies. (2022). *A Guide to Psychological First Aid for Red Cross and Red Crescent Societies*. <https://mhpsshub.org/resource/a-guide-to-psychological-first-aid-for-red-cross-red-crescent-societies/>
- [12] World Health Organization. (n.d.). *Prehospital Toolkit*. <https://www.who.int/teams/integrated-health-services/clinical-services-and-systems/emergency-and-critical-care/prehospital-toolkit>
- [13] World Health Organization. (2023). *Global Status Report on Road Safety 2023*. <https://www.who.int/teams/social-determinants-of-health/safety-and-mobility/global-status-report-on-road-safety-2023>
- [14] Centers for Disease Control and Prevention. (n.d.). *Heat and Health*. <https://www.cdc.gov/heat-health/>
- [15] Centers for Disease Control and Prevention. (n.d.). *Preventing Carbon Monoxide Poisoning*. <https://www.cdc.gov/carbon-monoxide/about/index.html>

- [16] National Institute for Health and Care Excellence. (n.d.). *Major trauma: assessment and initial management*. <https://www.nice.org.uk/guidance/ng39>
- [17] European Resuscitation Council. (2021). *European Resuscitation Council Guidelines 2021*. <https://cprguidelines.eu/>
- [18] American Heart Association. (2020). *2020 American Heart Association Guidelines for CPR and ECC*. Circulation. <https://doi.org/10.1161/CIR.0000000000000918>
- [19] International Committee of the Red Cross. (n.d.). *First Aid in Armed Conflict and Other Situations of Violence*. <https://www.icrc.org/en/document/first-aid-armed-conflict-and-other-situations-violence>
- [20] United Nations Office for the Coordination of Humanitarian Affairs. (n.d.). *Humanitarian Civil-Military Coordination and Emergency Response Resources*. <https://www.unocha.org/>
- [21] World Health Organization. (2019). *WHO Guidelines for the Health Sector Response to Child Maltreatment*. <https://www.who.int/publications/i/item/9789241550147>
- [22] JAMA Network Open. (2024). *Artificial Intelligence for Emergency Care Triage—Much Promise, but Important Questions Remain*. <https://jamanetwork.com/journals/jamanetworkopen/fullarticle/2818391>