

Assignment: Probability

T.Sai Raghavendra - FWC22087

13.4.3 ¹Let X represent the difference between the number of heads and the number of tails obtained when a coin is tossed 6 times. What are possible values of X?

Solution:

Variable	Value	Description
n	6	Number of trails
p_1	$\frac{1}{2}$	Probability of getting a head
q_1	$1 - p_1$	Probability of not getting a head
p_2	$\frac{1}{2}$	Probability of getting a tail
q_2	$1 - p_2$	Probability of not getting a tail
X_1	$\{0, 1, 2, 3, 4, 5, 6\}$	No.of heads in 6 tosses of a coin
X_2	$\{0, 1, 2, 3, 4, 5, 6\}$	No.of tails in 6 tosses of a coin

Table 13.4.3.2: Variable description.

(a) Number of heads in 6 tosses of a coin.

$$p_{X_1}(k) = \begin{cases} {}^nC_k p_1^k q_1^{n-k} & 0 \leq k \leq 6 \end{cases} \quad (13.4.1.1)$$

(b) Number of tails in 6 tosses of a coin.

$$p_{X_2}(k) = \begin{cases} {}^nC_k p_2^k q_2^{n-k} & 0 \leq k \leq 6 \end{cases} \quad (13.4.2.2)$$

Thus, the desired outcome is

$$X = X_1 - X_2 \quad (13.4.2.3)$$

¹Read question numbers as (CHAPTER NUMBER).(EXERCISE NUMBER).(QUESTION NUMBER)

$$X(6H, 0T) = |6 - 0| = 6 \quad (13.4.2.4)$$

$$X(5H, 1T) = |5 - 1| = 4 \quad (13.4.2.5)$$

$$X(4H, 2T) = |4 - 2| = 2 \quad (13.4.2.6)$$

$$X(3H, 3H) = |3 - 3| = 0 \quad (13.4.2.7)$$

$$X(2H, 4T) = |2 - 4| = 2 \quad (13.4.2.8)$$

$$X(1H, 5T) = |1 - 5| = 4 \quad (13.4.2.9)$$

$$X(0H, 6T) = |0 - 6| = 6 \quad (13.4.2.10)$$

Thus, the possible values of X are 0,2,4 and 6.