1

Assignment 3

Sairam V C Rebbapragada

Abstract—This document explains the concepts of Matrix multiplication, Matrix Addition and Matrix Inverse by solving a problem.

Download the python code from

https://github.com/Sairam13001/AI5006/blob/master/Assignment_3/assignment_3.py

and latex-tikz codes from

https://github.com/Sairam13001/AI5006/blob/master/Assignment 3/assignment 3.tex

1 Problem

$$\mathbf{A} = \begin{pmatrix} 3 & 1 \\ -1 & 2 \end{pmatrix} \tag{1.0.1}$$

Show that

$$\mathbf{A}^2 - 5\mathbf{A} + 7\mathbf{I} = 0. \tag{1.0.2}$$

Hence find A^{-1} .

2 Explanation

Square of a matrix is the product of matrix with itself:

$$\mathbf{A}^2 = \mathbf{A}\mathbf{A} \tag{2.0.1}$$

Product of a scalar with a matrix is the product of that scalar with every element of the matrix:

$$k\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} ka & kb \\ kc & kd \end{pmatrix} \tag{2.0.2}$$

Inverse of a matrix A is defined as:

$$\mathbf{A}\mathbf{A}^{-1} = \mathbf{I} \tag{2.0.3}$$

3 Solution

$$A^2 - 5A + 7I (3.0.1)$$

$$= \begin{pmatrix} 8 & 5 \\ -5 & 3 \end{pmatrix} - 5 \begin{pmatrix} 3 & 1 \\ -1 & 2 \end{pmatrix} + 7 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$
 (3.0.2)

$$= \begin{pmatrix} 8 & 5 \\ -5 & 3 \end{pmatrix} - \begin{pmatrix} 15 & 5 \\ -5 & 10 \end{pmatrix} + \begin{pmatrix} 7 & 0 \\ 0 & 7 \end{pmatrix}$$
 (3.0.3)

$$= \begin{pmatrix} 8 & 5 \\ -5 & 3 \end{pmatrix} - \begin{pmatrix} 8 & 5 \\ -5 & 3 \end{pmatrix} = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$$
 (3.0.4)

Thus, It is proved that

$$\mathbf{A}^2 - 5\mathbf{A} + 7\mathbf{I} = 0. {(3.0.5)}$$

Multiplying above equation with A^{-1} on both sides, We get :

$$\implies \mathbf{A}^2 \mathbf{A}^{-1} - 5\mathbf{A}\mathbf{A}^{-1} + 7\mathbf{I}\mathbf{A}^{-1} = 0$$
 (3.0.6)

$$\implies$$
 AAA⁻¹ – 5**I** + 7**A**⁻¹ = 0 (3.0.7)

$$\implies \mathbf{AI} - 5\mathbf{I} + 7\mathbf{A}^{-1} = 0 \tag{3.0.8}$$

$$\implies \mathbf{A}^{-1} = \frac{1}{7} (5\mathbf{I} - \mathbf{A})$$
(3.0.9)

Solving for A^{-1} , we get:

$$\mathbf{A}^{-1} = \frac{1}{7} \left(5 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} - \begin{pmatrix} 3 & 1 \\ -1 & 2 \end{pmatrix} \right)$$
 (3.0.10)

$$\implies \mathbf{A}^{-1} = \frac{1}{7} \left(\begin{pmatrix} 2 & -1 \\ 1 & 3 \end{pmatrix} \right) \tag{3.0.11}$$