Assignment 11

Sairam V C Rebbapragada

Abstract—This document uses the concepts of matrix inverse, trace and determinant in solving a problem.

Download Python code from

https://github.com/Sairam13001/AI5106/blob/main/ Assignment 11/assignment 11.py

Download latex-tikz codes from

https://github.com/Sairam13001/AI5106/blob/main/ Assignment 11/assignment 11.tex

1 Problem

Let $P_A(x)$ denote the characteristic polynomial of a matrix A, then for which of the following matrices $P_A(x) - P_{A^{-1}}(x)$ a constant?

3)
$$\begin{pmatrix} 3 & 2 \\ 4 & 3 \end{pmatrix}$$

4)
$$\begin{pmatrix} 2 & 3 \\ 3 & 4 \end{pmatrix}$$

2 Explanation

The characteristic polynomial of a matrix A is defined as

$$P_A(x) = det(xI - A) \tag{2.0.1}$$

3 Solution

Let matrix A be

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \tag{3.0.1}$$

$$\implies P_A(x) = det(xI - A) \tag{3.0.2}$$

$$= det \begin{pmatrix} x - a & -b \\ -c & x - d \end{pmatrix}$$
 (3.0.3)

$$= x^2 - (a+d)x + (ad - bc)$$
 (3.0.4)

$$= x^2 - tr(A)x + det(A)$$
 (3.0.5)

We know that for a 2×2 matrix, the inverse is calculated as:

$$A^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$$
 (3.0.6)

$$\implies P_{A^{-1}}(x) = det(xI - A^{-1})$$
 (3.0.7)

$$= x^2 - tr(A^{-1})x + det(A^{-1})$$
 (3.0.8)

So, $P_A(x) - P_{A^{-1}}(x)$ results in

$$(tr(A^{-1}) - tr(A))x + det(A) - det(A^{-1})$$
 (3.0.9)

It can be observed that $P_A(x) - P_{A^{-1}}(x)$ will be constant when

$$tr(A^{-1}) - tr(A) = 0 (3.0.10)$$

$$\implies tr(A^{-1}) = tr(A) \tag{3.0.11}$$

$$\implies \frac{a+d}{\det(A)} = a+d \tag{3.0.12}$$

(3.0.12) holds when either a + d = 0 or det(A) = 1. It is easy to observe that the correct answer among the given options is $\begin{pmatrix} 3 & 2 \\ 4 & 3 \end{pmatrix}$ as its determinant is 1.