



Logistic Regression

Quiz, 5 questions

Congratulations! You passed!

Next Item


1 / 1
point

1.
Suppose that you have trained a logistic regression classifier, and it outputs on a new example x a prediction $h_\theta(x) = 0.4$. This means (check all that apply):

1 / 1
point

2.
Suppose you have the following training set, and fit a logistic regression classifier $h_\theta(x) = g(\theta_0 + \theta_1 x_1 + \theta_2 x_2)$.

x_1	x_2	y
1	0.5	0
1	1.5	0
2	1	1
3	1	0



Which of the following are true? Check all that apply.

0 / 1
point

3.
For logistic regression, the gradient is given by $\frac{\partial}{\partial \theta_j} J(\theta) = \frac{1}{m} \sum_{i=1}^m (h_\theta(x^{(i)}) - y^{(i)}) x_j^{(i)}$. Which of these is a correct gradient descent update for logistic regression with a learning rate of α ? Check all that apply.

1 / 1
point

- 4.

Which of the following statements are true? Check all that apply.



Logistic Regression

Quiz, 5 questions



1 / 1
point

5.
Suppose you train a logistic classifier $h_{\theta}(x) = g(\theta_0 + \theta_1 x_1 + \theta_2 x_2)$. Suppose $\theta_0 = 6, \theta_1 = 0, \theta_2 = -1$. Which of the following figures represents the decision boundary found by your classifier?

