

Math 11, Fall 2007

Lecture 16

Scott Pauls¹

¹Department of Mathematics
Dartmouth College

10/31/07

Outline

- 1 Review and overview
 - Last class
- 2 Today's material
 - Integration in two variables in polar coordinates
- 3 Group Work
- 4 Next class

Outline

- 1 Review and overview
 - Last class
- 2 Today's material
 - Integration in two variables in polar coordinates
- 3 Group Work
- 4 Next class

Integration of a function of two variables

General domains

- Iterated integrals and Fubini's theorem

$$\int \int_R f(x, y) \, dA = \int_a^b \int_c^d f(x, y) \, dx \, dy$$

- Non-rectangular domains: parameterize boundary and introduce variables into the bounds of integration

Outline

- 1 Review and overview
 - Last class
- 2 Today's material
 - Integration in two variables in polar coordinates
- 3 Group Work
- 4 Next class

Polar coordinates

- $(x, y) \rightarrow (r, \theta)$
- $x = r \cos(\theta), y = r \sin(\theta)$
- $r^2 = x^2 + y^2, \theta = \tan^{-1}(y/x)$

Change of variables

If f is a continuous function defined on a polar rectangle
 $R = [a, b] \times [\alpha, \beta] = \{(r, \theta) | a \leq r \leq b, \alpha \leq \theta \leq \beta\}$ where
 $0 \leq \beta - \alpha \leq 2\pi$ then

$$\int \int_R f(x, y) \, dA = \int_{\alpha}^{\beta} \int_a^b f(r \cos(\theta), r \sin(\theta)) \, r \, dr \, d\theta$$

General change of variables

If we are trying to integrate f over a region D , we may always change to polar coordinates:

$$\int \int_D f \, dA = \int \int_{D^*} f \, r \, dr \, d\theta$$

where D^* is the same region of D , described with respect to the polar variables.

Examples

1

$$\iint_D (1 - x^2 - y^2) dA$$

where D is the circle of radius 2 centered at the origin.

2

Let $R = \{(x, y) | 1 \leq x^2 + y^2 \leq 4, 0 \leq y \leq x\}$ and find

$$\iint_R \tan^{-1}(y/x) dA$$

3

Find the volume of a sphere of radius a centered at the origin.

4

Find

$$\int_0^1 \int_0^{\sqrt{1-x^2}} e^{x^2+y^2} dy dx$$

Examples

A cylindrical drill with radius r_1 is used to bore a hole through the center of a solid ball of radius r_2 . Find the volume of the ring-shaped solid that remains. Express the volume in terms of the height, h , of the ring.

Work for next class

- Reading: 16.6
- f07hw17