

Eigenfunctions of the Laplacian: quantum ergodicity, chaos, and fast numerical methods

October 12, 2004

Alex Barnett

Courant Institute, NYU

Thanks to Peter Sarnak, Percy Deift, Jonathan Goodman...


Dirichlet eigenproblem

Normal modes of elastic membrane or ‘drum’

(150 yr old)

Eigenfunctions ϕ_j of Laplacian in bounded domain $\Omega \subset \mathbb{R}^2$

$$-\Delta\phi_j = E_j\phi_j, \quad \phi_j|_{\partial\Omega} = 0 \quad \langle\phi_i, \phi_j\rangle = \delta_{ij}$$



cavity modes,
quantum levels

‘level’ $j = 1 \dots \infty$

‘energy’ eigenvalue E_j



wavelength = $2\pi/E_j^{1/2}$

MOVIE $j \approx 3000$


- What happens to ϕ_j in the $E_j \rightarrow \infty$ (short wavelength) limit?
- How best to numerically calculate many modes efficiently?

Modern applications

- electromagnetic modes of waveguide ($\partial\Omega$ = metallic wall)
- acoustic resonances and duct transmission (Neumann BCs)
- spectral geometry *e.g.* Kac's drum problem, Riemann surfaces
- physics *e.g.* model cold electrons in ‘quantum dots’




quantum dots (Marcus)




resonant liquid surfaces (Kudrolli)

Dielectric micro-cavity lasers

quantum-cascade laser



mode and emission pattern




- 2D confinement due to total internal reflection, asymmetric cavity
- ‘scarred’ modes give power output $\times 10^3$ (*Gmachl et al ’98*)
- design & modeling: need compute many cavity modes (*Tureci ’03*)

Outline

- Background: ‘billiards’ and ‘quantum chaos’
- Quantum ergodicity
- Numerical results on quantum ergodicity
- Numerics: analysis of the ‘Method of Particular Solutions’
- Accelerated ‘scaling method’


Connection to dynamical systems

Drum problem is *quantized* equivalent of ‘billiards’
classical system: point particle, elastic reflection from $\partial\Omega$
phase space = (x, y, θ)




Connection to dynamical systems

Drum problem is *quantized* equivalent of ‘billiards’
classical system: point particle, elastic reflection from $\partial\Omega$
phase space = (x, y, θ)




Integrable:
conserved
quantities


classical




Ergodic:
covers all
phase space



eigenfunctions ϕ_j : ‘quantum’




localization (tori in phase space: EBK)



‘quantum chaos’ (Einstein 1917 realised!)


Connection to dynamical systems

Drum problem is *quantized* equivalent of ‘billiards’
classical system: point particle, elastic reflection from $\partial\Omega$
phase space = (x, y, θ)




Integrable:
conserved
quantities


classical




Ergodic:
covers all
phase space



eigenfunctions ϕ_j : ‘quantum’



localization (tori in phase space: EBK)



‘quantum chaos’ (Einstein 1917 realised!)

- So, shape of domain Ω is crucial to behaviour in $E \rightarrow \infty$ limit

Quantum ergodicity

If the domain Ω is ergodic, does the intensity ϕ_j^2 become *equidistributed* (uniform across space) as $E_j \rightarrow \infty$?

Quantum ergodicity

If the domain Ω is ergodic, does the intensity ϕ_j^2 become *equidistributed* (uniform across space) as $E_j \rightarrow \infty$?

More precisely, pick any test function $A = A(\mathbf{r})$, $\mathbf{r} \in \Omega$

- **classical** phase space average $\bar{A} = \frac{1}{\text{vol}(\Omega)} \int_{\Omega} A(\mathbf{r}) d\mathbf{r}$
- **quantum** intensity average

$$\int_{\Omega} A(\mathbf{r}) \phi_j(\mathbf{r})^2 d\mathbf{r} := \langle \phi_j, \hat{A} \phi_j \rangle \quad \begin{matrix} \text{diagonal matrix element of } \hat{A}, \\ \text{operator multiplying by } A(\mathbf{r}) \end{matrix}$$

Quantum ergodicity

If the domain Ω is ergodic, does the intensity ϕ_j^2 become *equidistributed* (uniform across space) as $E_j \rightarrow \infty$?

More precisely, pick any test function $A = A(\mathbf{r})$, $\mathbf{r} \in \Omega$

- **classical** phase space average $\bar{A} = \frac{1}{\text{vol}(\Omega)} \int_{\Omega} A(\mathbf{r}) d\mathbf{r}$
- **quantum** intensity average

$$\int_{\Omega} A(\mathbf{r}) \phi_j(\mathbf{r})^2 d\mathbf{r} := \langle \phi_j, \hat{A} \phi_j \rangle \quad \begin{matrix} \text{diagonal matrix element of } \hat{A}, \\ \text{operator multiplying by } A(\mathbf{r}) \end{matrix}$$

Quantum Ergodicity Theorem (QET): “almost all levels equidistribute”
(Schnirelman ’74, Colin de Verdière ’85, Zelditch ’87, Z-Zworski ’96)

For ergodic domain Ω and all well-behaved A ,

$$\lim_{E_j \rightarrow \infty} \langle \phi_j, \hat{A} \phi_j \rangle - \bar{A} = 0$$

holds for a subset of levels with density $\rightarrow 1$ as $E_j \rightarrow \infty$

- exceptional subset *can persist*, but must have vanishing density

Questions to investigate

1. How dense is exceptional set?
2. What is *rate* of convergence of $\langle \phi_j, \hat{A}\phi_j \rangle$ to \bar{A} ?

Determines *practical* applicability of QET:

How high in E do you have to go to get the “Correspondence Principle”?

Examine numerically for...


Sinai-type domain (proven uniformly hyperbolic)

Piecewise constant A (numerically convenient)


Up to very high level numbers $j \sim 10^6$

Collect as many levels as you can!

(B '04)



Results: Diagonal matrix elements



25000 levels
of first 50000


mean
 $\langle \phi_j, \hat{A}\phi_j \rangle \rightarrow \bar{A}$

- No exceptional levels! (or their density $\lesssim 3 \times 10^{-5}$)
- Variance slowly decreasing, but how?

Results: Rate of variance decay with E

Local variance $V_A(E) := \frac{1}{m} \sum_{\substack{N \leq n < N+m \\ E_n \approx E}} \left| \langle \phi_j, \hat{A} \phi_j \rangle - \bar{A} \right|^2$

- Hard to measure *e.g.* 1% needs $m \sim 2 \times 10^4$ indep samples!



consistent with
power law model
 $V_A(E) = aE^{-\gamma}$
fit $\gamma = 0.48 \pm 0.01$

Discussion

- Previous studies: $j < 6000$, found various $\gamma \approx 0.3\text{--}0.5$ (Bäcker '98)
... we go 100 times higher in j : true asymptotic regime

Discussion

- Previous studies: $j < 6000$, found various $\gamma \approx 0.3\text{--}0.5$ (Bäcker '98)
... we go 100 times higher in j : true asymptotic regime
- $\gamma = 1/2$ consistent with RANDOM WAVES model:

$$\phi_j \text{ statistically similar to } \lim_{N \rightarrow \infty} N^{-1/2} \sum_{i=1}^N \sin(k \mathbf{n}_i \cdot \mathbf{r}) \quad (\text{Berry '77})$$

all directions \mathbf{n}_i , wavenumber $k = E^{1/2}$

However prefactor *not* consistent with this model: 80% error

Discussion

- Previous studies: $j < 6000$, found various $\gamma \approx 0.3\text{--}0.5$ (Bäcker '98)
... we go 100 times higher in j : true asymptotic regime
- $\gamma = 1/2$ consistent with RANDOM WAVES model:

$$\phi_j \text{ statistically similar to } \lim_{N \rightarrow \infty} N^{-1/2} \sum_{i=1}^N \sin(k \mathbf{n}_i \cdot \mathbf{r}) \quad (\text{Berry '77})$$

all directions \mathbf{n}_i , wavenumber $k = E^{1/2}$

However prefactor *not* consistent with this model: 80% error

- To understand prefactor: need semiclassical theory (Feingold-Peres '86)
variance of off-diag matrix els $\langle \phi_i, \hat{A}\phi_j \rangle \leftrightarrow$ classical auto-correlation func of A
But find convergence surprisingly slow: 7% error at $E \sim 10^6$


High-energy eigenfunction ϕ_n

$E \approx 10^6$

wavenumber
 $k \approx 10^3$

level number
 $j \approx 5 \times 10^4$


Compare to...



Random plane waves

interesting
stringy ridge
structures
appear due to
 $|\mathbf{k}| = \text{const.}$

... not ‘scars’
(O’Connor-
Heller ’86)



Quantum Unique Ergodicity & ‘scars’

QUE Conjecture (Rudnick-Sarnak '94): *there are no exceptional levels*
... made in context of negatively-curved manifolds

- Found strong evidence for QUE in ergodic domain
- Number theorists care about this (*e.g.* automorphic forms)


Quantum Unique Ergodicity & ‘scars’

QUE Conjecture (Rudnick-Sarnak ’94): *there are no exceptional levels*
... made in context of negatively-curved manifolds

- Found strong evidence for QUE in ergodic domain
- Number theorists care about this (*e.g.* automorphic forms)

SCARS: enhanced intensity ϕ_j^2 along *unstable* periodic orbits


- surprise numerical discovery by physicist (Heller ’84)



- QUE \Rightarrow effect of scars dies out
- variance decay $\sim E^{-1/2}$ consistent with expected narrowing of scars

Scarred eigenfunction at high E


Ω = stadium (ergodic)



~ 300 wavelengths across, $E \sim 10^6$


Numerical method for eigenproblem

$10^2\text{-}10^3$ wavelengths across system: Direct discretization
(finite elements, etc) *extremely inefficient*—need boundary method



Numerical method for eigenproblem

10²-10³ wavelengths across system: Direct discretization
(finite elements, etc) *extremely inefficient*—need boundary method




Method of Particular Solutions (MPS):

(Fox '67, etc)

1. Guess energy parameter E .
2. Choose basis set $\{\xi_i\}$ obeying Helmholtz eqn: $-\Delta \xi_i = E \xi_i$
e.g. Bessels $J_i(kr) \sin l\theta$, plane waves $\sin(k\mathbf{n}_i \cdot \mathbf{r})$
3. Trial function $u = \sum_{i=1}^N x_i \xi_i$ coeff vector $\mathbf{x} \in \mathbb{R}^N$
4. Find $t(E) = \min_{|\mathbf{x}|=1} \|u\|_{\partial\Omega}$ lowest ‘boundary condition error’
use M quadrature points \mathbf{r}_m , $t(E)$ is min sing val of $B_{mi} := \xi_i(\mathbf{r}_m)$
5. Repeat & root search: levels E_j are where $t(E) \rightarrow 0$ (or v. small)

Numerical method for eigenproblem

10²-10³ wavelengths across system: Direct discretization
(finite elements, etc) *extremely inefficient*—need boundary method



Method of Particular Solutions (MPS):

(Fox '67, etc)

1. Guess energy parameter E .
2. Choose basis set $\{\xi_i\}$ obeying Helmholtz eqn: $-\Delta \xi_i = E \xi_i$
e.g. Bessels $J_i(kr) \sin l\theta$, plane waves $\sin(k\mathbf{n}_i \cdot \mathbf{r})$
3. Trial function $u = \sum_{i=1}^N x_i \xi_i$ coeff vector $\mathbf{x} \in \mathbb{R}^N$
4. Find $t(E) = \min_{|\mathbf{x}|=1} \|u\|_{\partial\Omega}$ lowest ‘boundary condition error’
use M quadrature points \mathbf{r}_m , $t(E)$ is min sing val of $B_{mi} := \xi_i(\mathbf{r}_m)$
5. Repeat & root search: levels E_j are where $t(E) \rightarrow 0$ (or v. small)

MPS largely abandoned by numerical analysts in 70's ... Why?

- increase N : *many* vanishing sing vals at *all* $E!$ \rightarrow useless

Domain norm vs boundary norm


Easy to fix: use Rayleigh quotient $\lambda_1(E) = \min_{u \neq 0} \frac{\|u\|_{\partial\Omega}^2}{\|u\|_{\Omega}^2}$ (B '00)
min eigenvalue of (singular) generalized matrix eigenproblem

Domain norm vs boundary norm


Easy to fix: use Rayleigh quotient $\lambda_1(E) = \min_{u \neq 0} \frac{\|u\|_{\partial\Omega}^2}{\|u\|_{\Omega}^2}$ (B '00)

min eigenvalue of (singular) generalized matrix eigenproblem

Look at E -dependent spectrum $\{\lambda_l(E)^{1/2}\}$:



around level $j \sim 700$




- λ_1 seems quadratic in $(E - E_j)$, but coefficients (slopes) differ
- higher eigenvalue structure!

Special boundary weighting

What if replace $\|u\|_{\partial\Omega}^2$ by $\|u\|_{w,\partial\Omega}^2 := \int_{\partial\Omega} w(s) u(s)^2 ds$?

Choose weight $w(s) = [\mathbf{r}(s) \cdot \mathbf{n}(s)]^{-1}$... requires Ω star-shaped



- special: all **coeffs** (slopes) equal \rightarrow predictive power!
- much reduced *avoided crossings* between eigenvalues

WHY?

Analysis of spectrum $\{\lambda_l(E)\}$ – sketch

Basis set $\{\xi_i\}$ approx spans $\mathcal{H}_\Omega(E) := \{u : -\Delta u = Eu \text{ in } \Omega\}$

Analysis of spectrum $\{\lambda_l(E)\}$ – sketch


Basis set $\{\xi_i\}$ approx spans $\mathcal{H}_\Omega(E) := \{u : -\Delta u = Eu \text{ in } \Omega\}$

Via complete set $\{\partial_n \phi_j\}$ on boundary, show spectrum identical to that of $D\mathbf{v} = \lambda Q_w \mathbf{v}$ ∞ -dim matrices: $D(E) = \text{diag } (E - E_j)^2$, $Q_w = \text{indep of } E$

Analysis of spectrum $\{\lambda_l(E)\}$ – sketch

Basis set $\{\xi_i\}$ approx spans $\mathcal{H}_\Omega(E) := \{u : -\Delta u = Eu \text{ in } \Omega\}$

Via complete set $\{\partial_n \phi_j\}$ on boundary, show spectrum identical to that of $D\mathbf{v} = \lambda Q_w \mathbf{v}$ ∞ -dim matrices: $D(E) = \text{diag } (E - E_j)^2$, $Q_w = \text{indep of } E$



Perturbation theory of matrix problem yields:

(coeffs) $^{-1}$ = diag of Q_w , avoided crossing sizes = off-diag of Q_w p.19

The matrix Q_w : quasi-orthogonality

$$(Q_w)_{ij} := \int_{\partial\Omega} w^{-1} \partial_n \phi_i \partial_n \phi_j \, ds$$

Eigenfunction normal derivs $\partial_n \phi_j$
not orthogonal on $\partial\Omega$, but...

For special choice $w = (\mathbf{r} \cdot \mathbf{n})^{-1}$:

$$(Q_w)_{ij} = 2\delta_{ij}E_j + c_{ij}$$

- Conjecture (Vergini '94): off-diag error term growth $|c_{ij}| \sim |E_i - E_j|$
- Semiclassics (B-Cohen-Heller '00): for Ω ergodic, $|c_{ij}| \sim (E_i - E_j)^2$

The matrix Q_w : quasi-orthogonality


$$(Q_w)_{ij} := \int_{\partial\Omega} w^{-1} \partial_n \phi_i \partial_n \phi_j \, ds$$

Eigenfunction normal derivs $\partial_n \phi_j$
not orthogonal on $\partial\Omega$, but...

For special choice $w = (\mathbf{r} \cdot \mathbf{n})^{-1}$:

$$(Q_w)_{ij} = 2\delta_{ij}E_j + c_{ij}$$

- Conjecture (Vergini '94): off-diag error term growth $|c_{ij}| \sim |E_i - E_j|$
- Semiclassics (B-Cohen-Heller '00): for Ω ergodic, $|c_{ij}| \sim (E_i - E_j)^2$
- Theorem (B '04): for all Ω , ergodic or not, $|c_{ij}| \leq C_\Omega(E_i - E_j)^2$




quasi-orthogonality:
locally, $Q_w \rightarrow$ Identity
as $E_i, E_j \rightarrow \infty$

explains coeffs, and
small avoided crossings

Result 1: Improved inclusion bounds

In practise $t(u) = \|u\|_{w,\partial\Omega}/\|u\|_\Omega$ never reaches zero.

But we can bound the distance to nearest level $\frac{|E - E_j|}{E_j^{1/2}} \leq C_\Omega t(u)$




- $O(E^{1/2})$ tighter than classical bounds (Kuttler-Sigillito '84)

Result 2: scaling method (Vergini '94, explained by B '00, '04)

Replace norm $\|u\|_{\Omega}^2$ in Rayleigh quotient by $\int_{\partial\Omega} (\mathbf{r} \cdot \mathbf{n})^{-1} u \mathbf{r} \cdot \nabla u \, ds$

Then E -dependent spectrum allows accurate prediction at high E :



- Many levels (E_j and ϕ_j) result from single matrix eigenproblem
- Speed gain (vs all known methods, incl integral eqns): $O(E^{1/2})$, can be 10^3
- Whole calculation $\sim 10^4$ levels at $j \sim 10^5$ takes only few CPU days

Basis set for $\mathcal{H}_\Omega(E)$ (B '02)

Plane waves

$$\xi_l(\mathbf{r}) = e^{i\mathbf{k}_l \cdot \mathbf{r}}$$

- inaccurate *e.g.* $\Omega = \text{stadium}$
- fail as $E \rightarrow \infty$ if Ω nonconvex

Basis set for $\mathcal{H}_\Omega(E)$ (B '02)

Plane waves

$$\xi_l(\mathbf{r}) = e^{i\mathbf{k}_l \cdot \mathbf{r}}$$

- inaccurate *e.g.* $\Omega = \text{stadium}$
- fail as $E \rightarrow \infty$ if Ω nonconvex


Neumann functions

$$\xi_l(\mathbf{r}) = Y_0(k|\mathbf{r} - \mathbf{x}_l|)$$

good for nonconvex

Why?

- $\partial\Omega^+$ layer potential complete in $\mathcal{H}_{\Omega^+}(E)$, for $k \neq$ Dir. eigenvalue of Ω^+
- $\mathcal{H}_{\Omega^+}(E)$ dense $\subset \mathcal{H}_\Omega(E)$ (Lax '56, Runge approx)



Basis set for $\mathcal{H}_\Omega(E)$ (B '02)

Plane waves

$$\xi_l(\mathbf{r}) = e^{i\mathbf{k}_l \cdot \mathbf{r}}$$

- inaccurate *e.g.* $\Omega = \text{stadium}$
- fail as $E \rightarrow \infty$ if Ω nonconvex

Neumann functions

$$\xi_l(\mathbf{r}) = Y_0(k|\mathbf{r} - \mathbf{x}_l|)$$

good for nonconvex

Why?



- $\partial\Omega^+$ layer potential complete in $\mathcal{H}_{\Omega^+}(E)$, for $k \neq$ Dir. eigenvalue of Ω^+
- $\mathcal{H}_{\Omega^+}(E)$ dense $\subset \mathcal{H}_\Omega(E)$ (Lax '56, Runge approx)

If no analytic continuation of ϕ_j outside Ω ? (Eckmann-Pillet '95)

e.g. corner, $\theta \neq \frac{\pi}{n}$

Empirical: errors small!

[MOVIE](#)



Conclusions

Dirichlet eigenproblem –

- Applications: technology, to physics, to number theory...

At high energy, wealth of interesting phenomena –

- Equidistribution of modes: unprecedented numerical tests
- Accelerated ($\times 10^3$) computational tools: analyzed and developed

Conclusions

Dirichlet eigenproblem –

- Applications: technology, to physics, to number theory...

At high energy, wealth of interesting phenomena –

- Equidistribution of modes: unprecedented numerical tests
- Accelerated ($\times 10^3$) computational tools: analyzed and developed

Many future directions:


- Break the star-shaped restriction?
- Can similarly accelerate integral equation eigenmethods?
- Resonance, open (scattering) problems? *e.g.* laser cavity

Preprints/talks: <http://www.cims.nyu.edu/~barnett>

Missing levels?

Weyl's estimate for $N(E)$, the # eigenvalues $E_j < E$:

$$N_{\text{Weyl}}(E) = \frac{\text{vol}(\Omega)}{4\pi} E - \frac{L}{4\pi} \sqrt{E} + O(1) \dots$$



- not one level missing in sequence of 6812