

Numerical study of the asymptotic rate of quantum ergodicity in chaotic billiards

MIT/Harvard Analysis Seminar, 17 March, 2006

Alex Barnett

Mathematics Department, Dartmouth College

Thanks to P. Sarnak (NYU), S. Zelditch (JHU),
N. Trefethen (Oxford), T. Betcke (Braunschweig)

Dirichlet eigenproblem

Normal modes of elastic membrane or ‘drum’ (Helmholtz, Germain, 19thC)


Eigenfunctions ϕ_j of Laplacian $\Delta := \partial_{x_1}^2 + \partial_{x_2}^2$ in bounded domain $\Omega \subset \mathbb{R}^2$

$$-\Delta\phi_j = E_j\phi_j \quad \phi_j|_{\partial\Omega} = 0 \text{ Dirichlet BC} \quad \langle\phi_i, \phi_j\rangle = \delta_{ij}$$

Dirichlet eigenproblem

Normal modes of elastic membrane or ‘drum’ (Helmholtz, Germain, 19thC)
Eigenfunctions ϕ_j of Laplacian $\Delta := \partial_{x_1}^2 + \partial_{x_2}^2$ in bounded domain $\Omega \subset \mathbb{R}^2$

$$-\Delta\phi_j = E_j\phi_j \quad \phi_j|_{\partial\Omega} = 0 \quad \text{Dirichlet BC} \quad \langle\phi_i, \phi_j\rangle = \delta_{ij}$$



‘energy’ eigenvalues
 $E_1 < E_2 \leq E_3 \leq \dots$


wavenumber $k_j = E_j^{1/2}$

wavelength $= 2\pi/k_j$

Dirichlet eigenproblem

Normal modes of elastic membrane or ‘drum’ (Helmholtz, Germain, 19thC)
Eigenfunctions ϕ_j of Laplacian $\Delta := \partial_{x_1}^2 + \partial_{x_2}^2$ in bounded domain $\Omega \subset \mathbb{R}^2$

$$-\Delta\phi_j = E_j\phi_j \quad \phi_j|_{\partial\Omega} = 0 \quad \text{Dirichlet BC} \quad \langle\phi_i, \phi_j\rangle = \delta_{ij}$$



‘energy’ eigenvalues
 $E_1 < E_2 \leq E_3 \leq \dots$

wavenumber $k_j = E_j^{1/2}$


wavelength $= 2\pi/k_j$

- Time-harmonic solns of wave eqn (acoustics, optics, quantum, etc)
- Asymptotics of ϕ_j as eigenvalue $E_j \rightarrow \infty$? Depends on shape...

Outline

- Eigenproblem (quantum)
- Billiard system (classical)
- ‘Quantum chaos’ conjectures and numerical results on
 - exceptional ‘scarred’ eigenmodes
 - rate at which eigenmodes become spatially uniform
- Numerical methods sketch if time

Billiards (the classical analogue)




state $z := (x_1, x_2, \xi_1, \xi_2)$
 $\in Z := \Omega \times \mathbb{R}^2$, phase space

free motion (geodesic flow)

trajectory $z_0(t) = G_t z_0$

Hamiltonian dynamical system: $H(x, \xi) = |\xi|^2$ conserved

Billiards (the classical analogue)




state $z := (x_1, x_2, \xi_1, \xi_2)$
 $\in Z := \Omega \times \mathbb{R}^2$, phase space

free motion (geodesic flow)

trajectory $z_0(t) = G_t z_0$


Hamiltonian dynamical system: $H(x, \xi) = |\xi|^2$ conserved

integrable:



d conserved quantities ($d=2$)

ergodic:



only H conserved

Properties of ergodic billiards

Ergodic: $\forall A \in L^1(Z)$, for almost every $z_0 \in Z$,

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T A(z_0(t)) dt = \frac{1}{\text{vol}(Z)} \int_Z A(z) dz =: \bar{A} \quad \text{spatial average}$$


Properties of ergodic billiards

Ergodic: $\forall A \in L^1(Z)$, for almost every $z_0 \in Z$,

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T A(z_0(t)) dt = \frac{1}{\text{vol}(Z)} \int_Z A(z) dz =: \bar{A} \quad \text{spatial average}$$

- ‘hyperbolic’: $|z_1(t) - z_2(t)| \sim ce^{\Lambda t}$ $0 < \Lambda = \text{Lyapunov exponent}$
- ‘Anosov’: uniformly hyperbolic (all periodic orbits unstable, isolated)

e.g. Bunimovich stadium ergodic but not Anosov...




Properties of ergodic billiards

Ergodic: $\forall A \in L^1(Z)$, for almost every $z_0 \in Z$,

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T A(z_0(t)) dt = \frac{1}{\text{vol}(Z)} \int_Z A(z) dz =: \bar{A} \quad \text{spatial average}$$

- ‘hyperbolic’: $|z_1(t) - z_2(t)| \sim ce^{\Lambda t}$ $0 < \Lambda = \text{Lyapunov exponent}$
- ‘Anosov’: uniformly hyperbolic (all periodic orbits unstable, isolated)

e.g. Bunimovich stadium ergodic but not Anosov...



- autocorrelation $C_A(\tau) := \frac{1}{\text{vol}(Z)} \int_Z A(z_0) A(z_0(\tau)) dz_0 - \bar{A}^2$
- flow is ‘mixing’ if $\forall A \in L^2(Z)$, $\lim_{\tau \rightarrow \infty} C_A(\tau) = 0$


Properties of ergodic billiards

Ergodic: $\forall A \in L^1(Z)$, for almost every $z_0 \in Z$,

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^T A(z_0(t)) dt = \frac{1}{\text{vol}(Z)} \int_Z A(z) dz =: \bar{A} \quad \text{spatial average}$$

- ‘hyperbolic’: $|z_1(t) - z_2(t)| \sim ce^{\Lambda t}$ $0 < \Lambda = \text{Lyapunov exponent}$
- ‘Anosov’: uniformly hyperbolic (all periodic orbits unstable, isolated)

e.g. Bunimovich stadium ergodic but not Anosov...



- autocorrelation $C_A(\tau) := \frac{1}{\text{vol}(Z)} \int_Z A(z_0) A(z_0(\tau)) dz_0 - \bar{A}^2$
- flow is ‘mixing’ if $\forall A \in L^2(Z)$, $\lim_{\tau \rightarrow \infty} C_A(\tau) = 0$

QUANTUM CHAOS: what are eigenmodes when flow ergodic? (Einstein 1917)

Modes ϕ_j are irregular: VIEW $j \sim 3000$: 45 wavelengths across

Quantum Ergodicity Theorem (QET)

For Ω with ergodic flow, and ‘well-behaved’ $A : \Omega \rightarrow \mathbb{R}$,

$$\lim_{E_j \rightarrow \infty} \langle \phi_j, A\phi_j \rangle - \overline{A} = 0 \quad \forall j \text{ except subseq. of vanishing density}$$

(Schnirelman '74, Colin de Verdière '85, Zelditch '87, Z-Zworski '96)


Quantum Ergodicity Theorem (QET)

For Ω with ergodic flow, and ‘well-behaved’ $A : \Omega \rightarrow \mathbb{R}$,

$$\lim_{E_j \rightarrow \infty} \langle \phi_j, A\phi_j \rangle - \overline{A} = 0 \quad \forall j \text{ except subseq. of vanishing density}$$

(Schnirelman '74, Colin de Verdière '85, Zelditch '87, Z-Zworski '96)

- measure $|\phi_j|^2 dx \xrightarrow{\text{weak in } L^1} \frac{1}{\text{vol}(\Omega)} dx$



- generally $A : Z \rightarrow \mathbb{R}$, ‘well-behaved’ operator $\text{Op}(A)$, 0^{th} -order Ψ DO
- proof: Egorov’s Thm (propagation of singularities of wave eqn)


Quantum Ergodicity Theorem (QET)

For Ω with ergodic flow, and ‘well-behaved’ $A : \Omega \rightarrow \mathbb{R}$,

$$\lim_{E_j \rightarrow \infty} \langle \phi_j, A\phi_j \rangle - \bar{A} = 0 \quad \forall j \text{ except subseq. of vanishing density}$$

(Schnirelman ’74, Colin de Verdière ’85, Zelditch ’87, Z-Zworski ’96)

- measure $|\phi_j|^2 dx \xrightarrow{\text{weak in } L^1} \frac{1}{\text{vol}(\Omega)} dx$



- generally $A : Z \rightarrow \mathbb{R}$, ‘well-behaved’ operator $\text{Op}(A)$, 0^{th} -order Ψ DO
- proof: Egorov’s Thm (propagation of singularities of wave eqn)

At what rate is limit reached?

At what rate does the density of excluded subsequence vanish?

Quantum Unique Ergodicity (QUE)


Conj. (Rudnick-Sarnak '94): There is no excluded subsequence in QET

- ‘Unique’ refers to $\frac{1}{\text{vol}(\Omega)}dx$ being unique limit measure of $|\phi_j|^2 dx$

Quantum Unique Ergodicity (QUE)

Conj. (Rudnick-Sarnak '94): There is no excluded subsequence in QET

- ‘Unique’ refers to $\frac{1}{\text{vol}(\Omega)}dx$ being unique limit measure of $|\phi_j|^2 dx$
- context: negatively curved manifolds (Anosov)




Recent analytic results:

- Proven to hold for *arithmetic* manifold $SL_2(\mathbb{Z}) \backslash \mathbb{H}$ (Lindenstrauss '03)
... very special system, symmetries, all $\Lambda = 1$
- Proven *not* to hold for some quantum Arnold cat maps (Faure et al '03)

Quantum Unique Ergodicity (QUE)

Conj. (Rudnick-Sarnak '94): There is no excluded subsequence in QET

- ‘Unique’ refers to $\frac{1}{\text{vol}(\Omega)}dx$ being unique limit measure of $|\phi_j|^2 dx$
- context: negatively curved manifolds (Anosov)



Recent analytic results:


- Proven to hold for *arithmetic* manifold $SL_2(\mathbb{Z}) \backslash \mathbb{H}$ (Lindenstrauss '03)
... very special system, symmetries, all $\Lambda = 1$
- Proven *not* to hold for some quantum Arnold cat maps (Faure et al '03)

QUE might be a surprise for Euclidean cavity eigenmodes because...

Scars: an exceptional subsequence?


high- j modes $|\phi_j|^2$ often localized on unstable periodic orbits (UPO)
e.g. numerical study of stadium eigenmodes

(Heller '84)



- Wavepacket heuristics \Rightarrow transverse width dies as $\sim E^{-1/4}$
- Note also exceptional ‘bouncing ball’ sequence (since not Anosov)

High-eigenvalue stadium scar



- ‘strong’ scar: \exists sequence $\{\phi_{j_n}\}$ with $O(1)$ mass on UPO
- QUE implies ‘weak’ scarring: UPO mass is $o(1)$

Which occurs? No analytic results in Euclidean billiards...


Numerical experiments

(B '04, to appear CPAM '06)

dispersing billiard, proven Anosov (Sinai '70)

desymmetrized, generic Λ exponents

$$\text{test function } A = \begin{cases} 1 & \text{in } \Omega_A, \\ 0 & \text{otherwise} \end{cases}$$




Numerical experiments

(B '04, to appear CPAM '06)

dispersing billiard, proven Anosov (Sinai '70)

desymmetrized, generic Λ exponents

$$\text{test function } A = \begin{cases} 1 & \text{in } \Omega_A, \\ 0 & \text{otherwise} \end{cases}$$



Large-scale study, 30,000 modes in range $j \sim 10^4$ to 10^6 , enabled by:

1. Efficient boundary-based numerics for ϕ_j ('scaling method')
 2. matrix elements $\langle \phi_j, A\phi_j \rangle = \int_{\Omega_A} \phi_j^2 dx$, using integrals on $\partial\Omega_A$
- 100 times higher in j than any previous studies (e.g. Bäcker '98)
 - only a few CPU-days total


Typical high-frequency mode

225 wavelengths
across system

level number
 $j \approx 5 \times 10^4$


$E_j \approx 10^6$

Note: making this
plot is much more
expensive than
finding $\int_{\Omega_A} \phi_j^2 dx$



Raw diagonal matrix element data

To reach high E , compute ϕ_j only in intervals $E_j \in [E, E + L(E)]$



- strong evidence for QUE (exceptional density $< 3 \times 10^{-5}$)

Rate of quantum ergodicity

energy interval $I_E = [E, E + L(E)]$, width scales as $L(E) = O(E^{1/2})$
mode count in interval $N(I_E) := \#\{j : E_j \in I_E\}$

‘quantum variance’ $V_A(E) := \frac{1}{N(I_E)} \sum_{E_j \in I_E} |\langle \phi_j, A\phi_j \rangle - \overline{A}|^2$

Rate of quantum ergodicity

energy interval $I_E = [E, E + L(E)]$, width scales as $L(E) = O(E^{1/2})$
mode count in interval $N(I_E) := \#\{j : E_j \in I_E\}$

$$\text{'quantum variance'} V_A(E) := \frac{1}{N(I_E)} \sum_{E_j \in I_E} |\langle \phi_j, A\phi_j \rangle - \bar{A}|^2$$

Conj. (Feingold-Peres '86): $V_A(E) \sim \frac{g\tilde{C}_A(0)}{\text{vol}(\Omega)} E^{-1/2}$

where $g = 2$, as in random matrix theory (RMT), GOE symmetry

$$\text{and } \tilde{C}_A(\omega) := \int_{-\infty}^{\infty} C_A(\tau) e^{i\omega\tau} d\tau$$

Heuristic argument relies on two steps:


- QET for operator $A(0)A(\tau)$: off-diagonal variance given by $\tilde{C}_A(\omega)$
- assume statistical independence of ϕ_j within I_E (as in RMT)

Classical variance

$$\tilde{C}_A(\omega) := \int_{-\infty}^{\infty} C_A(\tau) e^{i\omega\tau} d\tau$$

FT of autocorrelation =
power spectral density
of signal $A(z_0(t))$
along ergodic trajectory

LISTEN to it!




Classical variance

$$\tilde{C}_A(\omega) := \int_{-\infty}^{\infty} C_A(\tau) e^{i\omega\tau} d\tau$$

FT of autocorrelation =
power spectral density
of signal $A(z_0(t))$
along ergodic trajectory

LISTEN to it!




Physically: $\tilde{C}_A(\omega)$ is heating rate due to periodic driving of classical system, at frequency ω , by external field A (Cohen '99, B '00)

FP Conj.: $\omega \rightarrow 0$ limit gives prefactor for quantum ergodicity rate

- Note Ω symmetric, A antisymm: conj. fails, $V_A(E) = 0$ (Sarnak '03)
Does it hold in our nonsymmetric case?

Results on ergodicity rate



consistent with
power law model

$$V_A(E) = aE^{-\gamma}$$

$$\text{fit } \gamma = 0.48 \pm 0.01$$

$\gamma = 1/2$ agrees with

- FP conjecture
- random-wave model
- scar theory

- large numbers of modes \rightarrow unprecedented accuracy ($< 1\%$)
- consistent with FP prefactor, convergence slow: 7% off at $j = 10^5$
- asymptotic regime seen for first time (but more data needed!)

Boundary-Interior Method of Particular Solutions (B '00)

Task: find ϕ_j and E_j such that $(\Delta + E_j)\phi_j = 0$ and $\phi_j|_{\partial\Omega} = 0$

Boundary-Interior Method of Particular Solutions (B '00)

Task: find ϕ_j and E_j such that $(\Delta + E_j)\phi_j = 0$ and $\phi_j|_{\partial\Omega} = 0$

Given trial energy parameter $E\dots$

- $u = \sum_{i=1}^N x_i \xi_i$, where each basis function obeys $(\Delta + E)\xi_i = 0$ in Ω

Boundary-Interior Method of Particular Solutions (B '00)

Task: find ϕ_j and E_j such that $(\Delta + E_j)\phi_j = 0$ and $\phi_j|_{\partial\Omega} = 0$

Given trial energy parameter E ...

- $u = \sum_{i=1}^N x_i \xi_i$, where each basis function obeys $(\Delta + E)\xi_i = 0$ in Ω
- if can find coeff vector $\mathbf{x} \in \mathbb{R}^N$ giving $u|_{\partial\Omega} = 0$, but $u \neq 0$ in Ω
... then u is a mode ϕ_j and E is its eigenvalue E_j



Boundary-Interior Method of Particular Solutions (B '00)

Task: find ϕ_j and E_j such that $(\Delta + E_j)\phi_j = 0$ and $\phi_j|_{\partial\Omega} = 0$

Given trial energy parameter E ...

- $u = \sum_{i=1}^N x_i \xi_i$, where each basis function obeys $(\Delta + E)\xi_i = 0$ in Ω
- if can find coeff vector $\mathbf{x} \in \mathbb{R}^N$ giving $u|_{\partial\Omega} = 0$, but $u \neq 0$ in Ω
... then u is a mode ϕ_j and E is its eigenvalue E_j

'boundary error' $t(E) := \min_{u \in \text{Span}\{\xi_i\}, u \neq 0} \frac{\|u\|_{L^2(\partial\Omega)}}{\|u\|_{L^2(\Omega)}}$, (Rayleigh quotient)





Boundary-Interior Method of Particular Solutions (B '00)

Task: find ϕ_j and E_j such that $(\Delta + E_j)\phi_j = 0$ and $\phi_j|_{\partial\Omega} = 0$

Given trial energy parameter E ...

- $u = \sum_{i=1}^N x_i \xi_i$, where each basis function obeys $(\Delta + E)\xi_i = 0$ in Ω
- if can find coeff vector $\mathbf{x} \in \mathbb{R}^N$ giving $u|_{\partial\Omega} = 0$, but $u \neq 0$ in Ω
... then u is a mode ϕ_j and E is its eigenvalue E_j

$$\text{'boundary error'} t(E) := \min_{u \in \text{Span}\{\xi_i\}, u \neq 0} \frac{\|u\|_{L^2(\partial\Omega)}}{\|u\|_{L^2(\Omega)}}, \quad (\text{Rayleigh quotient})$$



- $t(E)$ is lowest generalized eigenvalue of pair of E -dep. matrices
- finally search in E for minima of $t(E)$

Scaling method


(Vergini '94, B '00, '04)

new Rayleigh quotient $\frac{f(u)}{g(u)}$ where

$$f(u, v) := \int_{\partial\Omega} (\mathbf{x} \cdot \mathbf{n})^{-1} uv,$$

$$g(u, v) := \int_{\partial\Omega} (\mathbf{x} \cdot \mathbf{n})^{-1} (u \mathbf{x} \cdot \nabla v + v \mathbf{x} \cdot \nabla u)$$


Generalized eigenvalues now approximately linear in $E - E_j$:



- solving $F\mathbf{x} = \hat{\lambda}G\mathbf{x}$ at **single** E value gives *all* nearest $O(N)$ modes
- no root search, no missing levels, efficiency gain $O(E^{1/2})$
- eigenvectors \mathbf{x} give *dilated* (scaled) approximations to modes ϕ_j



Mushroom billiard (mixed dynamics)

Inspired by AMS Notices March '06, Betcke & I computed...



- Due to re-entrant corner, Fourier-Bessel basis choice non-trivial
- We believe first high eigenmodes found for mushroom (last week)

High energy mushroom billiard



Conclusions

Studied nonsymmetric Anosov billiard eigenmodes. . .

- strong evidence for QUE
- slow asymptotic convergence to Feingold-Peres conjecture

Scaling method and boundary integral identities

- 10^3 faster than any other known method for eigenmodes
- cheaply get large number of high-eigenvalue matrix elements

Ongoing:

- bouncing-ball mode statistics (with A. Hassell)
- basis functions, scaling method numerical analysis (with T. Betcke)


Partial funding by: NSF (DMS-0507614)

Preprints/talks: <http://math.dartmouth.edu/~ahb>

Missing levels?

Weyl's estimate for $N(E)$, the # eigenvalues $E_j < E$:

$$N_{\text{Weyl}}(E) = \frac{\text{vol}(\Omega)}{4\pi} E - \frac{L}{4\pi} \sqrt{E} + O(1) \dots$$



- not one level missing in sequence of 6812