

Accurate and Efficient Numerical Methods for High-frequency 3D Wave Scattering and Periodic Geometries

CIMS Num. Analysis & Sci. Comput. Seminar 2/15/13


Alex Barnett (Dartmouth College)

Collaborators: Leslie Greengard (NYU), Andreas Klöckner (NYU), Mike O'Neil (NYU), Zydrunas Gimbutas (NYU), Adrianna Gillman (Dartmouth).




Barnett group research

- numerical methods for scattering of waves
 - time-harmonic (frequency domain)
 - high-order accurate, “fast” algorithms
 - periodic structures, corners, variable media



- eigenvalue problems and quantum chaos
 - fast algorithms for high freq.
 - photonic crystal band structure
 - PDE / numerical analysis



- other applied mathematics: ecology, medical imaging, music

Quantum chaos

Closed cavity (drum) modes

Shown: high mode numbers,
 $j = 1, 10, 10^2, 10^3, 10^4, 10^5$

Statistics as $j \rightarrow \infty$?

- methods $10^3 \times$ faster than any other known (B '06, B–Hassell '12)
- rigorous error analysis (B '09, B–Hassell '11)

Notices

of the American Mathematical Society


January 2008

Volume 55, Number 1

An Evaluation
 of Mathematics
 Competitions Using
 Item Response Theory
 page 8

Your Hit Parade:
 The Top Ten Most
 Fascinating Formulas
 in Ramanujan's Lost
 Notebook
 page 18

New York Meeting
 page 98




Quantum chaos (see page 41)

Nodal region counts of modes (w/ K. Konrad, snr thesis '12)

$\nu_j :=$ # of conn. regions of definite sign in mode j



Thm: $\nu_j \leq j$. (Courant '53) $\lim_{j \rightarrow \infty} \frac{\nu_j}{j} = r = ?$



Nodal region counts of modes (w/ K. Konrad, snr thesis '12)

$\nu_j :=$ # of conn. regions of definite sign in mode j

Thm: $\nu_j \leq j$. (Courant '53) $\lim_{j \rightarrow \infty} \frac{\nu_j}{j} = r = ?$



10^9 regions counted. $r \approx 0.0596$ shows conj. 5% off (Bogomolny–Schmit '02)

Today:

- I. New tool for solving high-frequency scattering in 2D and 3D
accurate near-field evaluation → efficient quadratures
- II. Robust schemes for solving waves in periodic structures
low rank property helps solve gratings and crystals

Scattering: motivation & applications


electromagnetics, radar cross section

acoustics, quantum, nanophotonics

devices, lithography, imaging

inverse probs: need good fwd solvers

many involve piecewise-uniform media



(Dolean '10)

Scattering: motivation & applications


electromagnetics, radar cross section

acoustics, quantum, nanophotonics

devices, lithography, imaging

inverse probs: need good fwd solvers

many involve piecewise-uniform media



(Dolean '10)

Need solvers:



- robust
don't break at certain parameters or geometries
- high-order convergent
reliability, sensitivity anal, optimization
- efficient
optimal complexity: ideally linear in # unknowns

Applications of periodic scattering

Gratings, filters, antennae, photonic crystals, meta-materials, solar...

Applications of periodic scattering



Gratings, filters, antennae, photonic crystals, meta-materials, solar...




multi-layer dielectric diffraction
grating, NIF lasers (LLNL)
 2×10^6 periods! (Barty '04)

Applications of periodic scattering


Gratings, filters, antennae, photonic crystals, meta-materials, solar...



multi-layer dielectric diffraction
grating, NIF lasers (LLNL)
 2×10^6 periods! (Barty '04)



plasmonic solar cell (Atwater '10)





Si microwires absorber (Kelzenberg '10)

↑ high
aspect
ratio


- Design optimization
- Simulation at $>10^3$ inc. angles, frequencies

Applications of periodic scattering


Gratings, filters, antennae, photonic crystals, meta-materials, solar...



multi-layer dielectric diffraction
grating, NIF lasers (LLNL)
 2×10^6 periods! (Barty '04)



plasmonic solar cell (Atwater '10)



Si microwires absorber (Kelzenberg '10)

↑ high
aspect
ratio


- Design optimization
- Simulation at $>10^3$ inc. angles, frequencies

But first some old and new tools for the non-periodic case...

I. Scattering of time-harmonic waves

freq. ω , wavelength $\lambda = \frac{2\pi}{\omega}$

obstacle $\Omega \subset \mathbb{R}^d$, $d = 2, 3$, bdry $\partial\Omega$



plane wave $u^{\text{inc}}(\mathbf{x}) = e^{i\omega\hat{n}\cdot\mathbf{x}}$

physical field $u^{\text{tot}} = u^{\text{inc}} + u$

scattered field u obeys BVP:

$$(\Delta + \omega^2)u = 0 \quad \text{in } \mathbb{R}^d \setminus \overline{\Omega}$$

$$u = -u^{\text{inc}} \quad \text{on } \partial\Omega$$

Sommerfeld radiation condition:


$$\frac{\partial u}{\partial r} - i\omega u = o(r^{-(d-1)/2}), \quad r \rightarrow \infty$$

\exists unique soln. Sound-soft acoustics / Maxwell TM ($u=E_z, \partial\Omega=\text{PEC}$)

I. Scattering of time-harmonic waves

freq. ω , wavelength $\lambda = \frac{2\pi}{\omega}$

obstacle $\Omega \subset \mathbb{R}^d$, $d = 2, 3$, bdry $\partial\Omega$



plane wave $u^{\text{inc}}(\mathbf{x}) = e^{i\omega\hat{n}\cdot\mathbf{x}}$

physical field $u^{\text{tot}} = u^{\text{inc}} + u$

scattered field u obeys BVP:

$$(\Delta + \omega^2)u = 0 \quad \text{in } \mathbb{R}^d \setminus \overline{\Omega}$$

$$u = -u^{\text{inc}} \quad \text{on } \partial\Omega$$

Sommerfeld radiation condition:

$$\frac{\partial u}{\partial r} - i\omega u = o(r^{-(d-1)/2}), \quad r \rightarrow \infty$$

\exists unique soln. Sound-soft acoustics / Maxwell TM ($u = E_z$, $\partial\Omega = \text{PEC}$)

Errors 10^{-10} (as here) demands high-order ... how do this cheaply?

Discretizing: volume vs boundary?

Conventional numerical PDE solution: mesh volume, N unknowns
e.g. finite differencing, finite elements

Discretizing: volume vs boundary?

Conventional numerical PDE solution: mesh volume, N unknowns

e.g. finite differencing, finite elements

Problems:

- (1) needs many unknowns per wavelength: $N > O(\omega^d)$
“pollution” (Babuska–Sauter)
- (2) meshing expensive ($d=3$)
- (3) must truncate to finite domain w/ approximate rad. cond.
- (4) matrix sparse, but ill-conditioned (even though BVP isn’t!)

Better to discretize boundaries: $N \sim O(\omega^{d-1})$ much smaller

Discretizing: volume vs boundary?

Conventional numerical PDE solution: mesh volume, N unknowns

e.g. finite differencing, finite elements

Problems:

(1) needs many unknowns per wavelength: $N > O(\omega^d)$

“pollution” (Babuska–Sauter)

(2) meshing expensive ($d=3$)

(3) must truncate to finite domain w/ approximate rad. cond.

(4) matrix sparse, but ill-conditioned (even though BVP isn’t!)

Better to discretize boundaries: $N \sim O(\omega^{d-1})$ much smaller

Fredholm integral equation reminder: given kernel $k(t, s)$ cont. for $s \neq t$

solve for func. τ : $c\tau(t) + \int_0^{2\pi} k(t, s)\tau(s)ds = f(t), \quad \forall t \in (0, 2\pi)$

- if $c = 0$: 1st-kind, ill-posed. Discretize \rightarrow ill-cond. as N grows
- if $c \neq 0$: 2nd-kind, always \exists unique solution; well-cond. indep of N

Representing PDE solns: potential theory

‘charge’ (source of waves) distributed along curve Γ w/ density func.


Single-, double-layer potentials, $\mathbf{x} \in \mathbb{R}^2$

$$v(\mathbf{x}) = \int_{\Gamma} G(\mathbf{x}, \mathbf{y}) \sigma(\mathbf{y}) ds_{\mathbf{y}} := (\mathcal{S}\sigma)(\mathbf{x})$$

$$u(\mathbf{x}) = \int_{\Gamma} \frac{\partial G}{\partial n_{\mathbf{y}}}(\mathbf{x}, \mathbf{y}) \tau(\mathbf{y}) ds_{\mathbf{y}} := (\mathcal{D}\tau)(\mathbf{x})$$

$$G(\mathbf{x}, \mathbf{y}) := G(\mathbf{x} - \mathbf{y}) = \frac{i}{4} H_0^{(1)}(\omega |\mathbf{x} - \mathbf{y}|)$$

kernel is Helmholtz fundamental soln
a.k.a. free space Greens func



Representing PDE solns: potential theory

‘charge’ (source of waves) distributed along curve Γ w/ density func.


Single-, double-layer potentials, $\mathbf{x} \in \mathbb{R}^2$

$$v(\mathbf{x}) = \int_{\Gamma} G(\mathbf{x}, \mathbf{y}) \sigma(\mathbf{y}) ds_{\mathbf{y}} := (\mathcal{S}\sigma)(\mathbf{x})$$

$$u(\mathbf{x}) = \int_{\Gamma} \frac{\partial G}{\partial n_{\mathbf{y}}}(\mathbf{x}, \mathbf{y}) \tau(\mathbf{y}) ds_{\mathbf{y}} := (\mathcal{D}\tau)(\mathbf{x})$$

$$G(\mathbf{x}, \mathbf{y}) := G(\mathbf{x} - \mathbf{y}) = \frac{i}{4} H_0^{(1)}(\omega |\mathbf{x} - \mathbf{y}|)$$

kernel is Helmholtz fundamental soln
a.k.a. free space Greens func



Jump relations: field limit as $\mathbf{x} \rightarrow \Gamma$ can depend on which side (\pm):

$$v^\pm = S\sigma \quad \text{no jump}$$

$$u^\pm = D\tau \pm \frac{1}{2}\tau \quad \text{jump}$$

S, D bdry integral ops w/ above kernels,
both bounded in $C(\Gamma) \rightarrow C^1(\Gamma)$

Representing PDE solns: potential theory

‘charge’ (source of waves) distributed along curve Γ w/ density func.


Single-, double-layer potentials, $\mathbf{x} \in \mathbb{R}^2$

$$v(\mathbf{x}) = \int_{\Gamma} G(\mathbf{x}, \mathbf{y}) \sigma(\mathbf{y}) ds_{\mathbf{y}} := (\mathcal{S}\sigma)(\mathbf{x})$$

$$u(\mathbf{x}) = \int_{\Gamma} \frac{\partial G}{\partial n_{\mathbf{y}}}(\mathbf{x}, \mathbf{y}) \tau(\mathbf{y}) ds_{\mathbf{y}} := (\mathcal{D}\tau)(\mathbf{x})$$

$$G(\mathbf{x}, \mathbf{y}) := G(\mathbf{x} - \mathbf{y}) = \frac{i}{4} H_0^{(1)}(\omega |\mathbf{x} - \mathbf{y}|)$$

kernel is Helmholtz fundamental soln
a.k.a. free space Greens func



Jump relations: field limit as $\mathbf{x} \rightarrow \Gamma$ can depend on which side (\pm):


$$v^\pm = S\sigma \quad \text{no jump}$$

$$u^\pm = D\tau \pm \frac{1}{2}\tau \quad \text{jump}$$

S, D bdry integral ops w/ above kernels,
both bounded in $C(\Gamma) \rightarrow C^1(\Gamma)$

- From now fix $\Gamma = \partial\Omega$ i.e. densities live on obstacle boundary

Single obstacle




scatt. field $u = (\mathcal{D} - i\omega\mathcal{S})\tau$ “combined field repr.”

BC & JR: $A\tau := \left(\frac{1}{2}I + \underbrace{D - i\omega S}_{K}$) $\tau = -u^{\text{inc}}|_{\partial\Omega}$

2nd-kind IE ↗ K , kernel k weakly singular



Single obstacle



scatt. field $u = (\mathcal{D} - i\omega\mathcal{S})\tau$ “combined field repr.”
BC & JR: $A\tau := (\frac{1}{2}I + \underbrace{D - i\omega S}_{K}, \text{ kernel } k \text{ weakly singular})\tau = -u^{\text{inc}}|_{\partial\Omega}$
2nd-kind IE ↗


Smooth quadrature: $j = 1, \dots, N$
nodes $\mathbf{y}_j \in \partial\Omega$, weights w_j

Approx. by dense $N \times N$ linear system:



$$A\tau = b \quad \text{unknown density vector } \tau \approx \{\tau(\mathbf{y}_j)\}_{j=1}^N \quad (\text{Nyström '30})$$


Single obstacle



scatt. field $u = (\mathcal{D} - i\omega\mathcal{S})\tau$ “combined field repr.”
BC & JR: $A\tau := (\frac{1}{2}I + \underbrace{D - i\omega S}_{K}, \text{ kernel } k \text{ weakly singular})\tau = -u^{\text{inc}}|_{\partial\Omega}$
2nd-kind IE ↗

Smooth quadrature: $j = 1, \dots, N$
nodes $\mathbf{y}_j \in \partial\Omega$, weights w_j


Approx. by dense $N \times N$ linear system:



$$A\tau = b \quad \text{unknown density vector } \tau \approx \{\tau(\mathbf{y}_j)\}_{j=1}^N \quad (\text{Nyström '30})$$

- large $N > 10^4$: solve **iteratively** by applying A to vectors (GMRES)
- 2nd-kind \Rightarrow # iters indep. of N $\text{spec}(A)$ clusters at $\frac{1}{2}$, not near 0


Single obstacle



scatt. field $u = (\mathcal{D} - i\omega\mathcal{S})\tau$ “combined field repr.”
 BC & JR: $A\tau := (\frac{1}{2}I + \underbrace{D - i\omega S}_{K}, \text{ kernel } k \text{ weakly singular})\tau = -u^{\text{inc}}|_{\partial\Omega}$
 2nd-kind IE ↗

Smooth quadrature: $j = 1, \dots, N$
 nodes $\mathbf{y}_j \in \partial\Omega$, weights w_j

Approx. by dense $N \times N$ linear system:




$$A\tau = b \quad \text{unknown density vector } \tau \approx \{\tau(\mathbf{y}_j)\}_{j=1}^N \quad (\text{Nyström '30})$$

- large $N > 10^4$: solve **iteratively** by applying A to vectors (GMRES)
- 2nd-kind \Rightarrow # iters indep. of N $\text{spec}(A)$ clusters at $\frac{1}{2}$, not near 0

Distant panels easy: $A_{ij} = k(\mathbf{y}_i, \mathbf{y}_j)w_j$ high-order accurate
 Self / neighbor panels hard, esp. in 3D singularity e.g. $k \sim 1/r$ in 3D


Single obstacle



scatt. field $u = (\mathcal{D} - i\omega\mathcal{S})\tau$ “combined field repr.”
 BC & JR: $A\tau := (\frac{1}{2}I + \underbrace{D - i\omega S}_{K}, \text{ kernel } k \text{ weakly singular})\tau = -u^{\text{inc}}|_{\partial\Omega}$
 2nd-kind IE ↗

Smooth quadrature: $j = 1, \dots, N$
 nodes $\mathbf{y}_j \in \partial\Omega$, weights w_j

Approx. by dense $N \times N$ linear system:



$$A\tau = b \quad \text{unknown density vector } \tau \approx \{\tau(\mathbf{y}_j)\}_{j=1}^N \quad (\text{Nyström '30})$$

- large $N > 10^4$: solve **iteratively** by applying A to vectors (GMRES)
- 2nd-kind \Rightarrow # iters indep. of N $\text{spec}(A)$ clusters at $\frac{1}{2}$, not near 0

Distant panels easy: $A_{ij} = k(\mathbf{y}_i, \mathbf{y}_j)w_j$ high-order accurate

Self / neighbor panels hard, esp. in 3D singularity e.g. $k \sim 1/r$ in 3D

Exploit: applying A = evaluating u^+ due to τ 's interpolant density

Problem of near-boundary field evaluation

Simpler case: 2D interior Laplace ($\omega = 0$) $\partial\Omega$ param by $Z(s)$, $s \in [0, 2\pi)$


say want eval. $u = \mathcal{D}\tau$ $u = \operatorname{Re} v$, $v(z) = \frac{i}{2\pi} \int_{\partial\Omega} \frac{\tau(y)}{z - y} dy$

Problem of near-boundary field evaluation

Simpler case: 2D interior Laplace ($\omega = 0$) $\partial\Omega$ param by $Z(s)$, $s \in [0, 2\pi]$

say want eval. $u = \mathcal{D}\tau$ $u = \operatorname{Re} v$, $v(z) = \frac{i}{2\pi} \int_{\partial\Omega} \frac{\tau(y)}{z - y} dy$

Use periodic trapezoid rule (N nodes uniform in param s):




- exponential convergent, but rate arbitrarily slow as $z \rightarrow \partial\Omega$

Thm (B '12): rate = dist. of $Z^{-1}(z)$ from real axis in complex s plane

Why is close evaluation hard?


$$u = \operatorname{Re} v, \quad v(z) = \frac{i}{2\pi} \int_{\partial\Omega} \frac{\tau(y)}{z - y} dy, \quad \text{approx. by } N\text{-node quadr}$$



target $z \rightarrow \partial\Omega$: integrand non-smooth

Why is close evaluation hard?

$$u = \operatorname{Re} v, \quad v(z) = \frac{i}{2\pi} \int_{\partial\Omega} \frac{\tau(y)}{z - y} dy, \quad \text{approx. by } N\text{-node quadr}$$




target $z \rightarrow \partial\Omega$: integrand non-smooth

- increase N ? wasteful, always a z where error terrible
- adaptive quad? v. slow as $z \rightarrow \partial\Omega$ done afresh for each target point

Why is close evaluation hard?

$$u = \operatorname{Re} v, \quad v(z) = \frac{i}{2\pi} \int_{\partial\Omega} \frac{\tau(y)}{z - y} dy, \quad \text{approx. by } N\text{-node quadr}$$



target $z \rightarrow \partial\Omega$: integrand non-smooth

- increase N ? wasteful, always a z where error terrible
- adaptive quad? v. slow as $z \rightarrow \partial\Omega$ done afresh for each target point

2D case recently solved efficiently
(Helsing–Ojala '08, Helsing '12)

But want scheme which generalizes simply to 3D ...

Quadrature By eXpansion (QBX)

(B '12)
(Klöckner-B-Greengard-O'Neil '12)

$\tau, v|_{\partial\Omega}$ analytic \Rightarrow v extends analytically some dist. outside Ω


Quadrature By eXpansion (QBX)

(B '12)

(Klöckner-B-Greengard-O'Neil '12)

$\tau, v|_{\partial\Omega}$ analytic \Rightarrow v extends analytically some dist. outside Ω

Taylor exp. $v(z) = \sum_{n=0}^{\infty} c_n(z - z_0)$ rad. of conv. ρ takes you beyond $\partial\Omega$




Quadrature By eXpansion (QBX)

(B '12)
(Klöckner-B-Greengard-O'Neil '12)

$\tau, v|_{\partial\Omega}$ analytic $\Rightarrow v$ extends analytically some dist. outside Ω

Taylor exp. $v(z) = \sum_{n=0}^{\infty} c_n(z - z_0)^n$ rad. of conv. ρ takes you beyond $\partial\Omega$



- pick center z_0 about $2.5h$ from $\partial\Omega$
- eval. p terms via Cauchy,

$$c_n = \frac{v^{(n)}(z_0)}{n!} = \frac{i}{2\pi} \int_{\partial\Omega} \frac{\tau(y)}{(z - y)^{n+1}} dy$$

integrand more osc. \Rightarrow need βN nodes, $\beta \approx 4$
interpolate τ from original N


- eval. Taylor exp. in $|z - z_0| \leq R < \rho$
- repeat for z_0 's all around $\partial\Omega$

Quadrature By eXpansion (QBX)

(B '12)
(Klöckner-B-Greengard-O'Neil '12)

$\tau, v|_{\partial\Omega}$ analytic $\Rightarrow v$ extends analytically some dist. outside Ω

Taylor exp. $v(z) = \sum_{n=0}^{\infty} c_n(z - z_0)^n$ rad. of conv. ρ takes you beyond $\partial\Omega$



- pick center z_0 about $2.5h$ from $\partial\Omega$
- eval. p terms via Cauchy,

$$c_n = \frac{v^{(n)}(z_0)}{n!} = \frac{i}{2\pi} \int_{\partial\Omega} \frac{\tau(y)}{(z - y)^{n+1}} dy$$

integrand more osc. \Rightarrow need βN nodes, $\beta \approx 4$
interpolate τ from original N

- eval. Taylor exp. in $|z - z_0| \leq R < \rho$
- repeat for z_0 's all around $\partial\Omega$


Thm. (B) error $\leq C \left(\frac{R}{\rho}\right)^p + Cp \left(\frac{C\beta}{p}\right)^p e^{-C\beta}$ asymp. exponential conv. in p, β

QBX for close eval: interior Laplace BVP


As before, double-layer rep. $u = \mathcal{D}\tau$

exact soln $u(x, y) = e^y \cos x$

\log_{10} relative err, $N = 130$



using QBX, same N




refinement $\beta = 4$, order $p = 10$, typ.


QBX for close eval: interior Laplace BVP

As before, double-layer rep. $u = \mathcal{D}\tau$ exact soln $u(x, y) = e^y \cos x$

\log_{10} relative err, $N = 130$



using QBX, same N



refinement $\beta = 4$, order $p = 10$, typ.

Helmholtz ($\omega > 0$):

- Taylor \rightarrow local expansion $\sum_{|n| < p} c_n J_n(\omega r) e^{in\theta}$
- Cauchy \rightarrow Graf's addition theorem

QBX, high freq.
close eval. for
Helmholtz

100 λ diam.
700 λ perimeter




$N=9000$ unknowns
fill + solve 87 sec

FMM eval in 24 sec
(8×10^6 pts)

QBX in 28 sec
(2×10^5 pts)


rel. error $< 10^{-11}$

(B '12)



Fast algorithms for evaluation & applying A

Kernels (Green's funcs) special between “distant” boxes:




$$A_{ij} = k(\mathbf{x}_i, \mathbf{y}_j) \quad N^2 \text{ elements}$$

$$\text{rank}(A) := \{\# \text{ sing. vals.} > 10^{-16}\} = ?$$

Fast algorithms for evaluation & applying A

Kernels (Green's funcs) special between “distant” boxes:



$$A_{ij} = k(\mathbf{x}_i, \mathbf{y}_j) \quad N^2 \text{ elements}$$


$$\text{rank}(A) := \{\# \text{ sing. vals.} > 10^{-16}\} = ?$$

≈ 20 , indep of N ! low rank

Proof? $u|_{\text{targets}} \approx \text{loc. exp. in ball}$

Fast algorithms for evaluation & applying A

Kernels (Green's funcs) special between “distant” boxes:



$$A_{ij} = k(\mathbf{x}_i, \mathbf{y}_j) \quad N^2 \text{ elements}$$


$$\text{rank}(A) := \{\# \text{ sing. vals. } > 10^{-16}\} = ?$$

≈ 20 , indep of N ! low rank

Proof? $u|_{\text{targets}} \approx \text{loc. exp. in ball}$

Fast Multipole Method (FMM):

(Greengard–Rokhlin '87)



does dense matrix-vector prod. in $O(N)$

hierarchical subdivision (quad-tree)

distant interactions: LE's & multipoles

combine MP up tree, split LE down tree

Summary of panel-based $O(N)$ scattering solver

- (1) For each target panel, use QBX to precompute A_{ij} for self & neighbor source panels via a few nearby centers

Summary of panel-based $O(N)$ scattering solver

- (1) For each target panel, use QBX to precompute A_{ij} for self & neighbor source panels via a few nearby centers
- (2) Iteratively solve $A\tau = b$ apply A via FMM & precomputed entries

Summary of panel-based $O(N)$ scattering solver

- (1) For each target panel, use QBX to precompute A_{ij} for self & neighbor source panels via a few nearby centers
- (2) Iteratively solve $\mathbf{A}\boldsymbol{\tau} = \mathbf{b}$ apply \mathbf{A} via FMM & precomputed entries
- (3) Use density $\boldsymbol{\tau}$ to evaluate scattered field $u(x)$ where needed

Summary of panel-based $O(N)$ scattering solver


- (1) For each target panel, use QBX to precompute A_{ij} for self & neighbor source panels via a few nearby centers
- (2) Iteratively solve $\mathbf{A}\boldsymbol{\tau} = \mathbf{b}$ apply \mathbf{A} via FMM & precomputed entries
- (3) Use density $\boldsymbol{\tau}$ to evaluate scattered field $u(x)$ where needed

3D: panels $q \times q$ Gauss nodes

Local expansion $u(r, \theta, \phi) =$


$$\sum_{|n| \leq p} \sum_{m=-n}^n c_{nm} j_n(\omega r) Y_n^m(\theta, \phi)$$

spherical harmonic addn thm



- $p+1$ th order proven, $\boldsymbol{\tau} \in W^{p+3+\epsilon, 2}$ (Epstein–Greengard–Klöckner '12)

3D high-freq. torus scattering result



30 λ diameter
(1800 λ^2 surf. area)


$N \approx 145000$
 $q=8, p=10, \beta=4.5$

QBX quad 1.2 hr
GMRES 1 hr (57 its)
laptop (4-core i7)

relative error 10^{-5}


- FEM/FDTD at this high accuracy & freq. prohibitive
- Corners also handled in 2D; 3D edges/corners ongoing work ...

II. Quasi-periodic scattering from grating



u^{inc} , angle θ^i , wavevector (κ^i, k^i)

II. Quasi-periodic scattering from grating




u^{inc} , angle θ^i , wavevector (κ^i, k^i)

u^{inc} , hence solution, *quasi-periodic*:

$$u(x + d, y) = \alpha u(x, y)$$

↖ Bloch phase $\alpha = e^{i\kappa^i d}$

II. Quasi-periodic scattering from grating



Rayleigh–Bloch radiation condition for u :


set of QP plane waves, x -wavenumbers

$$\kappa_n := \kappa^i + 2\pi n/d$$


Only outwards-prop. or decaying:

$$u(x, y > y_0) = \sum_{n \in \mathbb{Z}} a_n e^{i(\kappa_n x + \sqrt{\omega^2 - \kappa_n^2} y)}$$


$$u(x, y < y_0) = \sum_{n \in \mathbb{Z}} b_n e^{i(\kappa_n x - \sqrt{\omega^2 - \kappa_n^2} y)}$$



Standard way to periodize integral equations



Standard way to periodize integral equations




- replaces A by $A_{\text{QP}} := \sum_n A_n$

Replace free-space kernel
 $G(x, y) := G(x, y; \mathbf{0})$ by
$$G_{\text{QP}}(x, y) := \sum_{n \in \mathbb{Z}} \alpha^n G(x, y - nd)$$

makes IE on $\partial\Omega$ solve full QP prob.

Standard way to periodize integral equations



Replace free-space kernel
 $G(x, y) := G(x, y; \mathbf{0})$ by
$$G_{\text{QP}}(x, y) := \sum_{n \in \mathbb{Z}} \alpha^n G(x, y - nd)$$


- replaces A by $A_{\text{QP}} := \sum_n A_n$ makes IE on $\partial\Omega$ solve full QP prob.

Issues:

- (1) G_{QP} slow to evaluate accurately (e.g. lattice sums)
much effort spent! (McPhedran, Linton, Kurkcu–Reitich, Bruno, ...)
- (2) Large-scale problems: never want to fill N^2 entries of dense A_{QP}
need something else if want fast algorithms
- (3) G_{QP} doesn't exist at, blows up near, “Wood anomalies”: **not robust!**
 (ω, θ^i) params where one or more Bragg angles along the grating (Wood 1902)

New low-rank periodizing scheme

(B–Greengard ’11)



replace “far” effect on u by


$$\text{basis } \sum_{m=1}^M c_m \varphi_m$$

each φ_n solves Helmholtz eqn

far \Rightarrow low rank $\Rightarrow M$ small

New low-rank periodizing scheme

(B-Greengard '11)



replace “far” effect on u by basis $\sum_{m=1}^M c_m \varphi_m$
each φ_n solves Helmholtz eqn
 $\text{far} \Rightarrow \text{low rank} \Rightarrow M \text{ small}$

Must now impose QP on walls: $u_R - \alpha u_L = 0, \quad u_{n,R} - \alpha u_{n,L} = 0$


2×2 block linear sys:

$$\begin{bmatrix} A & B \\ C & Q \end{bmatrix} \begin{bmatrix} \tau \\ \mathbf{c} \end{bmatrix} = \begin{bmatrix} -u^{\text{inc}}|_{\partial\Omega} \\ 0 \end{bmatrix} \quad \begin{aligned} \leftarrow \text{usual BC on } \partial\Omega \\ \leftarrow \text{QP wall conditions} \end{aligned}$$

cures 3 issues: robust, good w/ FMM, all free-space G , no lattice sums

New low-rank periodizing scheme

(B-Greengard '11)



replace “far” effect on u by basis $\sum_{m=1}^M c_m \varphi_m$
each φ_n solves Helmholtz eqn
 $\text{far} \Rightarrow \text{low rank} \Rightarrow M \text{ small}$


Must now impose QP on walls: $u_R - \alpha u_L = 0, \quad u_{n,R} - \alpha u_{n,L} = 0$

2×2 block linear sys:

$$\begin{bmatrix} A & B \\ C & Q \end{bmatrix} \begin{bmatrix} \tau \\ c \end{bmatrix} = \begin{bmatrix} -u^{\text{inc}}|_{\partial\Omega} \\ 0 \end{bmatrix} \quad \begin{array}{l} \leftarrow \text{usual BC on } \partial\Omega \\ \leftarrow \text{QP wall conditions} \end{array}$$

cures 3 issues: robust, good w/ FMM, all free-space G , no lattice sums


L, R unbounded \Rightarrow Fourier transform $y \leftrightarrow k$, on Sommerfeld contour:




$$G(x, y) = \frac{i}{4\pi} \int_{-\infty}^{\infty} e^{iky} \frac{e^{i\sqrt{\omega^2 - k^2}|x|}}{\sqrt{\omega^2 - k^2}} dk$$

k -quadrature exponentially convergent in M nodes (φ_m = complex plane waves)


Small demo (10-line code in MATLAB toolbox MPSpack, B–Betcke '09)



Small demo (10-line code in MATLAB toolbox MPSpack, B–Betcke '09)




Exponential convergence:




$$d = 1.6\lambda \quad N = 70 \quad M = 80 \quad \text{error } 10^{-14} \quad t_{\text{fill+dense solve}} = 0.04 \text{ s (laptop)}$$

Small demo (10-line code in MATLAB toolbox MPSpack, B–Betcke '09)





Exponential convergence:



$$d = 1.6\lambda \quad N = 70 \quad M = 80 \quad \text{error } 10^{-14} \quad t_{\text{fill+dense solve}} = 0.04 \text{ s (laptop)}$$



MOVIE 1

Dielectrics, interfaces, high aspect ratios



$$\begin{aligned} (\Delta + \omega^2)u &= 0 && \text{outside} \\ (\Delta + n^2\omega^2)u &= 0 && \text{inside} \\ u^+ - u^- &= -u^i && \left. \right\} \text{on } \partial\Omega \quad \text{transmission} \\ u_n^+ - u_n^- &= -u_n^i && \left. \right\} \text{(TM Maxwell)} \end{aligned}$$

Dielectrics, interfaces, high aspect ratios



$$(\Delta + \omega^2)u = 0 \quad \text{outside}$$

$$(\Delta + n^2\omega^2)u = 0 \quad \text{inside}$$

$$\begin{aligned} u^+ - u^- &= -u^i \\ u_n^+ - u_n^- &= -u_n^i \end{aligned} \quad \left. \right\} \text{on } \partial\Omega \quad \begin{array}{l} \text{transmission} \\ (\text{TM Maxwell}) \end{array}$$

2nd kind rep: (Müller '69, Rokhlin '83)

$$u = \begin{cases} \mathcal{D}\tau + \mathcal{S}\sigma & \text{inside} \\ \mathcal{D}_i\tau + \mathcal{S}_i\sigma & \text{outside} \end{cases}$$


A now 2×2 ; must periodize both layers

High aspect easy $M \sim N \sim$ height
unlike lattice sums (Nishimura '08)

32λ tall $N=700$ $M=300$ err 10^{-8} 4 s

Applications care about multiple incident angles

Power fractions scattered into each transmitted/reflected Bragg order:



$d = 1.6\lambda$ diel. obst. error 10^{-12} 3000 angles in 8 mins

- square-root type cusps at each Wood anomaly (dotted red)

Fast direct periodic scattering solver (Gillman–B ’12)

Iterative methods not ideal:

- start afresh for each RHS $f = -u^{\text{inc}}|_{\partial\Omega}$
- resonances, complex geom \rightarrow many iters

Fast direct periodic scattering solver (Gillman–B ’12)

Iterative methods not ideal:

- start afresh for each RHS $f = -u^{\text{inc}}|_{\partial\Omega}$
- resonances, complex geom \rightarrow many iters

Direct solve possible in $O(N)$:

nested compressed rep. of A^{-1}

exploits low-rank offdiag. blocks

can then apply A^{-1} v. cheaply

(Hackbusch, Rokhlin–Martinsson)

Fast direct periodic scattering solver (Gillman–B ’12)

Iterative methods not ideal:

- start afresh for each RHS $\mathbf{f} = -u^{\text{inc}}|_{\partial\Omega}$
- resonances, complex geom \rightarrow many iters

Direct solve possible in $O(N)$:

nested compressed rep. of \mathbf{A}^{-1}

exploits low-rank offdiag. blocks

can then apply \mathbf{A}^{-1} v. cheaply

(Hackbusch, Rokhlin–Martinsson)

Direct soln of periodic 2×2 sys:

$$\mathbf{c} = (Q - CA^{-1}B)^{-1}A^{-1}\mathbf{f}$$

$$\boldsymbol{\tau} = A^{-1}(\mathbf{f} - B\mathbf{c})$$

cf (Ho-Greengard ’12) redoes A_{QP}^{-1}

Fast direct periodic scattering solver (Gillman–B ’12)

Iterative methods not ideal:

- start afresh for each RHS $\mathbf{f} = -u^{\text{inc}}|_{\partial\Omega}$
- resonances, complex geom \rightarrow many iters

Direct solve possible in $O(N)$:


nested compressed rep. of \mathbf{A}^{-1}
exploits low-rank offdiag. blocks
can then apply \mathbf{A}^{-1} v. cheaply
(Hackbusch, Rokhlin–Martinsson)

Direct soln of periodic 2×2 sys:

$$\begin{aligned}\mathbf{c} &= (Q - CA^{-1}B)^{-1}A^{-1}\mathbf{f} \\ \boldsymbol{\tau} &= A^{-1}(\mathbf{f} - B\mathbf{c})\end{aligned}$$


cf (Ho-Greengard ’12) redoes A_{QP}^{-1}

E.g.: rough obstacle, Wood’s
needs $N = 10^5$ for error 10^{-10}



- 4 min for A^{-1} , 6 sec per θ^i : $600\times$ faster than even FMM + GMRES


Towards 3D: simpler “box scheme” & folding up



in box: $u = (\mathcal{D} - i\omega\mathcal{S})\tau + \sum_{m=1}^M c_m \varphi_m$
 near CFIEs loc. exp.
 above & below: $u =$ radiating Rayleigh–Bloch

BC:	A	B			$=$	τ	f
L–R QP:	C	Q				c	0
T cont:	C_T	Q_T	P_T			a	0
B cont:	C_B	Q_B	P_B			b	0

Towards 3D: simpler “box scheme” & folding up



in box: $u = (\mathcal{D} - i\omega\mathcal{S})\tau + \sum_{m=1}^M c_m \varphi_m$
 near CFIEs loc. exp.

above & below: $u =$ radiating Rayleigh–Bloch


BC:	A	B					=	$\begin{matrix} \mathbf{f} \\ 0 \\ 0 \\ 0 \end{matrix}$
L–R QP:	C	Q						
T cont:	C_T	Q_T	P_T					
B cont:	C_B	Q_B	P_B					

- Q not square, exp. ill. cond: bad to iterate on whole sys!

Eliminate QP unknowns: block Schur w/ small pseudoinverse

$$\begin{bmatrix} A & B \\ C & Q \end{bmatrix} \begin{bmatrix} \tau \\ \mathbf{c} \end{bmatrix} = \begin{bmatrix} \mathbf{b} \\ 0 \end{bmatrix} \quad \Leftrightarrow \quad \underbrace{(A - BQ^\dagger C)}_{A_{QP}} \tau = \mathbf{f}$$

Towards 3D: simpler “box scheme” & folding up



in box: $u = (\mathcal{D} - i\omega\mathcal{S})\tau + \sum_{m=1}^M c_m \varphi_m$
 near CFIEs loc. exp.
 above & below: $u =$ radiating Rayleigh–Bloch

BC:	A	B							
L–R QP:	C	Q							
T cont:	C_T	Q_T	P_T						
B cont:	C_B	Q_B	P_B						

$$\begin{matrix} \boldsymbol{\tau} \\ \mathbf{c} \\ \mathbf{a} \\ \mathbf{b} \end{matrix} = \begin{matrix} \mathbf{f} \\ 0 \\ 0 \\ 0 \end{matrix}$$

- Q not square, exp. ill. cond: bad to iterate on whole sys!


Eliminate QP unknowns: block Schur w/ small pseudoinverse

$$\begin{bmatrix} A & B \\ C & Q \end{bmatrix} \begin{bmatrix} \boldsymbol{\tau} \\ \mathbf{c} \end{bmatrix} = \begin{bmatrix} \mathbf{b} \\ 0 \end{bmatrix} \quad \Leftrightarrow \quad \underbrace{(A - BQ^\dagger C)}_{A_{QP}} \boldsymbol{\tau} = \mathbf{f}$$

- robust @ Wood's? Dirichlet/impedance half-space A_{QP} always exists

Prelim. results: 3D bi-periodic scattering

φ_m basis = 3D loc. exp. Match QP (4 faces) & rad. cond. (2 faces)



Dirichlet obstacles

$$d = 2.4\lambda$$

$$N = 4032 \text{ (small)}$$

$$\text{QBX } 20 \text{ s}$$

$$Q^\dagger \text{ 5 sec } (4000 \times 1339)$$

$$31 \text{ its } 60 \text{ s}$$

$$\text{error (flux) } 10^{-5}$$

- $O(N)$ scheme: small dense lin. alg. replaces messy lattice sums!
- high aspect simple via $\varphi_m = \text{plane waves}$
- Apps: modeling photonic crystal slab, metamaterials

Conclusions

Accurate, efficient, analysis-based scattering solvers, optimal $O(N)$ work

- QBX: high-order quadrature to apply IE ops on surfaces
 - simple new idea: use local expansion near boundary
- Periodic problems: use low-rank property of distant interactions
 - only free-space G , robust at Wood's, works with fast solvers

Ongoing/future: multilayer dielectrics, Neumann BC, Maxwell,
solar cells (w/ J. Liu), periodic Stokes

B, *subm.* SIAM J. Sci. Comp.

Klöckner–B–Greengard–O’Neil, *subm.*

Gillman–B, *subm.* J. Comput. Phys.

B–Greengard, BIT Numer. Math. (2011)

<http://math.dartmouth.edu/~ahb>

funding: NSF DMS-1216656

code: (try it!)

MPSpack toolbox for MATLAB

<http://code.google.com/p/mpspack>