

*Robust and efficient computation of  
two-dimensional photonic crystal band structure  
using second-kind integral equations*

LSU, March 1, 2010

Alex Barnett

Mathematics Department, Dartmouth College  
joint with Leslie Greengard (Courant)

# Photonic crystals

periodic dielectric structures

period  $\approx$  wavelength of light  $\approx 1\mu\text{m}$

control optical propagation in ways  
impossible in homogeneous media



(Joannopoulos group)

# Photonic crystals


periodic dielectric structures

period  $\approx$  wavelength of light  $\approx 1\mu\text{m}$

control optical propagation in ways impossible in homogeneous media




(Joannopoulos group)



2D lattice of cylinders (INFM, U. Pavia)

- e.g. ‘bandgap’ medium:  $\exists$  freqs. s.t.  
all waves evanescent (non-propagating)
- ‘insulators’ with embedded waveguides
  - unlike dielectric guides, sharp bends ok


# Photonic crystal examples



- Slab w/ 2D-periodic air holes couples to external dielectric guide  
manipulate guide dispersion to give  
v slow group velocity ( $c/300$ )

Si,  $\lambda = 1.6\mu\text{m}$  (Vlasov '05)


# Photonic crystal examples



Si,  $\lambda = 1.6\mu\text{m}$  (Vlasov '05)

- Full 3D bandgap (all polarizations)  
'Yablonovite' (cm scale) (Yablonovich '91)  
'woodpile'  $\lambda = 12\mu\text{m}$  (Lin *et al.* '98)  
'inverse opals' (spherical air 'bubbles')  
stacked slabs (built  $\lambda = 1.3\mu\text{m}$ , Qi *et al.* '04)
- complex geometry (not just cylinders!)

- Slab w/ 2D-periodic air holes  
couples to external dielectric guide  
manipulate guide dispersion to give  
 $v$  slow group velocity ( $c/300$ )



(Johnson *et al.* '00)


# Applications

Build **low-loss** optical signal paths on  $1\mu\text{m}$  scale:  
integrated optical devices, signal-processing,

Big goal: optical (*high* speed!) computing

*e.g.* high-Q resonators, couplers, junctions

channel-drop filter in 2D crystal




(Johnson *et al.* '00)

# Applications

Build **low-loss** optical signal paths on  $1\mu\text{m}$  scale:  
integrated optical devices, signal-processing,  
Big goal: optical (*high* speed!) computing  
*e.g.* high-Q resonators, couplers, junctions

**channel-drop filter in 2D crystal**



(Johnson *et al.* '00)


- Meta-materials *e.g.* negative refractive index ( $-1$  = 'perfect' lens)
- Solar cells and LEDs: control the density of states  
⇒ spontaneous emission/absorption rates, directions (S. Fan '97)



# Applications

Build **low-loss** optical signal paths on  $1\mu\text{m}$  scale:  
integrated optical devices, signal-processing,  
Big goal: optical (*high* speed!) computing  
*e.g.* high-Q resonators, couplers, junctions

**channel-drop filter in 2D crystal**



(Johnson *et al.* '00)

- Meta-materials *e.g.* negative refractive index ( $-1$  = 'perfect' lens)
- Solar cells and LEDs: control the density of states  
 $\Rightarrow$  spontaneous emission/absorption rates, directions (S. Fan '97)

## Common features

- piecewise-homogeneous dielectric media, wavenumber low
- each medium linear, may be dispersive *e.g.* metals (plasmons)
- manufacturing costly  $\Rightarrow$  accurate **numerical** modeling key



# Outline

1. Band structure: eigenmodes on a torus
2. Boundary integral equations
3. Periodizing: standard way & new way
4. Interpolation of bands
5. Software environment


# 2D dielectric crystal

( $z$ -invariant Maxwell, TM polarization)

unit cell  $U$       smooth inclusion  $\Omega \in U$ , refractive index  $n$

lattice  $\Lambda := \{m\mathbf{e}_1 + n\mathbf{e}_2 : n, m \in \mathbb{Z}\}$

dielectric inclusions  $\Omega_\Lambda := \{\mathbf{x} : \mathbf{x} + \mathbf{d} \in \Omega \text{ for some } \mathbf{d} \in \Lambda\}$




# 2D dielectric crystal

( $z$ -invariant Maxwell, TM polarization)

unit cell  $U$       smooth inclusion  $\Omega \in U$ , refractive index  $n$

lattice  $\Lambda := \{m\mathbf{e}_1 + n\mathbf{e}_2 : n, m \in \mathbb{Z}\}$

dielectric inclusions  $\Omega_\Lambda := \{\mathbf{x} : \mathbf{x} + \mathbf{d} \in \Omega \text{ for some } \mathbf{d} \in \Lambda\}$



scalar wave  $u : \mathbb{R}^2 \rightarrow \mathbb{C}$  obeys

**PDE (fixed frequency  $\omega$ ):**

$$(\Delta + n^2\omega^2)u = 0 \text{ in } \Omega_\Lambda$$

$$(\Delta + \omega^2)u = 0 \text{ in } \mathbb{R}^2 \setminus \Omega_\Lambda$$

**material matching conditions:**

$$u^+ - u^- = 0 \text{ on } \partial\Omega_\Lambda$$

$$u_n^+ - u_n^- = 0 \text{ on } \partial\Omega_\Lambda$$

# Bloch ‘theorem’

Solutions to PDE w/ periodic coeffs have form (or are sum of forms)

$$u(\mathbf{x}) = e^{i\mathbf{k}\cdot\mathbf{x}}\tilde{u}(\mathbf{x}), \quad \tilde{u} \text{ is periodic}$$

- called **Bloch waves**,  $\mathbf{k} \in \mathbb{R}^2$  Bloch wavevector

*‘When I started to think about it, I felt that the main problem was to explain how the electrons could sneak by all the ions in a metal... By straight Fourier analysis I found to my delight that the wave differed from the plane wave of free electrons only by a periodic modulation’*

(F. Bloch, 1928)

(indep. prediscovers by Hill 1877, Floquet 1883, Lyapunov 1892)


# Bloch wave and eigenvalue problem

- Bloch eigenvalues: set of  $(\omega, \mathbf{k})$  s.t. non-trivial Bloch waves  $u$  exist

# Bloch wave and eigenvalue problem

- Bloch eigenvalues: set of  $(\omega, \mathbf{k})$  s.t. non-trivial Bloch waves  $u$  exist

Example generalized eigenfunction  $u$  of Bloch wave form  $e^{i\mathbf{k}\cdot\mathbf{x}}\tilde{u}(\mathbf{x})$ :




Shown:  $\text{Re}[u]$  for

$$\omega = 5, \quad \mathbf{k} = (-0.39, 2.08)$$

# Bloch wave and eigenvalue problem

- Bloch eigenvalues: set of  $(\omega, \mathbf{k})$  s.t. non-trivial Bloch waves  $u$  exist

Example generalized eigenfunction  $u$  of Bloch wave form  $e^{i\mathbf{k}\cdot\mathbf{x}}\tilde{u}(\mathbf{x})$ :




Shown:  $\text{Re}[u]$  for

$$\omega = 5, \quad \mathbf{k} = (-0.39, 2.08)$$

$\mathbf{k}$  equiv. to  $\mathbf{k} + \mathbf{q}$ ,  $\forall \mathbf{q} \in 2\pi\Lambda^*$


$\Lambda^* = \text{dual (reciprocal) lattice}$

$\mathbf{k}$  lives on a torus, consider only *fundamental domain* (FD):





# Band structure



For each wavevector  $\mathbf{k} \in \text{FD}$ ,  
 $\exists$  discrete Bloch eigenvalues

$$\omega_1(\mathbf{k}) \leq \omega_2(\mathbf{k}) \leq \cdots \nearrow \infty$$

- form ‘sheets’ above the FD

note: conical at low freq  $\omega$

note: bandgap

- is most important property of photonic crystal for applications

# Recast problem on compact domain (torus)

- Bloch wave condition equiv. to quasi-periodic BCs on  $\partial U$


Require vanishing unit cell **discrepancy**:

$$f := u|_L - \alpha^{-1}u|_{L+\mathbf{e}_1} = 0$$

$$f' := u_n|_L - \alpha^{-1}u_n|_{L+\mathbf{e}_1} = 0$$

$$g := u|_B - \beta^{-1}u|_{B+\mathbf{e}_2} = 0$$

$$g' := u_n|_B - \beta^{-1}u_n|_{B+\mathbf{e}_2} = 0$$



Bloch phase parameters  $\alpha := e^{i\mathbf{k} \cdot \mathbf{e}_1}$ ,  $\beta := e^{i\mathbf{k} \cdot \mathbf{e}_2}$ ,  $|\alpha| = |\beta| = 1$

- Task: find Bloch eigenvalue triples  $(\omega, k_x, k_y)$ , i.e.  $(\omega, \alpha, \beta)$

# Main numerical approaches

## Time domain

- a) time-stepping on finite-difference grid (FDTD) (e.g. Yee '66)
  - low order (inaccurate); close freqs  $\rightarrow$  need large  $t$  (inefficient)

# Main numerical approaches

## Time domain

- a) time-stepping on finite-difference grid (FDTD) (e.g. Yee '66)  
low order (inaccurate); close freqs  $\rightarrow$  need large  $t$  (inefficient)

## Freq domain

- b) multiple-scattering, KKR, cylinders only (McPhedran, Moroz)
- c) Plane-wave method: all in Fourier space (Joannopoulos, Johnson, Sözüer)  
discont. dielectric  $\Rightarrow$  Gibbs phenom, slow ( $1/N$  or  $1/N^2$ ) convergence
- d) Finite element (FEM) discretization in  $U$  (Chew, Dobson, Dossou)  
better for discontinuity,  $N$  large, meshing complicated

# Main numerical approaches

## Time domain

- a) time-stepping on finite-difference grid (FDTD) (e.g. Yee '66)  
low order (inaccurate); close freqs  $\rightarrow$  need large  $t$  (inefficient)

## Freq domain

- b) multiple-scattering, KKR, cylinders only (McPhedran, Moroz)
- c) Plane-wave method: all in Fourier space (Joannopoulos, Johnson, Sözüer)  
discont. dielectric  $\Rightarrow$  Gibbs phenom, slow ( $1/N$  or  $1/N^2$ ) convergence
- d) Finite element (FEM) discretization in  $U$  (Chew, Dobson, Dossou)  
better for discontinuity,  $N$  large, meshing complicated
- e) Integral equations: formulate problem *on* the discontinuity  $\partial\Omega$   
reduced dimensionality (small  $N$ )  
high order (quadratures): high accuracy w/ small effort ( $\Rightarrow$  sensitivity analysis)  
scarcely used for band structure (Yuan '08)

# Potential theory

‘charge’ (sources of waves) distributed along curve  $\Gamma$  w/ density func.


single-, double-layer potentials,  $\mathbf{x} \in \mathbb{R}^2$ :

$$u(\mathbf{x}) = \int_{\Gamma} \Phi_{\omega}(\mathbf{x}, \mathbf{y}) \sigma(\mathbf{y}) ds_{\mathbf{y}} := (\mathcal{S}\sigma)(\mathbf{x})$$

$$v(\mathbf{x}) = \int_{\Gamma} \frac{\partial \Phi_{\omega}}{\partial n_{\mathbf{y}}}(\mathbf{x}, \mathbf{y}) \tau(\mathbf{y}) ds_{\mathbf{y}} := (\mathcal{D}\tau)(\mathbf{x})$$

$$\Phi_{\omega}(\mathbf{x}, \mathbf{y}) := \Phi_{\omega}(\mathbf{x} - \mathbf{y}) := \frac{i}{4} H_0^{(1)}(k|\mathbf{x} - \mathbf{y}|)$$

Helmholtz fundamental soln  
aka free space Greens func



# Potential theory

‘charge’ (sources of waves) distributed along curve  $\Gamma$  w/ density func.


single-, double-layer potentials,  $\mathbf{x} \in \mathbb{R}^2$ :

$$u(\mathbf{x}) = \int_{\Gamma} \Phi_{\omega}(\mathbf{x}, \mathbf{y}) \sigma(\mathbf{y}) ds_{\mathbf{y}} := (\mathcal{S}\sigma)(\mathbf{x})$$

$$v(\mathbf{x}) = \int_{\Gamma} \frac{\partial \Phi_{\omega}}{\partial n_{\mathbf{y}}}(\mathbf{x}, \mathbf{y}) \tau(\mathbf{y}) ds_{\mathbf{y}} := (\mathcal{D}\tau)(\mathbf{x})$$

$$\Phi_{\omega}(\mathbf{x}, \mathbf{y}) := \Phi_{\omega}(\mathbf{x} - \mathbf{y}) := \frac{i}{4} H_0^{(1)}(k|\mathbf{x} - \mathbf{y}|)$$

Helmholtz fundamental soln  
aka free space Greens func



**Jump relations:** limit as  $\mathbf{x} \rightarrow \Gamma$  may depend on which side ( $\pm$ ):

$$u^{\pm} = \mathcal{S}\sigma$$

$$u_n^{\pm} = \mathcal{D}^T \sigma \mp \frac{1}{2} \sigma$$

$$v^{\pm} = \mathcal{D}\tau \pm \frac{1}{2} \tau$$

$$v_n^{\pm} = T\tau$$

$\mathcal{S}, \mathcal{D}$  are integral ops with above kernels  
but defined on  $C(\Gamma) \rightarrow C(\Gamma)$


$T$  has kernel  $\frac{\partial^2 \Phi_{\omega}}{\partial n_{\mathbf{x}} \partial n_{\mathbf{y}}}(\mathbf{x}, \mathbf{y})$



# Integral equations for scattering (sketch)

e.g. Dirichlet obstacle: represent  $u = u^{\text{inc}} + \mathcal{D}\tau$


DLP on  $\partial\Omega$



# Integral equations for scattering (sketch)

e.g. Dirichlet obstacle: represent  $u = u^{\text{inc}} + \mathcal{D}\tau$

DLP on  $\partial\Omega$



**BC**  $0 = u^+ = u^{\text{inc}}|_{\partial\Omega} + (D + \frac{1}{2})\tau$

by JR3

integral eqn on  $\partial\Omega$ :  $(I + 2D)\tau = -2u^{\text{inc}}$

2nd-kind,  $D$  compact op so  $(I + 2D)$  sing. vals.  $\rightarrow 0$


Why important? when scale up...

condition # small, iterative solvers (GMRES) fast

# Integral equations for scattering (sketch)

e.g. Dirichlet obstacle: represent  $u = u^{\text{inc}} + \mathcal{D}\tau$

DLP on  $\partial\Omega$



**BC**  $0 = u^+ = u^{\text{inc}}|_{\partial\Omega} + (D + \frac{1}{2})\tau$

by JR3

integral eqn on  $\partial\Omega$ :  $(I + 2D)\tau = -2u^{\text{inc}}$

2nd-kind,  $D$  compact op so  $(I + 2D)$  sing. vals.  $\nrightarrow 0$

Why important? when scale up...

condition # small, iterative solvers (GMRES) fast

Quadrature scheme: choose  $N$  nodes  $\mathbf{y}_j \in \partial\Omega$ , weights  $w_j$


Nyström discretization:  $N$ -by- $N$  linear system for vector  $\{\tau_k^{(N)}\}_{k=1}^N$

$$\tau_k^{(N)} + 2 \sum_{j=1}^N w_j D(\mathbf{y}_k, \mathbf{y}_j) \tau_j^{(N)} = -2u^{\text{inc}}(\mathbf{y}_k), \quad k = 1, \dots, N$$

# Integral equations for scattering (sketch)

e.g. Dirichlet obstacle: represent  $u = u^{\text{inc}} + \mathcal{D}\tau$

DLP on  $\partial\Omega$



**BC**  $0 = u^+ = u^{\text{inc}}|_{\partial\Omega} + (D + \frac{1}{2})\tau$

by JR3

integral eqn on  $\partial\Omega$ :  $(I + 2D)\tau = -2u^{\text{inc}}$

2nd-kind,  $D$  compact op so  $(I + 2D)$  sing. vals.  $\nrightarrow 0$

Why important? when scale up...

condition # small, iterative solvers (GMRES) fast

Quadrature scheme: choose  $N$  nodes  $\mathbf{y}_j \in \partial\Omega$ , weights  $w_j$


Nyström discretization:  $N$ -by- $N$  linear system for vector  $\{\tau_k^{(N)}\}_{k=1}^N$

$$\tau_k^{(N)} + 2 \sum_{j=1}^N w_j D(\mathbf{y}_k, \mathbf{y}_j) \tau_j^{(N)} = -2u^{\text{inc}}(\mathbf{y}_k), \quad k = 1, \dots, N$$

Thm: (Anselone, Kress)  $\|\tau^{(N)} - \tau\|_{\infty}$  converges at *same rate* as quadrature scheme for the true integrand  $D(\mathbf{y}, \cdot)\tau$ .


- Analytic curve & data, periodic trapezoid rule: spectral convergence
- e.g. above:  $N = 60$  enough for  $10^{-6}$  error,  $N = 100$  for  $10^{-12}$
- error =  $O(e^{-\gamma N})$ , rate  $\gamma \approx$  distance to nearest singularity of  $\tau$  in  $\mathbb{C}$

# Dielectric (transmission) scattering



Represent  $u = u^{\text{inc}} + \mathcal{D}\tau + \mathcal{S}\sigma$  outside wavenumber  $\omega$   
 $u = \mathcal{D}_i\tau + \mathcal{S}_i\sigma$  inside wavenumber  $n\omega$

# Dielectric (transmission) scattering



Represent  $u = u^{\text{inc}} + \mathcal{D}\tau + \mathcal{S}\sigma$  outside wavenumber  $\omega$   
 $u = \mathcal{D}_i\tau + \mathcal{S}_i\sigma$  inside wavenumber  $n\omega$

mismatch on  $\partial\Omega$ :  $h := u^+ - u^-$ ,  $h' := u_n^+ - u_n^-$

BCs: mismatch  $m := [h; h']$  vanishes, use JRs...

$$\begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} u^{\text{inc}}|_{\partial\Omega} \\ u_n^{\text{inc}}|_{\partial\Omega} \end{bmatrix} + \underbrace{\left( \begin{bmatrix} I & 0 \\ 0 & I \end{bmatrix} + \begin{bmatrix} D - D_i & S_i - S \\ T - T_i & D_i^T - D^T \end{bmatrix} \right)}_A \underbrace{\begin{bmatrix} \tau \\ -\sigma \end{bmatrix}}_{\eta}$$

block 2nd-kind

$A$  maps densities to their effect on mismatch

- hypersingular part of  $T$  cancels, so  $A = \text{Id} + \text{compact}$  (Rokhlin '83)
- kernel weakly singular, but exists spectral product quadrature  
 for  $f(s) + \log(4 \sin^2 \frac{s}{2})g(s)$ ,  $f, g$  analytic  $2\pi$ -periodic (Kress '91)

# The standard way to periodize

replace kernel  $\Phi_\omega(\mathbf{x})$  by  $\Phi_{\omega,\text{QP}}(\mathbf{x}) := \sum_{m,n \in \mathbb{Z}} \alpha^m \beta^n \Phi(\mathbf{x} - m\mathbf{e}_1 - n\mathbf{e}_2)$

thus integral operator  $A$  becomes  $A_{\text{QP}}$



# The standard way to periodize

replace kernel  $\Phi_\omega(\mathbf{x})$  by  $\Phi_{\omega, \text{QP}}(\mathbf{x}) := \sum_{m,n \in \mathbb{Z}} \alpha^m \beta^n \Phi(\mathbf{x} - m\mathbf{e}_1 - n\mathbf{e}_2)$

thus integral operator  $A$  becomes  $A_{\text{QP}}$

- is standard approach for scattering from gratings (McPhedran, Otani)
- seems natural for band structure problem . . . not yet used in literature

# The standard way to periodize

replace kernel  $\Phi_\omega(\mathbf{x})$  by  $\Phi_{\omega, \text{QP}}(\mathbf{x}) := \sum_{m,n \in \mathbb{Z}} \alpha^m \beta^n \Phi(\mathbf{x} - m\mathbf{e}_1 - n\mathbf{e}_2)$

thus integral operator  $A$  becomes  $A_{\text{QP}}$

- is standard approach for scattering from gratings (McPhedran, Otani)
- seems natural for band structure problem ... not yet used in literature

**Theorem** (*integral formulation of band structure*) :

|   |
|---|
| If $A_{\text{QP}}$ exists, $\text{Nul } A_{\text{QP}} \neq \{0\} \iff (\omega, k_x, k_y)$ is eigenvalue |
|---|

note: no  $u^{\text{inc}}$  since eigenvalue problem homogeneous

proof: non-physical fields are a swapped-media transmission BVP

# The standard way to periodize

replace kernel  $\Phi_\omega(\mathbf{x})$  by  $\Phi_{\omega, \text{QP}}(\mathbf{x}) := \sum_{m,n \in \mathbb{Z}} \alpha^m \beta^n \Phi(\mathbf{x} - m\mathbf{e}_1 - n\mathbf{e}_2)$

thus integral operator  $A$  becomes  $A_{\text{QP}}$

- is standard approach for scattering from gratings (McPhedran, Otani)
- seems natural for band structure problem ... not yet used in literature

**Theorem** (*integral formulation of band structure*) :

If  $A_{\text{QP}}$  exists,  $\text{Nul } A_{\text{QP}} \neq \{0\} \iff (\omega, k_x, k_y)$  is eigenvalue

note: no  $u^{\text{inc}}$  since eigenvalue problem homogeneous

proof: non-physical fields are a swapped-media transmission BVP

**Not** a robust method:  $A_{\text{QP}}$  does **not exist** for certain parameters  $(\omega, k_x, k_y)$   
since there  $\Phi_{\omega, \text{QP}}(\mathbf{x}) \rightarrow \infty, \forall \mathbf{x}$

why...?

# Failure at spurious resonances

$\Phi_{\omega, \text{QP}}(\mathbf{x})$  is Helmholtz Greens function in *empty* (index 1) torus

$$= \frac{1}{\text{Vol}(U)} \sum_{\mathbf{q} \in 2\pi\Lambda^*} \frac{e^{i(\mathbf{k}+\mathbf{q}) \cdot \mathbf{x}}}{\omega^2 - |\mathbf{k} + \mathbf{q}|^2} \quad \text{spectral representation on torus}$$




has simple pole wherever  $(\omega, k_x, k_y)$  is eigenvalue of empty torus...  
but physical field  $u$  well-behaved here: breakdown is **non-physical**!

# Failure at spurious resonances

$\Phi_{\omega, \text{QP}}(\mathbf{x})$  is Helmholtz Greens function in *empty* (index 1) torus

$$= \frac{1}{\text{Vol}(U)} \sum_{\mathbf{q} \in 2\pi\Lambda^*} \frac{e^{i(\mathbf{k}+\mathbf{q}) \cdot \mathbf{x}}}{\omega^2 - |\mathbf{k} + \mathbf{q}|^2} \quad \text{spectral representation on torus}$$

has simple pole wherever  $(\omega, k_x, k_y)$  is eigenvalue of empty torus...  
but physical field  $u$  well-behaved here: breakdown is **non-physical**!



# Our cure: robust way to periodize

represent  $u = \mathcal{D}\tau + \mathcal{S}\sigma +$  (densities  $\xi$  on walls of  $U$ ) outside



can enforce mismatch  $m = 0$



can enforce discrepancy  $d := [f; f'; g; g'] = 0$

## Our cure: robust way to periodize

represent  $u = \mathcal{D}\tau + \mathcal{S}\sigma +$  (densities  $\xi$  on walls of  $U$ )      outside

$\uparrow$                            $\uparrow$

can enforce mismatch  $m = 0$       can enforce discrepancy  $d := [f; f'; g; g'] = 0$

In block operator form

$$\underbrace{\begin{bmatrix} A & B \\ C & Q \end{bmatrix}}_M \begin{bmatrix} \eta \\ \xi \end{bmatrix} = \begin{bmatrix} m \\ d \end{bmatrix}$$

- added extra degrees of freedom (a small #, indep. of complexity of  $\Omega$ )



# Our cure: robust way to periodize

represent  $u = \mathcal{D}\tau + \mathcal{S}\sigma +$  (densities  $\xi$  on walls of  $U$ ) outside

can enforce mismatch  $m = 0$

↑  
can enforce discrepancy  $d := [f; f'; g; g'] = 0$

In block operator form

$$\underbrace{\begin{bmatrix} A & B \\ C & Q \end{bmatrix}}_M \begin{bmatrix} \eta \\ \xi \end{bmatrix} = \begin{bmatrix} m \\ d \end{bmatrix}$$

- added extra degrees of freedom (a small #, indep. of complexity of  $\Omega$ )
- gain robustness: no matrix element blow-up at spurious resonances

## Observe:

$$\text{Nul } M \neq \{0\} \quad \Leftrightarrow \quad (\omega, k_x, k_y) \text{ Bloch eigenvalue}$$

- idea of extra sources of waves not new (*e.g.* Hafner '02)
- what is new:  $M = \text{Id} + \text{compact}$       ideal for large-scale, iterative, FMM

# How choose new densities on unit cell walls?

- to control 4 discrepancies ( $f, f', g, g'$ )


need 4 densities  $\xi = [\tau_L; \sigma_L; \tau_B; \sigma_B]$

$$Q = \frac{1}{2}\text{Id} + (\text{self-interactions}) + (\text{other interactions})$$

JRs


$$\sigma_L \rightarrow u|_L$$

$$\sigma_L \rightarrow u|_B$$




# How choose new densities on unit cell walls?

- to control 4 discrepancies ( $f, f', g, g'$ )  
 need 4 densities  $\xi = [\tau_L; \sigma_L; \tau_B; \sigma_B]$   
 $Q = \frac{1}{2}\text{Id} + (\text{self-interactions}) + (\text{other interactions})$   
 $\text{JR}s \quad \sigma_L \rightarrow u|_L \quad \sigma_L \rightarrow u|_B$




- add phased **ghost** copies on other 2 walls  
 recall  $f := u|_L - \alpha^{-1}u|_{L+\mathbf{e}_1}$   
 effect of  $\sigma_L$  on  $u_n|_L$   
 effect of  $\alpha\sigma_L$  on  $\alpha^{-1}u_n|_{L+\mathbf{e}_1}$  } **cancel** apart from Id






# How choose new densities on unit cell walls?


- to control 4 discrepancies ( $f, f', g, g'$ )  
need 4 densities  $\xi = [\tau_L; \sigma_L; \tau_B; \sigma_B]$   
 $Q = \frac{1}{2}\text{Id} + (\text{self-interactions}) + (\text{other interactions})$   
JRs  $\sigma_L \rightarrow u|_L$   $\sigma_L \rightarrow u|_B$



- add phased **ghost** copies on other 2 walls  
recall  $f := u|_L - \alpha^{-1}u|_{L+\mathbf{e}_1}$   
effect of  $\sigma_L$  on  $u_n|_L$   
effect of  $\alpha\sigma_L$  on  $\alpha^{-1}u_n|_{L+\mathbf{e}_1}$  } **cancel** apart from Id




- add more ‘sticking-out’ ghost images  
effect of  on  $u_n|_L$   
effect of  $\alpha$   on  $\alpha^{-1}u_n|_{L+\mathbf{e}_1}$  } **cancel** apart from Id  
 $\Rightarrow$  all corner interactions vanish!



- result:  $Q = I + (\text{interactions of distance } \geq 1)$   
 $\Rightarrow$  low rank, rapid convergence: 20-pt Gauss quadr. on  $L, B \Rightarrow 10^{-12}$  error


# Full scheme

Finally we add 3x3 phased image copies of densities on  $\partial\Omega$ , giving:



# Full scheme

Finally we add 3x3 phased image copies of densities on  $\partial\Omega$ , giving:




- Careful cancellations:  $B, C, Q$  have only interactions of distance  $\geq 1$
- Large dist increases convergence **rate**, i.e. large  $c$  in error =  $O(e^{-cN})$

*Philosophy: sum neighboring image sources directly  
so fields due to remainder of lattice have distant singularities*

# Error convergence


$\log_{10}$  min sing. val  $M$  for known Bloch eigenvalue (should be zero):



Note: is eigenvalue error  
up to  $O(1)$  const

$$\omega = 5, \mathbf{k} \approx (-0.39, 2.08)$$

mode:



- spectral (exponential) convergence in inclusion & wall # dofs


# Crude results: small inclusion

band structure: simply plot log min sing. val. of  $M$  vs  $(\omega, k_x, k_y) \dots$



# Crude results: small inclusion


band structure: simply plot log min sing. val. of  $M$  vs  $(\omega, k_x, k_y) \dots$



0.1 sec per eval  
pre-store  $\alpha, \beta$  coeffs  
30 sec per  
const- $\omega$  slice  
 $24 \times 24$  evals

- errors  $10^{-9}$  for 40 pts on  $\partial\Omega$ , 20 on each wall (total  $N = 160$ )


# Large inclusion passing through unit cell




As  $\text{dist}(\Omega, \partial U) \rightarrow 0$  standard quadrature v. poor

- fix via adaptive quadrature of Lagrange interpolant
- faster: project wall densities onto J-expansion using Graf addition thm (needs  $N=35$  per wall)


# Large inclusion passing through unit cell



- As  $\text{dist}(\Omega, \partial U) \rightarrow 0$  standard quadrature v. poor
- fix via adaptive quadrature of Lagrange interpolant
  - faster: project wall densities onto J-expansion using Graf addition thm (needs  $N=35$  per wall)



Amazingly (due to far singularities), J-exp analytically continues the field to outside  $U$ :




$$\omega = 4.47$$

$$\mathbf{k} \approx (0.17, 2.11)$$


$$n=1 \text{ inside}$$

$$n=3.3 \text{ outside}$$


# Large inclusion passing through unit cell



- As  $\text{dist}(\Omega, \partial U) \rightarrow 0$  standard quadrature v. poor
- fix via adaptive quadrature of Lagrange interpolant
  - faster: project wall densities onto J-expansion using Graf addition thm (needs  $N=35$  per wall)



Amazingly (due to far singularities), J-exp analytically continues the field to outside  $U$ :



$$\omega = 4.47$$

$$\mathbf{k} \approx (0.17, 2.11)$$

$$n=1 \text{ inside}$$

$$n=3.3 \text{ outside}$$

Sampling fine 3D grid is crude & slow: how find bands to spectral acc?



# Interpolation across the Brillouin zone

How find eigenvalue sheets in the volume  $S^1 \times S^1 \times (0, \omega_{\max})$  ?

# Interpolation across the Brillouin zone

How find eigenvalue sheets in the volume  $S^1 \times S^1 \times (0, \omega_{\max})$  ?

E.g. iterative rootfinding on min sing. val. of  $M$ :





$\omega$ -slice  
( $\alpha, \beta$  fixed)

- slow (20 its per root), unreliable (misses nearby roots)

# Interpolation across the Brillouin zone

How find eigenvalue sheets in the volume  $S^1 \times S^1 \times (0, \omega_{\max})$  ?

E.g. iterative rootfinding on min sing. val. of  $M$ :



$\omega$ -slice  
( $\alpha, \beta$  fixed)

- slow (20 its per root), unreliable (misses nearby roots)

Realize:  $M = I + (\text{cpt op-valued analytic func of } \omega, k_x \text{ and } k_y)$

$\det M$  is a **Fredholm determinant**, also analytic


- rootfinding a real-analytic function is nice...

(J. Boyd '02)

# Spectral rootfinding of analytic functions

(Boyd '02)

$f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $2\pi$ -periodic






# Spectral rootfinding of analytic functions

(Boyd '02)

$f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $2\pi$ -periodic

use trigonometric poly interpolant



$$f(\theta) \approx \sum_{n=-N}^N c_n e^{in\theta}$$

exponentially convergent in  $N$


width of strip about  $\mathbb{R}$  in which  $f$  analytic

# Spectral rootfinding of analytic functions

(Boyd '02)

$f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $2\pi$ -periodic

use trigonometric poly interpolant



$$f(\theta) \approx \sum_{n=-N}^N c_n e^{in\theta}$$

exponentially convergent in  $N$

width of strip about  $\mathbb{R}$  in which  $f$  analytic


- $c_n$  found by FFT of grid samples  $f(\pi n/N)$ ,  $n = 1, \dots, 2N$

# Spectral rootfinding of analytic functions

(Boyd '02)

$f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $2\pi$ -periodic

use trigonometric poly interpolant



$$f(\theta) \approx \sum_{n=-N}^N c_n e^{in\theta}$$

exponentially convergent in  $N$

width of strip about  $\mathbb{R}$  in which  $f$  analytic

- $c_n$  found by FFT of grid samples  $f(\pi n/N)$ ,  $n = 1, \dots, 2N$

map  $e^{i\theta} = z \in \mathbb{C}$

$\Rightarrow$  Laurent  $q(z) = \sum_{n=-N}^N c_n z^n$

roots of  $f$  lie on  $|z| = 1$


has roots *near*  $|z| = 1$

# Spectral rootfinding of analytic functions

(Boyd '02)

$f : \mathbb{R} \rightarrow \mathbb{R}$ ,  $2\pi$ -periodic

use trigonometric poly interpolant



$$f(\theta) \approx \sum_{n=-N}^N c_n e^{in\theta}$$

exponentially convergent in  $N$

width of strip about  $\mathbb{R}$  in which  $f$  analytic

- $c_n$  found by FFT of grid samples  $f(\pi n/N)$ ,  $n = 1, \dots, 2N$

map  $e^{i\theta} = z \in \mathbb{C}$

roots of  $f$  lie on  $|z| = 1$

$\Rightarrow$  Laurent  $q(z) = \sum_{n=-N}^N c_n z^n$

has roots *near*  $|z| = 1$


“Degree doubling”:  $z^N q(z)$  is degree- $2N$  poly, so...

- use Matlab `roots`    QR for eigvals of companion matrix,  $O(N^3)$  but v. stable
- extract the angles  $\theta$  of roots near unit circle

(Boyd nonlin EVP; Trefethen-Battles '06 `chebfun`)

# Rootfinding $\det M$ in the $\omega$ direction


$\det M(\omega, k_x, k_y)$  not periodic in  $\omega$ : map  $\omega = \omega_0 + a \cos \theta$       periodic  $\theta$   
this is Chebyshev interpolation on interval  $[\omega_0 - a, \omega_0 + a]$




Fix  $\mathbf{k}$ , eval  $\det M$  at Cheby pts, get  $\omega_j(\mathbf{k})$  in interval  
25 evals covers  $\omega \in [4, 6]$ , i.e. 10-20 evals per root found

# Rootfinding $\det M$ in the $\omega$ direction


$\det M(\omega, k_x, k_y)$  not periodic in  $\omega$ : map  $\omega = \omega_0 + a \cos \theta$  **periodic  $\theta$**   
this is **Chebyshev interpolation** on interval  $[\omega_0 - a, \omega_0 + a]$



Fix  $\mathbf{k}$ , eval  $\det M$  at Cheby pts, get  $\omega_j(\mathbf{k})$  in interval  
25 evals covers  $\omega \in [4, 6]$ , i.e. 10-20 evals per root found




Also analytic in  $k_x, k_y \Rightarrow$  interpolate in 3D!



Robust spectrally-accurate bands via small # grid evals  
e.g.  $25 \times 24 \times 24$  for  $\omega \in [4, 6]$  and whole Brillouin zone, error  $10^{-8}$

# Band structure to spectral accuracy



$n=0.3$  inside  
 $n=1$  outside  
large inclusion

eval only  $24 \times 24$  samples in  $\mathbf{k}$   
but contains much finer details

$10^{-8}$  errors, 1 hour on laptop

- Note: eigenvalues  $\omega_j(\mathbf{k})$  are **not** analytic!  
 $\exists$  conical (diabolical) points ...interpolates poorly

- like level set method: handle smooth func

*movie 1*

# Software environment (teaser)



# Software environment (teaser)

MPSPack: object-oriented 2D PDE toolbox in Matlab (B-Betcke '09)

- implements above & more: Helmholtz, Laplace, scattering
- intuitive interface: curves, domains, basis sets, problems, are **objects**

E.g.

# Software environment (teaser)

MPSPack: object-oriented 2D PDE toolbox in Matlab (B-Betcke '09)

- implements above & more: Helmholtz, Laplace, scattering
- intuitive interface: curves, domains, basis sets, problems, are **objects**

E.g.

```
s=segment.radialfunc(100,{@(q)1+.3*cos(3*q),@(q)-.9*sin(3*q)});  
plot(s);  
title('piecewise analytic curves')
```

# Software environment (teaser)

MPSPack: object-oriented 2D PDE toolbox in Matlab (B-Betcke '09)

- implements above & more: Helmholtz, Laplace, scattering
- intuitive interface: curves, domains, basis sets, problems, are **objects**

E.g.

```
s=segment.radialfunc(100,{@(q)1+.3*cos(3*q),@(q)-.9*sin(3*q)});  
    piecewise analytic curves
```

```
d = domain([], [], s, -1);  
    make exterior domain using segment
```

# Software environment (teaser)


MPSPack: object-oriented 2D PDE toolbox in Matlab (B-Betcke '09)

- implements above & more: Helmholtz, Laplace, scattering
- intuitive interface: curves, domains, basis sets, problems, are **objects**

E.g.

```
s=segment.radialfunc(100,{@(q)1+.3*cos(3*q),@(q)-.9*sin(3*q)},  
    piecewise analytic curves
```

```
d = domain([], [], s, -1);  
    make exterior domain using segment
```



# Software environment (teaser)

MPSpack: object-oriented 2D PDE toolbox in Matlab (B-Betcke '09)


- implements above & more: Helmholtz, Laplace, scattering
- intuitive interface: curves, domains, basis sets, problems, are **objects**

E.g.

```
s=segment.radialfunc(100,{@(q)1+.3*cos(3*q),@(q)-.9*sin(3*q)},  
    piecewise analytic curves
```

```
d = domain([], [], s, -1);  
    make exterior domain using segment
```

```
d.setbc(1, 'N', []);  
    Neumann BCs on exterior
```



# Software environment (teaser)

MPSpack: object-oriented 2D PDE toolbox in Matlab (B-Betcke '09)

- implements above & more: Helmholtz, Laplace, scattering
- intuitive interface: curves, domains, basis sets, problems, are **objects**


E.g.

```
s=segment.radialfunc(100,{@(q)1+.3*cos(3*q),@(q)-.9*sin(3*q)},  
    piecewise analytic curves
```

```
d = domain([], [], s, -1);  
    make exterior domain using segment
```

```
d.setbc(1, 'N', []);  
    Neumann BCs on exterior
```

```
d.addmfsbasis(s, 200, struct('tau',0.05));  
    choose basis set for solution
```



# Software environment (teaser)

MPSpack: object-oriented 2D PDE toolbox in Matlab (B-Betcke '09)

- implements above & more: Helmholtz, Laplace, scattering
- intuitive interface: curves, domains, basis sets, problems, are **objects**


E.g.

```
s=segment.radialfunc(100,{@(q)1+.3*cos(3*q),@(q)-.9*sin(3*q)},  
    piecewise analytic curves
```

```
d = domain([], [], s, -1);  
    make exterior domain using segment
```

```
d.setbc(1, 'N', []);  
    Neumann BCs on exterior
```

```
d.addmfsbasis(s, 200, struct('tau',0.05));  
    choose basis set for solution
```



```
p = scattering(d, [1]);
```

make a scattering problem from domain  $\mathcal{d}$



```
p = scattering(d, []);
```

make a scattering problem from domain  $\bar{d}$

```
p.setoverallwavenumber(30); p.setincidentwave(pi/6);
```

```
p = scattering(d, []);
```

make a scattering problem from domain  $\mathcal{d}$

```
p.setoverallwavenumber(30); p.setincidentwave(pi/6);
```

```
p.solvecoeffs; p.bcresidualnorm
```

fills matrix, solves in 0.1 sec,  $L^2$  error  $6 \times 10^{-9}$

```
p = scattering(d, []);
```


make a scattering problem from domain  $d$

```
p.setoverallwavenumber(30); p.setincidentwave(pi/6);
```

```
p.solvecoeffs; p.bcreidualnorm
```

fills matrix, solves in 0.1 sec,  $L^2$  error  $6 \times 10^{-9}$

```
p.showfullfield(struct('bb',[-2 2 -2 2]));
```



```
p = scattering(d, []);
```


make a scattering problem from domain  $d$

```
p.setoverallwavenumber(30); p.setincidentwave(pi/6);
```

```
p.solvecoeffs; p.bcresidualnorm
```

fills matrix, solves in 0.1 sec,  $L^2$  error  $6 \times 10^{-9}$

```
p.showfullfield(struct('bb', [-2 2 -2 2]));
```




- was easy case: 8 lines (could have done in 80 lines of Matlab)
- multiple (sub)domains: basis, quadrature, bookkeeping hidden  
e.g. dielectric band structure still only 20 lines of code
- human-readable, rapid to code, sensible defaults (you can change)

To do: automatic meshing, Dirichlet EVP, ...

# Current work: grating scattering

Quasi-periodize in  $x$ -direction only: layer-potentials infinite in  $y \dots$



- $N = 50$  unknowns on inclusion,  $M = 200$  unknowns to periodize
- accuracy  $10^{-13}$

# Conclusions

- efficient 2nd-kind integral equations for photonic crystal EVP
- periodize via small # extra degrees of freedom on cell walls
- more robust and flexible than quasi-periodic Greens function:
  - no spurious blow-up at empty resonances
  - extends simply to 3D (unlike lattice sums)
- interpolate Fredholm det, not Bloch eigenvalues themselves

## Future:

- 3D; drop in FMM for inclusion; gratings with substrate ...

### code:

<http://code.google.com/p/mpspack>

### funding:

NSF DMS-0507614  
DMS-0811005

### Preprints, talks, movies:


<http://math.dartmouth.edu/~ahb>

made with: Linux, L<sup>A</sup>T<sub>E</sub>X, Prosper

# EXTRA SLIDES

# Equal-frequency curves

Complexity of const- $\omega$  slice across Brillouin zone:



- only  $24 \times 24$  evaluations of  $\det M$ , Boyd's spectral rootfinding
- Apps: Snell's Law for reflection off semi- $\infty$  crystal