Taylor and Maclaurin Series

October 20, 2006

Recall

ullet The Taylor series of the function f at a is

$$f(x) = f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \cdots$$

• The Maclaurin series

$$f(x) = f(0) + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \cdots$$

When does a function equals its Taylor series?

Consider the partial sums of the Taylor series:

$$T_n(x) = f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \dots + \frac{f^n(a)}{n!}(x-a)^n.$$

- T_n is called the nth-degree Taylor polynomial of f at a.
- Then f equals its Taylor series if $f(x) = \lim_{n \to 0} T_n(x)$.

The remainder of the Taylor series

• Let
$$R_n(x) = f(x) - T_n(x)$$

• If

$$\lim_{n \to 0} R_n(x) = 0$$

for |x-a| < R, then f is equal to the sum of its Taylor series on the interval |x-a| < R.

Useful formulas

• (Taylor's inequality) If $|f^{(n+1)}(x)| \leq M$ for $|x-a| \leq d$, then the remainder $R_n(x)$ satisfies the inequality

$$|R_n(x)| \le \frac{M}{(n+1)!} |x-a|^{n+1}$$

for $|x - a| \le d$.

• Often we use the following fact:

$$\lim_{n \to \infty} \frac{x^n}{n!} = 0$$

for every real number x.

ullet Prove that e^x is equal to the sum of its Maclaurin series.

ullet Prove that e^x is equal to the sum of its Maclaurin series.

• Find the Taylor series for $f(x) = e^x$ at a = 1.

- ullet Prove that e^x is equal to the sum of its Maclaurin series.
- Find the Taylor series for $f(x) = e^x$ at a = 1.
- \bullet Find the Maclaurin series for $\sin x$ and prove that it represents $\sin x$ for all x.

- ullet Prove that e^x is equal to the sum of its Maclaurin series.
- Find the Taylor series for $f(x) = e^x$ at a = 1.
- \bullet Find the Maclaurin series for $\sin x$ and prove that it represents $\sin x$ for all x.

$$\sin x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}$$

ullet Find the Maclaurin series for $\cos x$.

Lecture 14

ullet Find the Maclaurin series for $\cos x$.

$$\cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$

 \bullet Find the Maclaurin series for $\cos x$.

$$\cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$

• Find the Maclaurin series for $x^2 \cos x$.

 \bullet Find the Maclaurin series for $\cos x$.

$$\cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}$$

- Find the Maclaurin series for $x^2 \cos x$.
- \bullet Evaluate $\int e^{-x^2} \mathrm{d}x$ as an infinite series.

ullet Use the Maclaurin series for e^x to evaluate

$$\lim_{x \to 0} \frac{e^x - 1 - x}{x^2}$$

ullet Use the Maclaurin series for e^x to evaluate

$$\lim_{x \to 0} \frac{e^x - 1 - x}{x^2}$$

• Find the sum of the series

$$\sum_{n=0}^{\infty} \frac{5^n}{3^n n!}$$