

Quantum mushrooms, scars, and high frequency cavity eigenmodes

UMass Amherst, CAM Seminar, 28 Nov, 2006

Alex Barnett

Department of Mathematics, Dartmouth College

Dirichlet eigenproblem

Normal modes of elastic membrane or ‘drum’ (Helmholtz, Germain, 19thC)

Eigenfunctions ϕ_j of Laplacian $\Delta := \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$ in bounded domain $\Omega \subset \mathbb{R}^2$

$$-\Delta \phi_j = E_j \phi_j \quad \phi_j|_{\partial\Omega} = 0 \text{ Dirichlet BC} \quad \int_{\Omega} \phi_i \phi_j \, d\mathbf{x} = \delta_{ij}$$

Dirichlet eigenproblem


Normal modes of elastic membrane or ‘drum’ (Helmholtz, Germain, 19thC)

Eigenfunctions ϕ_j of Laplacian $\Delta := \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$ in bounded domain $\Omega \subset \mathbb{R}^2$

$$-\Delta\phi_j = E_j\phi_j$$

$$\phi_j|_{\partial\Omega} = 0 \quad \text{Dirichlet BC}$$

$$\int_{\Omega} \phi_i \phi_j \, d\mathbf{x} = \delta_{ij}$$



mode $j = 1 \dots \infty$

discrete eigenvalues
 $E_1 < E_2 \leq E_3 \leq \dots \infty$

wavenumber $k_j = E_j^{1/2}$

wavelength $\frac{2\pi}{k_j}$

Dirichlet eigenproblem


Normal modes of elastic membrane or ‘drum’ (Helmholtz, Germain, 19thC)

Eigenfunctions ϕ_j of Laplacian $\Delta := \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$ in bounded domain $\Omega \subset \mathbb{R}^2$

$$-\Delta\phi_j = E_j\phi_j$$

$$\phi_j|_{\partial\Omega} = 0 \quad \text{Dirichlet BC}$$

$$\int_{\Omega} \phi_i \phi_j \, d\mathbf{x} = \delta_{ij}$$



mode $j = 1 \dots \infty$

discrete eigenvalues
 $E_1 < E_2 \leq E_3 \leq \dots \infty$


wavenumber $k_j = E_j^{1/2}$

wavelength $\frac{2\pi}{k_j}$

- What are asymptotic properties of modes in $E_j \rightarrow \infty$ limit ?


Some history of (related) normal modes

Ernst Chladni (1756–1827) sprinkles sand on metal plates
‘Plays’ them with violin bow: visualizes nodal lines ($\phi = 0$)



Some history of (related) normal modes

Ernst Chladni (1756–1827) sprinkles sand on metal plates
‘Plays’ them with violin bow: visualizes nodal lines ($\phi = 0$)



- popular lectures all over Europe
 - Napoleon impressed: offers 1 kg gold to explain patterns
 - Napoleon realised: irregularly shaped plate harder to understand!
 - Sophie Germain got prize in 1816



Note: rigid plate \neq membrane
(biharmonic Δ^2 rather than Laplacian Δ)

Modern applications

- electromagnetic waveguides (TM modes)
- acoustic resonances (Neumann BCs) *e.g.* aircraft cabin design
- paradigm for trickier resonance problems *e.g.* 3D RF/optical cavities


Modern applications

- electromagnetic waveguides (TM modes)
- acoustic resonances (Neumann BCs) *e.g.* aircraft cabin design
- paradigm for trickier resonance problems *e.g.* 3D RF/optical cavities
- single-particle quantum mechanics *e.g.* modeling quantum dots



quantum dots (Marcus)

resonant transmission, often single closed-dot mode responsible




transport statistics (Akis)


Dielectric micro-cavity laser design

leaky resonant cavities

quantum-cascade laser (Gmachl)



mode and emission pattern



- 2D cavity confinement due to total internal reflection ($n = 3.3$)
- approximated by Dirichlet eigenmodes; system size $10\text{--}50 \lambda$
- optimize design: need many modes for many shapes (Tureci '03)

Einstein-Brillouin-Keller (EBK)

(Keller '60)

Recall $\{\phi_j\}$ are products of 1D modes if Δ separable (ellipse, rectangle)


If not, can use short wavelength asymptotic mode approximation:

$$\phi(x, y) \approx \sum_{m=1}^M A_m(x, y) e^{ikS_m(x, y)}$$

mode \approx sum of traveling waves, ‘rays’

S_m = phase function

A_m = spatially-varying amplitude



Einstein-Brillouin-Keller (EBK)

(Keller '60)

Recall $\{\phi_j\}$ are products of 1D modes if Δ separable (ellipse, rectangle)


If not, can use short wavelength asymptotic mode approximation:

$$\phi(x, y) \approx \sum_{m=1}^M A_m(x, y) e^{ikS_m(x, y)}$$

mode \approx sum of traveling waves, ‘rays’

S_m = phase function

A_m = spatially-varying amplitude



Substitute into $-\Delta\phi = E\phi$ gives

$$|\nabla S_m| = 1 \quad \text{eikonal: phase grows along straight rays}$$

For such modes to exist:

- i) rays reflect off boundary, giving ray families which must close
- ii) quantization: round-trip phase = $2\pi n + \frac{\pi}{2}(\# \text{ focal points}) + \pi(\# \text{ reflections})$

Einstein-Brillouin-Keller (EBK)

(Keller '60)

Recall $\{\phi_j\}$ are products of 1D modes if Δ separable (ellipse, rectangle)


If not, can use short wavelength asymptotic mode approximation:

$$\phi(x, y) \approx \sum_{m=1}^M A_m(x, y) e^{ikS_m(x, y)}$$

mode \approx sum of traveling waves, ‘rays’

S_m = phase function

A_m = spatially-varying amplitude



Substitute into $-\Delta\phi = E\phi$ gives


$$|\nabla S_m| = 1 \quad \text{eikonal: phase grows along straight rays}$$

For such modes to exist:

- i) rays reflect off boundary, giving ray families which must close
- ii) quantization: round-trip phase = $2\pi n + \frac{\pi}{2}(\# \text{ focal points}) + \pi(\# \text{ reflections})$

But, do bouncing ray paths always form closed families...?

Bouncing rays: the game of billiards



phase space: $(x, y, \mathbf{p}_x, \mathbf{p}_y)$

$(x, y) = \mathbf{x}$ = position


$(\mathbf{p}_x, \mathbf{p}_y) = \mathbf{p}$ = momentum

trajectory $\mathbf{x}(t)$

free motion with reflection

Hamiltonian dynamical system: energy $H(\mathbf{x}, \mathbf{p}) = |\mathbf{p}|^2$ conserved

Bouncing rays: the game of billiards



phase space: (x, y, p_x, p_y)

$(x, y) = \mathbf{x}$ = position

$(p_x, p_y) = \mathbf{p}$ = momentum


trajectory $\mathbf{x}(t)$

free motion with reflection

Hamiltonian dynamical system: energy $H(\mathbf{x}, \mathbf{p}) = |\mathbf{p}|^2$ conserved


TWO BROAD CLASSES

integrable:



d conserved quantities ($d = 2$)
(tori \rightarrow closed ray families)

ergodic:



energy alone conserved: chaos!


Quantum chaos

Class of ray dynamics in Ω has profound effect on eigenmodes...


Quantum chaos

Class of ray dynamics in Ω has profound effect on eigenmodes...

Integrable:
 d conserved
quantities




Ergodic:
covers whole
energy shell




rays (classical)

eigenmodes ϕ_j (quantum)



regular modes: separable or EBK




irregular modes: ‘quantum chaos’


Quantum chaos

Class of ray dynamics in Ω has profound effect on eigenmodes...

Integrable:
 d conserved
quantities




Ergodic:
covers whole
energy shell




rays (classical)

eigenmodes ϕ_j (quantum)



regular modes: separable or EBK



irregular modes: ‘quantum chaos’

(phase space may contain both integrable and ergodic pieces, other subtle categories)

Mathematical questions

1. Quantum chaos: what happens in $E \rightarrow \infty$ semiclassical limit?
(Einstein 1917; Gutzwiller, Berry '80s; Zelditch...)


- impact: atomic, molecular, mesoscopic physics...
- more general Riemann surfaces: modular forms, number theory
- eigenvalue E_j statistics \leftrightarrow Random Matrix Theory (unproven!)

Mathematical questions

1. Quantum chaos: what happens in $E \rightarrow \infty$ semiclassical limit?
(Einstein 1917; Gutzwiller, Berry '80s; Zelditch...)

- impact: atomic, molecular, mesoscopic physics...
- more general Riemann surfaces: modular forms, number theory
- eigenvalue E_j statistics \leftrightarrow Random Matrix Theory (unproven!)

2. Spectral geometry: can one hear the shape of a drum? (Kac '66)



Answer: no !
(Gordon *et al.* '92)

Isospectral drums,
numerics to 14 digits
(Driscoll '97)

Engineering, physics, and mathematical questions all pose the following *numerical* challenges:

1. Complex geometry and cavity shapes, *e.g.* corners, 3D
2. Higher frequencies: **multiscale** problem
(wavelength \ll system size)

Outline of rest of talk

- I. Quantum ergodicity
- II. A new type of ‘scarring’ of eigenmodes
- III. The numerical methods that make it all possible

I. Quantum ergodicity

Quantum Ergodicity Theorem (QET)

If Ω ergodic, $A(\mathbf{x})$ ‘well-behaved’ function on Ω , with mean value
 $\overline{A} := \frac{1}{\text{vol}(\Omega)} \int_{\Omega} A(\mathbf{x}) d\mathbf{x}$, then

$$\lim_{E_j \rightarrow \infty} \langle \phi_j, A\phi_j \rangle - \overline{A} = 0 \quad \forall j \text{ except subseq. of vanishing density}$$

(Schnirelman '74, Colin de Verdière '85, Zelditch '87, Z-Zworski '96)


Quantum Ergodicity Theorem (QET)

If Ω ergodic, $A(\mathbf{x})$ ‘well-behaved’ function on Ω , with mean value
 $\overline{A} := \frac{1}{\text{vol}(\Omega)} \int_{\Omega} A(\mathbf{x}) d\mathbf{x}$, then

$$\lim_{E_j \rightarrow \infty} \langle \phi_j, A\phi_j \rangle - \overline{A} = 0 \quad \forall j \text{ except subseq. of vanishing density}$$

(Schnirelman '74, Colin de Verdière '85, Zelditch '87, Z-Zworski '96)

- prob. density $|\phi_j|^2 \xrightarrow{\text{weakly}} 1/\text{vol}(\Omega)$



$j = 1$

$j = 10$

$j = 10^2$

$j = 10^3$

$j \approx 5 \times 10^4$

Proof: Egorov's Thm (microlocal analysis, time propagation of wave eqn)


Quantum Ergodicity Theorem (QET)

If Ω ergodic, $A(\mathbf{x})$ ‘well-behaved’ function on Ω , with mean value
 $\overline{A} := \frac{1}{\text{vol}(\Omega)} \int_{\Omega} A(\mathbf{x}) d\mathbf{x}$, then

$$\lim_{E_j \rightarrow \infty} \langle \phi_j, A\phi_j \rangle - \overline{A} = 0 \quad \forall j \text{ except subseq. of vanishing density}$$

(Schnirelman '74, Colin de Verdière '85, Zelditch '87, Z-Zworski '96)

- prob. density $|\phi_j|^2 \xrightarrow{\text{weakly}} 1/\text{vol}(\Omega)$



$j = 1$

$j = 10$

$j = 10^2$

$j = 10^3$

$j \approx 5 \times 10^4$

Proof: Egorov's Thm (microlocal analysis, time propagation of wave eqn)

Of practical importance: At what *rate* is limit reached?

How fast does the density of excluded subsequence vanish?


Quantum Unique Ergodicity (QUE)

Conj. (Rudnick-Sarnak '94): There is *no* excluded subsequence in QET

Quantum Unique Ergodicity (QUE)

Conj. (Rudnick-Sarnak '94): There is *no* excluded subsequence in QET

- context: negatively curved manifolds (chaos comes from metric)




Recent rigorous analytic results:

- QUE proven for *arithmetic* manifold $SL_2(\mathbb{Z}) \backslash \mathbb{H}$ (Lindenstrauss '03)
- Proven *not* to hold for some quantum Arnold cat maps (Faure et al '03)

Quantum Unique Ergodicity (QUE)

Conj. (Rudnick-Sarnak '94): There is *no* excluded subsequence in QET

- context: negatively curved manifolds (chaos comes from metric)



Recent rigorous analytic results:


- QUE proven for *arithmetic* manifold $SL_2(\mathbb{Z}) \backslash \mathbb{H}$ (Lindenstrauss '03)
- Proven *not* to hold for some quantum Arnold cat maps (Faure et al '03)

QUE might be a surprise for chaotic cavity eigenmodes because of...

Scars: an exceptional subsequence?


High- j modes $|\phi_j|^2$ sometimes localize on unstable periodic orbits (UPO)

- discovered by *numerical* study of quarter-stadium modes (Heller '84)



- Wavepacket arguments \Rightarrow transverse width dies as $\sim E^{-1/4}$
- Note here also exceptional ‘bouncing ball’ sequence (neutrally stable)

High-eigenvalue scarred stadium mode



- ‘strong’ scar if there is a sequence $\{\phi_{j_n}\}$ with $O(1)$ mass on UPO
- QUE implies ‘weak’ scarring: mass on UPO vanishes as $E \rightarrow \infty$


Which occurs? No analytic results in 2D cavities...

Numerical experiments

(B, Comm. Pure Appl. Math. '06)

dispersing billiard, proven chaotic (Sinai '70)

$$\text{test function } A = \begin{cases} 1 & \text{in } \Omega_A, \\ 0 & \text{otherwise} \end{cases}$$




Numerical experiments

(B, Comm. Pure Appl. Math. '06)

dispersing billiard, proven chaotic (Sinai '70)

$$\text{test function } A = \begin{cases} 1 & \text{in } \Omega_A, \\ 0 & \text{otherwise} \end{cases}$$



Large-scale study, 30,000 modes in range $j \sim 10^4$ to 10^6 , enabled by:

1. Efficient boundary-based numerics for ϕ_j : ‘scaling method’
 2. matrix elements $\langle \phi_j, A\phi_j \rangle = \int_{\Omega_A} \phi_j^2 d\mathbf{x}$ via boundary integrals
- 100 times higher in j than any previous studies (e.g. Bäcker '98)
 - only a few CPU-days total


Typical high-frequency ergodic mode

225 λ across

quantum number
 $j \approx 5 \times 10^4$

$E_j \approx 10^6$

not very uniform,
rather, stringy...



(compare: random sum of plane waves)

$$\operatorname{Re} \sum_m a_m e^{i\mathbf{k}_m \cdot \mathbf{x}}$$

all wavenumbers


$$|\mathbf{k}_m| = \sqrt{E} = \text{const.}$$

also stringy...

interesting


to the eye only?

(a mystery)



Raw diagonal matrix element data

To reach high E , calculate $\langle \phi_j, A\phi_j \rangle$ only in certain eigenvalue intervals



- strong evidence for QUE (exceptional fraction $< 3 \times 10^{-5}$)

Rate of quantum ergodicity

E -dependent variance $V_A(E) :=$ local average of $|\langle \phi_j, A\phi_j \rangle - \bar{A}|^2$

At what rate does this ‘quantum variance’ die as $E \rightarrow \infty$?

Rate of quantum ergodicity

E -dependent variance $V_A(E) :=$ local average of $|\langle \phi_j, A\phi_j \rangle - \bar{A}|^2$

At what rate does this ‘quantum variance’ die as $E \rightarrow \infty$?

Conjecture (Feingold-Peres ’86): $V_A(E) \sim \frac{2\tilde{C}_A(0)}{\text{vol}(\Omega)} E^{-1/2}$


Rate of quantum ergodicity

E -dependent variance $V_A(E) :=$ local average of $|\langle \phi_j, A\phi_j \rangle - \bar{A}|^2$

At what rate does this ‘quantum variance’ die as $E \rightarrow \infty$?

Conjecture (Feingold-Peres ’86): $V_A(E) \sim \frac{2\tilde{C}_A(0)}{\text{vol}(\Omega)} E^{-1/2}$

- prefactor involves classical quantity $\tilde{C}_A(\omega) := \int_{-\infty}^{\infty} C_A(\tau) e^{i\omega\tau} d\tau$
- autocorrelation $C_A(\tau) := \frac{1}{2\pi\text{vol}(\Omega)} \int_{\Omega} \int_0^{2\pi} A(\mathbf{x}_0(0)) A(\mathbf{x}_0(\tau)) d\mathbf{x}_0 d\theta_0 - \bar{A}^2$



notation: $\mathbf{x}_0(t)$ is the unit-speed trajectory
launched at position \mathbf{x}_0 , angle θ_0


The integral is over the energy shell

Classical variance

$$\tilde{C}_A(\omega) := \int_{-\infty}^{\infty} C_A(\tau) e^{i\omega\tau} d\tau$$

FT of autocorrelation =
power spectral density
of noisy signal $A(\mathbf{x}(t))$
along ergodic trajectory

LISTEN to it!




Classical variance

$$\tilde{C}_A(\omega) := \int_{-\infty}^{\infty} C_A(\tau) e^{i\omega\tau} d\tau$$

FT of autocorrelation =
power spectral density
of noisy signal $A(\mathbf{x}(t))$
along ergodic trajectory

LISTEN to it!




Physically: $\tilde{C}_A(\omega)$ is heating rate due to periodic driving of classical system, at frequency ω , by external field A (Cohen '99, B '00)

- $\tilde{C}_A(\omega)$ gives band profile of off-diagonal elements $\langle \phi_i, A\phi_j \rangle$

FP Conj.: $\omega \rightarrow 0$ limit gives prefactor for quantum ergodicity rate

Numerical test: does FP Conj. hold in this ergodic cavity?

Results on ergodicity rate




consistent with
power law
 $V_A(E) = aE^{-\gamma}$
best fit $\gamma = 0.48 \pm 0.01$
(FP Conj. $\gamma = 1/2$)

- large numbers of modes → unprecedented accuracy ($< 1\%$)
- asymptotic regime seen for first time (but more data needed!)
- consistent with FP Conj., convergence very slow: 7% off at $j = 10^5$


II. New type of ‘moving scar’

Mushroom billiard




Divided phase space, free
from usual KAM hierarchy
(Bunimovich '01)

regular:




ergodic:



Eigenmodes have not been computed before now...

High freq. mushroom modes


(B-Betcke, sub. to Chaos)




$$\begin{aligned} k &= 100 \\ j &\approx 2000 \end{aligned}$$

- Percival's conjecture verified:
modes localize to either regular or chaotic region

Very high freq. mushroom modes




ergodic, equidistributed



ergodic, strongly scarred

Moving scar in boundary functions

$|\partial_n \phi_j(q)|^2$, boundary location q :




- sloping streak effect

MODES

SMOOTHED

Moving scar in boundary functions

$|\partial_n \phi_j(q)|^2$, boundary location q :




- sloping streak effect

MODES

SMOOTHED

take FT along k axis, gives:
wave autocorrelation in time



- refocused returning orbits
- return length varies with q

EVOLVE

III. Numerics (sketch)

Method of Particular Solutions

Given trial energy parameter $E > 0$:

- choose basis function set $\{\xi_i(E; \mathbf{x})\}_{i=1\dots N}$ with $-\Delta\xi_i = E\xi_i$ in Ω
global Helmholtz solutions with wrong BCs
then $u = \sum_{i=1}^N a_i \xi_i$ obeys $-\Delta u = Eu$ in Ω

Method of Particular Solutions

Given trial energy parameter $E > 0$:

- choose basis function set $\{\xi_i(E; \mathbf{x})\}_{i=1\dots N}$ with $-\Delta\xi_i = E\xi_i$ in Ω
global Helmholtz solutions with wrong BCs
then $u = \sum_{i=1}^N a_i \xi_i$ obeys $-\Delta u = Eu$ in Ω
- if can find coeff vector $\mathbf{a} \in \mathbb{R}^N$ giving $u|_{\partial\Omega} = 0$, but $u \neq 0$ in Ω
... then u is a mode ϕ_j and E is its eigenvalue E_j

Method of Particular Solutions

Given trial energy parameter $E > 0$:

- choose basis function set $\{\xi_i(E; \mathbf{x})\}_{i=1\dots N}$ with $-\Delta \xi_i = E \xi_i$ in Ω
global Helmholtz solutions with wrong BCs
then $u = \sum_{i=1}^N a_i \xi_i$ obeys $-\Delta u = Eu$ in Ω
- if can find coeff vector $\mathbf{a} \in \mathbb{R}^N$ giving $u|_{\partial\Omega} = 0$, but $u \neq 0$ in Ω
... then u is a mode ϕ_j and E is its eigenvalue E_j


Rayleigh quotient $t(E)^2 := \min_u \frac{\int_{\partial\Omega} u^2 ds}{\int_{\Omega} u^2 d\mathbf{x}} = \min_{\mathbf{a} \neq 0} \frac{\mathbf{a}^T F_E \mathbf{a}}{\mathbf{a}^T G_E \mathbf{a}} = \lambda_1(E)$

Method of Particular Solutions


Given trial energy parameter $E > 0$:

- choose basis function set $\{\xi_i(E; \mathbf{x})\}_{i=1\dots N}$ with $-\Delta \xi_i = E \xi_i$ in Ω
global Helmholtz solutions with wrong BCs
then $u = \sum_{i=1}^N a_i \xi_i$ obeys $-\Delta u = Eu$ in Ω
- if can find coeff vector $\mathbf{a} \in \mathbb{R}^N$ giving $u|_{\partial\Omega} = 0$, but $u \neq 0$ in Ω
... then u is a mode ϕ_j and E is its eigenvalue E_j

Rayleigh quotient $t(E)^2 := \min_u \frac{\int_{\partial\Omega} u^2 ds}{\int_{\Omega} u^2 d\mathbf{x}} = \min_{\mathbf{a} \neq 0} \frac{\mathbf{a}^T F_E \mathbf{a}}{\mathbf{a}^T G_E \mathbf{a}} = \lambda_1(E)$



$\Omega = \text{peanut}, j \approx 700$




- Each E_j found at a $t(E)$ minimum

Basis functions

Each basis func $\xi_i(E; \mathbf{x})$ is a *global* Helmholtz soln at energy E . . .

Basis functions

Each basis func $\xi_i(E; \mathbf{x})$ is a *global* Helmholtz soln at energy E . . .




$$\left\{ \begin{array}{ll} \text{Plane waves} & \sin(k\mathbf{n}_i \cdot \mathbf{x}), \\ \text{Fourier-Bessel} & J_l(kr) \sin(l\theta) \end{array} \right.$$

Thm: Ω smooth \Rightarrow exponential convergence (Eisenstat '74)

Practice: fail for nonconvex Ω (coeff sizes $|\mathbf{a}| \gg 10^{16}$)

Basis functions


Each basis func $\xi_i(E; \mathbf{x})$ is a *global* Helmholtz soln at energy E . . .



$$\begin{cases} \text{Plane waves} & \sin(k\mathbf{n}_i \cdot \mathbf{x}), \quad k^2 = E \\ \text{Fourier-Bessel} & J_l(kr) \sin(l\theta) \end{cases}$$

Thm: Ω smooth \Rightarrow exponential convergence (Eisenstat '74)

Practice: fail for nonconvex Ω (coeff sizes $|\mathbf{a}| \gg 10^{16}$)



Fundamental solutions


$Y_0(k|\mathbf{x} - \mathbf{x}_i|)$ with $\{\mathbf{x}_i\}$ on outer boundary

Practice: excellent, also w/ non-reentrant corners (B '06)

Ongoing analysis of coeff sizes (w/ Timo Betcke)

Basis functions

Each basis func $\xi_i(E; \mathbf{x})$ is a *global* Helmholtz soln at energy E . . .



$$\begin{cases} \text{Plane waves} & \sin(k\mathbf{n}_i \cdot \mathbf{x}), \quad k^2 = E \\ \text{Fourier-Bessel} & J_l(kr) \sin(l\theta) \end{cases}$$

Thm: Ω smooth \Rightarrow exponential convergence (Eisenstat '74)

Practice: fail for nonconvex Ω (coeff sizes $|\mathbf{a}| \gg 10^{16}$)

Fundamental solutions

$Y_0(k|\mathbf{x} - \mathbf{x}_i|)$ with $\{\mathbf{x}_i\}$ on outer boundary


Practice: excellent, also w/ non-reentrant corners (B '06)

Ongoing analysis of coeff sizes (w/ Timo Betcke)

Corner-adapted Fourier-Bessel:

singular corners $\theta \neq \frac{\pi}{n}$

Practice: exp. conv. for multiple corners (Betcke '05)



Scaling method


Recall $F_E \mathbf{a} = \lambda G_E \mathbf{a}$ F_E, G_E basis reps. of Rayleigh quotient $\|u\|_{L^2(\partial\Omega)}^2 / \|u\|_{L^2(\Omega)}^2$
Minimizing $\lambda_1(E)$ slow; nearby minima easily missed—can do better?

Scaling method

Recall $F_E \mathbf{a} = \lambda G_E \mathbf{a}$ F_E, G_E basis reps. of Rayleigh quotient $\|u\|_{L^2(\partial\Omega)}^2 / \|u\|_{L^2(\Omega)}^2$

Minimizing $\lambda_1(E)$ slow; nearby minima easily missed—can do better?

Include higher generalized eigenvalues...



- Clue: spectrum at single E has info about many nearby $\hat{\lambda}_1$ minima

Scaling method

(Vergini-Saraceno '94; B '00, '06)


New Rayleigh quotient: $\int_{\partial\Omega} (\mathbf{x} \cdot \mathbf{n})^{-1} |u|^2 ds / \int_{\partial\Omega} (\mathbf{x} \cdot \mathbf{n})^{-1} u \mathbf{x} \cdot \nabla u ds$

Scaling method

(Vergini-Saraceno '94; B '00, '06)

New Rayleigh quotient: $\int_{\partial\Omega} (\mathbf{x} \cdot \mathbf{n})^{-1} |u|^2 ds / \int_{\partial\Omega} (\mathbf{x} \cdot \mathbf{n})^{-1} u \mathbf{x} \cdot \nabla u ds$

Generalized eigenvalues $\lambda_l(E)$ linear in $E - E_j$ (for Ω star-shaped):



- solving $F_E \mathbf{a} = \lambda G_E \mathbf{a}$ at **single** E value gives *all* nearest $O(k)$ modes
- no root search, no missing levels, speed gain $O(k) \sim 10^3$ over MPS
- eigenvectors \mathbf{a}_l give *dilated* (rescaled) approximations to modes ϕ_j
- errors grow like $t \sim |E_j - E|^3$ (3rd-order convergence with effort)

Bifurcated genealogy

NUMERICAL ANALYSIS: high accuracy

Vekua ('60s) MPS (Fox–Henrici–Moler '67)

Complex approximation theory
Schryer, Eisenstat ('70s)
Still ('80s)

Trefethen

Driscoll

GSVD (Betcke–Trefethen '05)

Fundamental solns (MFS)
(Karageorghis '00)


QUANTUM PHYSICS: high freq

Plane wave method
(Heller '80s)
Scaling method
(Vergini–Saraceno '94)

Barnett ('00, '06)

GSVD (Betcke–Trefethen '05)  Barnett ('00, '06)

Bifurcated genealogy



Researchers using my code: M. Porter (Caltech), P. Doyle (Dartmouth), M. Wright (ISVR, UK), J. Belmonte (UCLM, Spain), H. Mosadegh (Isfahan, Iran), Vinayak (Nehru U., India) ...

Conclusion

Dirichlet eigenproblem: a paradigm linear wave resonance problem

Verified numerically in high-frequency regime . . .

- asymptotic decay $E^{-1/4}$ of fluctuations in chaotic modes
- found in mushroom: new moving scar fringe effect at focal points

Numerics: global basis methods excel at high frequency $E \gg 1$:

- scaling method $O(E^{1/2})$ (typ. 10^3) faster than any known method
- working on: basis choice; errors; dielectric resonator problem . . .

Thanks: T. Betcke (Manchester)

P. Sarnak (Princeton)

L. N. Trefethen (Oxford)

S. Zelditch (JHU)

P. Deift (NYU)

Funding: NSF (DMS-0507614)

Preprints, talks, movies:

<http://math.dartmouth.edu/~ahb>


Made with: Linux, L^AT_EX, Prosper

Laser modes

In reality the laser modes leak out into free space (they radiate)
Let's approximate by close cavity (Dirichlet BC):


$E = 16417.2$

MODES




CPU / mode	method
60 s	MPS, root search
10 s	scaling, ϕ_j across Ω
0.6 s	scaling, ϕ_j bdry only

Bizarre laser shapes: spiral cavity



micrograph (Kneissl *et al.* '04)



numerics (Chern *et al.* '03)

- what is optimal shape/pumping?
- hard numerical calculations

Random plane waves

Conjecture (Berry '77): statistical model of eigenmodes

$$\phi_n \sim \lim_{N \rightarrow \infty} \frac{1}{\sqrt{N}} \sum_{j=1}^N a_j \sin(\mathbf{k}_j \cdot \mathbf{r} + \alpha_j)$$

iid amplitudes $a_j \in \mathbb{R}$
iid phases $\alpha_j \in [0, 2\pi)$

Wavevectors \mathbf{k}_j , spaced uniformly in direction, $|\mathbf{k}_j| = k$.

- Ray analogue of classical ergodicity.
- uniform measure as $E = k^2 \rightarrow \infty$