

A new integral representation of quasi-periodic fields and its application to scattering and Bloch eigenvalue problems

FACM, May 22, 2010

Alex Barnett (Dartmouth College)

joint work with Leslie Greengard (Courant Institute, NYU)




Scattering in 2D from periodic grating

time-harmonic linear waves, obey $(\Delta + \omega^2)u = 0$

Helmholtz, freq ω

incident plane wave $u^i(\mathbf{x}) = e^{i\mathbf{k}\cdot\mathbf{x}}$ wavevector $\mathbf{k} = (\kappa^i, k^i)$ $|\mathbf{k}| = \omega$, unit speed


$\Omega \subset \mathbb{R}^2$ obstacle, $\Omega_{\mathbb{Z}} = \{\mathbf{x} : (x + nd, y) \in \Omega \text{ for some } n \in \mathbb{Z}\}$



Scattering in 2D from periodic grating

time-harmonic linear waves, obey $(\Delta + \omega^2)u = 0$ Helmholtz, freq ω
incident plane wave $u^i(\mathbf{x}) = e^{i\mathbf{k}\cdot\mathbf{x}}$ wavevector $\mathbf{k} = (\kappa^i, k^i)$ $|\mathbf{k}| = \omega$, unit speed

$\Omega \subset \mathbb{R}^2$ obstacle, $\Omega_{\mathbb{Z}} = \{\mathbf{x} : (x + nd, y) \in \Omega \text{ for some } n \in \mathbb{Z}\}$



total field $u^t = u^i + u$, where

scattered field u solves BVP:

$$(\Delta + \omega^2)u = 0 \quad \text{in } \mathbb{R}^2 \setminus \overline{\Omega_{\mathbb{Z}}}$$


$$u = -u^i \quad \text{on } \partial\Omega_{\mathbb{Z}} \quad \text{Dirichlet}$$

u ‘radiative’ as $y \rightarrow \pm\infty$

Scattering in 2D from periodic grating

time-harmonic linear waves, obey $(\Delta + \omega^2)u = 0$ Helmholtz, freq ω
incident plane wave $u^i(\mathbf{x}) = e^{i\mathbf{k}\cdot\mathbf{x}}$ wavevector $\mathbf{k} = (\kappa^i, k^i)$ $|\mathbf{k}| = \omega$, unit speed

$\Omega \subset \mathbb{R}^2$ obstacle, $\Omega_{\mathbb{Z}} = \{\mathbf{x} : (x + nd, y) \in \Omega \text{ for some } n \in \mathbb{Z}\}$



total field $u^t = u^i + u$, where

scattered field u solves BVP:

$$(\Delta + \omega^2)u = 0 \quad \text{in } \mathbb{R}^2 \setminus \overline{\Omega_{\mathbb{Z}}}$$

$$u = -u^i \quad \text{on } \partial\Omega_{\mathbb{Z}} \quad \text{Dirichlet}$$


u ‘radiative’ as $y \rightarrow \pm\infty$

u^i ‘quasi-periodic’, so u is too: $u(x + d, y) = \alpha u(x, y)$ Bloch $\alpha = e^{i\kappa^i d}$

Scattering in 2D from periodic grating

time-harmonic linear waves, obey $(\Delta + \omega^2)u = 0$ Helmholtz, freq ω
incident plane wave $u^i(\mathbf{x}) = e^{i\mathbf{k}\cdot\mathbf{x}}$ wavevector $\mathbf{k} = (\kappa^i, k^i)$ $|\mathbf{k}| = \omega$, unit speed

$\Omega \subset \mathbb{R}^2$ obstacle, $\Omega_{\mathbb{Z}} = \{\mathbf{x} : (x + nd, y) \in \Omega \text{ for some } n \in \mathbb{Z}\}$



total field $u^t = u^i + u$, where

scattered field u solves BVP:

$$(\Delta + \omega^2)u = 0 \quad \text{in } \mathbb{R}^2 \setminus \overline{\Omega_{\mathbb{Z}}}$$

$$u = -u^i \quad \text{on } \partial\Omega_{\mathbb{Z}} \quad \text{Dirichlet}$$

u ‘radiative’ as $y \rightarrow \pm\infty$

u^i ‘quasi-periodic’, so u is too: $u(x + d, y) = \alpha u(x, y)$ Bloch $\alpha = e^{i\kappa^i d}$

- classical BVP: acoustics, z -invariant Maxwell (Rayleigh 1897,...)
- Apps: optics, filters, lithography, photonics, plasmons, surface meas...

Plane waves of same quasi-periodicity


u^i is QP, but so are other plane waves, wavevectors (κ_n, k_n) :

$$\kappa_n = \kappa^i + 2\pi n/d \quad k_n = +\sqrt{\omega^2 - \kappa_n^2} \quad \text{positive real or positive imag}$$

Plane waves of same quasi-periodicity

u^i is QP, but so are other plane waves, wavevectors (κ_n, k_n) :


$$\kappa_n = \kappa^i + 2\pi n/d \quad k_n = +\sqrt{\omega^2 - \kappa_n^2} \quad \text{positive real or positive imag}$$



Plane waves of same quasi-periodicity

u^i is QP, but so are other plane waves, wavevectors (κ_n, k_n) :

$$\kappa_n = \kappa^i + 2\pi n/d \quad k_n = +\sqrt{\omega^2 - \kappa_n^2} \quad \text{positive real or positive imag}$$



k_n real: propagating


k_n imag: evanescent (decaying or growing in y)

$k_n = 0$: so-called Wood's anomaly rapid change wrt ω or inc. angle θ^i

Plane waves of same quasi-periodicity

u^i is QP, but so are other plane waves, wavevectors (κ_n, k_n) :

$$\kappa_n = \kappa^i + 2\pi n/d \quad k_n = +\sqrt{\omega^2 - \kappa_n^2} \quad \text{positive real or positive imag}$$




k_n real: propagating

k_n imag: evanescent (decaying or growing in y)

$k_n = 0$: so-called Wood's anomaly rapid change wrt ω or inc. angle θ^i

Rayleigh–Bloch radiation conditions



scattered u only outgoing or decaying channel modes:


$y > y_0$: upwards-prop. (or -decay)

$$u(x, y) = \sum_{n \in \mathbb{Z}} c_n e^{i\kappa_n x} e^{ik_n(y-y_0)}$$

$y < -y_0$: downwards-prop. (or -decay)

$$u(x, y) = \sum_{n \in \mathbb{Z}} d_n e^{i\kappa_n x} e^{ik_n(-y-y_0)}$$

Rayleigh–Bloch radiation conditions



- c_n, d_n scattered amplitudes

scattered u only outgoing or decaying channel modes:

$y > y_0$: upwards-prop. (or -decay)


$$u(x, y) = \sum_{n \in \mathbb{Z}} c_n e^{i\kappa_n x} e^{ik_n(y-y_0)}$$

$y < -y_0$: downwards-prop. (or -decay)

$$u(x, y) = \sum_{n \in \mathbb{Z}} d_n e^{i\kappa_n x} e^{ik_n(-y-y_0)}$$

vertical channel/waveguide with QP BC

Rayleigh–Bloch radiation conditions



scattered u only outgoing or decaying channel modes:

$y > y_0$: upwards-prop. (or -decay)

$$u(x, y) = \sum_{n \in \mathbb{Z}} c_n e^{i\kappa_n x} e^{ik_n(y-y_0)}$$

$y < -y_0$: downwards-prop. (or -decay)

$$u(x, y) = \sum_{n \in \mathbb{Z}} d_n e^{i\kappa_n x} e^{ik_n(-y-y_0)}$$

- c_n, d_n scattered amplitudes

vertical channel/waveguide with QP BC

Numerical PDE methods \exists many: finite difference, finite elements...

Formulate BVP as integral equation on $\partial\Omega$ to solve numerically for u

adv: efficient rep (small # unknowns), rad. cond. for free, high-order accurate

Potential theory (review)

‘charge’ (source of waves) distributed along curve Γ w/ density func.


single-, double-layer potentials, $\mathbf{x} \in \mathbb{R}^2$:

$$v(\mathbf{x}) = \int_{\Gamma} \Phi_{\omega}(\mathbf{x}, \mathbf{y}) \sigma(\mathbf{y}) ds_{\mathbf{y}} := (\mathcal{S}\sigma)(\mathbf{x})$$

$$u(\mathbf{x}) = \int_{\Gamma} \frac{\partial \Phi_{\omega}}{\partial n_y}(\mathbf{x}, \mathbf{y}) \tau(\mathbf{y}) ds_{\mathbf{y}} := (\mathcal{D}\tau)(\mathbf{x})$$

$$\Phi_{\omega}(\mathbf{x}, \mathbf{y}) := \Phi_{\omega}(\mathbf{x} - \mathbf{y}) = \frac{i}{4} H_0^{(1)}(\omega |\mathbf{x} - \mathbf{y}|)$$

Helmholtz fundamental soln
a.k.a. free space Greens func



Potential theory (review)

‘charge’ (source of waves) distributed along curve Γ w/ density func.


single-, double-layer potentials, $\mathbf{x} \in \mathbb{R}^2$:

$$v(\mathbf{x}) = \int_{\Gamma} \Phi_{\omega}(\mathbf{x}, \mathbf{y}) \sigma(\mathbf{y}) ds_{\mathbf{y}} := (\mathcal{S}\sigma)(\mathbf{x})$$

$$u(\mathbf{x}) = \int_{\Gamma} \frac{\partial \Phi_{\omega}}{\partial n_y}(\mathbf{x}, \mathbf{y}) \tau(\mathbf{y}) ds_{\mathbf{y}} := (\mathcal{D}\tau)(\mathbf{x})$$

$$\Phi_{\omega}(\mathbf{x}, \mathbf{y}) := \Phi_{\omega}(\mathbf{x} - \mathbf{y}) = \frac{i}{4} H_0^{(1)}(\omega |\mathbf{x} - \mathbf{y}|)$$

Helmholtz fundamental soln
a.k.a. free space Greens func



Jump relations: limit as $\mathbf{x} \rightarrow \Gamma$ may depend on which side (\pm):

$$v^{\pm} = S\sigma$$

$$v_n^{\pm} = D^T \sigma \mp \frac{1}{2} \sigma$$

$$u^{\pm} = D\tau \pm \frac{1}{2}\tau$$

$$u_n^{\pm} = T\tau$$


S, D are integral ops with above kernels
but defined on $C(\Gamma) \rightarrow C(\Gamma)$

T has kernel $\frac{\partial^2 \Phi_{\omega}(\mathbf{x}, \mathbf{y})}{\partial n_x \partial n_y}$

Integral equations for scattering (sketch)


single Dirichlet obstacle: represent scattered $u = \mathcal{D}\tau$

DLP on $\partial\Omega$



Integral equations for scattering (sketch)

single Dirichlet obstacle: represent scattered $u = \mathcal{D}\tau$ DLP on $\partial\Omega$




BC & JR3: $A\tau := (\frac{1}{2}I + D)\tau = -u^i$

2nd-kind IE on $\partial\Omega$, D compact so A sing. vals. $\not\rightarrow 0$

- why important? large scale problems...
condition # small, iterative solvers (GMRES) fast

Integral equations for scattering (sketch)

single Dirichlet obstacle: represent scattered $u = \mathcal{D}\tau$ DLP on $\partial\Omega$



BC & JR3: $A\tau := (\frac{1}{2}I + D)\tau = -u^i$

2nd-kind IE on $\partial\Omega$, D compact so A sing. vals. $\not\rightarrow 0$

- why important? large scale problems...
condition # small, iterative solvers (GMRES) fast


Quadrature scheme: nodes $\mathbf{y}_j \in \partial\Omega$, weights w_j , $j = 1, \dots, N$

Nyström discretization: N -by- N linear system for vector $\{\tau_j^{(N)}\}_{j=1}^N$

$$\frac{1}{2}\tau_k^{(N)} + \sum_{j=1}^N w_j D(\mathbf{y}_k, \mathbf{y}_j) \tau_j^{(N)} = -u^i(\mathbf{y}_k), \quad k = 1, \dots, N$$

Integral equations for scattering (sketch)

single Dirichlet obstacle: represent scattered $u = \mathcal{D}\tau$ DLP on $\partial\Omega$



BC & JR3: $A\tau := (\frac{1}{2}I + D)\tau = -u^i$

2nd-kind IE on $\partial\Omega$, D compact so A sing. vals. $\not\rightarrow 0$

- why important? large scale problems...
condition # small, iterative solvers (GMRES) fast

Quadrature scheme: nodes $\mathbf{y}_j \in \partial\Omega$, weights w_j , $j = 1, \dots, N$

Nyström discretization: N -by- N linear system for vector $\{\tau_j^{(N)}\}_{j=1}^N$


$$\frac{1}{2}\tau_k^{(N)} + \sum_{j=1}^N w_j D(\mathbf{y}_k, \mathbf{y}_j) \tau_j^{(N)} = -u^i(\mathbf{y}_k), \quad k = 1, \dots, N$$

Thm: (Anselone, Kress) $\|\tau^{(N)} - \tau\|_\infty$ converges at *same rate* as the quadrature scheme for true integrand $D(\mathbf{y}, \cdot)\tau$

e.g. here exponential: $N = 60$ enough for 10^{-6} error, $N = 100$ for 10^{-12}

Integral equations for scattering (sketch)

single Dirichlet obstacle: represent scattered $u = \mathcal{D}\tau$ DLP on $\partial\Omega$



BC & JR3: $A\tau := (\frac{1}{2}I + D)\tau = -u^i$

2nd-kind IE on $\partial\Omega$, D compact so A sing. vals. $\not\rightarrow 0$

- why important? large scale problems...
condition # small, iterative solvers (GMRES) fast

Quadrature scheme: nodes $\mathbf{y}_j \in \partial\Omega$, weights w_j , $j = 1, \dots, N$

Nyström discretization: N -by- N linear system for vector $\{\tau_j^{(N)}\}_{j=1}^N$

$$\frac{1}{2}\tau_k^{(N)} + \sum_{j=1}^N w_j D(\mathbf{y}_k, \mathbf{y}_j) \tau_j^{(N)} = -u^i(\mathbf{y}_k), \quad k = 1, \dots, N$$

Thm: (Anselone, Kress) $\|\tau^{(N)} - \tau\|_\infty$ converges at *same rate* as
the quadrature scheme for true integrand $D(\mathbf{y}, \cdot)\tau$

e.g. here exponential: $N = 60$ enough for 10^{-6} error, $N = 100$ for 10^{-12}

- efficient (2D became 1D), \exists fast $O(N \log N)$ schemes for $N > 10^4$
- in practice for unique solution $\forall \omega$: replace \mathcal{D} by $\mathcal{D} - i\omega \mathcal{S}$

The standard way to periodize

replace kernel $\Phi_\omega(\mathbf{x})$ by $\Phi_{\omega,\text{QP}}(\mathbf{x}) := \sum_{m \in \mathbb{Z}} \alpha^m \Phi_\omega(\mathbf{x} - m\mathbf{d})$

thus integral operator A becomes A_{QP} (is still 2nd kind)

The standard way to periodize

replace kernel $\Phi_\omega(\mathbf{x})$ by $\Phi_{\omega,\text{QP}}(\mathbf{x}) := \sum_{m \in \mathbb{Z}} \alpha^m \Phi_\omega(\mathbf{x} - m\mathbf{d})$

thus integral operator A becomes A_{QP} (is still 2nd kind)

Three problems

- not robust: $\Phi_{\omega,\text{QP}}$ does **not exist** for Wood's anomaly params (ω, θ^i)
blows up $\sim (\omega - \omega_{\text{Wood}})^{-1/2}$, round-off error too; but soln u well-behaved!
one fix (for surfaces): use half-space $\Phi_{\omega,\text{QP}}$ (Chandler-Wilde, Arens, ...)

The standard way to periodize

replace kernel $\Phi_\omega(\mathbf{x})$ by $\Phi_{\omega,\text{QP}}(\mathbf{x}) := \sum_{m \in \mathbb{Z}} \alpha^m \Phi_\omega(\mathbf{x} - m\mathbf{d})$

thus integral operator A becomes A_{QP} (is still 2nd kind)

Three problems

- not robust: $\Phi_{\omega,\text{QP}}$ does **not exist** for Wood's anomaly params (ω, θ^i)
blows up $\sim (\omega - \omega_{\text{Wood}})^{-1/2}$, round-off error too; but soln u well-behaved!
one fix (for surfaces): use half-space $\Phi_{\omega,\text{QP}}$ (Chandler-Wilde, Arens,...)
- $\Phi_{\omega,\text{QP}}(\mathbf{x})$ hard to evaluate accurately
 - e.g. 50 evals per sec (Kurkcu–Reitich '09)
 - e.g. by lattice sums: Fourier-Bessel coeffs s_l of regular part,

$$\Phi_{\omega,\text{QP}}(\mathbf{x}) - \Phi_\omega(\mathbf{x}) = \sum_{l \in \mathbb{Z}} s_l J_l(\omega r) e^{il\theta}, \quad \mathbf{x} = (r, \theta) \quad (*)$$

slow; messy in 3D (McPhedran, Linton, Moroz,...)

The standard way to periodize

replace kernel $\Phi_\omega(\mathbf{x})$ by $\Phi_{\omega,\text{QP}}(\mathbf{x}) := \sum_{m \in \mathbb{Z}} \alpha^m \Phi_\omega(\mathbf{x} - m\mathbf{d})$

thus integral operator A becomes A_{QP} (is still 2nd kind)


Three problems

- not robust: $\Phi_{\omega,\text{QP}}$ does **not exist** for Wood's anomaly params (ω, θ^i)
blows up $\sim (\omega - \omega_{\text{Wood}})^{-1/2}$, round-off error too; but soln u well-behaved!
one fix (for surfaces): use half-space $\Phi_{\omega,\text{QP}}$ (Chandler-Wilde, Arens,...)
- $\Phi_{\omega,\text{QP}}(\mathbf{x})$ hard to evaluate accurately
e.g. 50 evals per sec (Kurkcu–Reitich '09)
e.g. by lattice sums: Fourier-Bessel coeffs s_l of regular part,

$$\Phi_{\omega,\text{QP}}(\mathbf{x}) - \Phi_\omega(\mathbf{x}) = \sum_{l \in \mathbb{Z}} s_l J_l(\omega r) e^{il\theta}, \quad \mathbf{x} = (r, \theta) \quad (*)$$

slow; messy in 3D (McPhedran, Linton, Moroz,...)

- $(*)$ converges in disc \Rightarrow high aspect ratio Ω is bad:
expand disc? sum near neighbors directly, needs large #




New way to periodize

use only free-space Φ_ω , add densities on unit cell walls, enforce QP
fixes 3 problems: avoids blow-up, no lattice sums, no aspect ratio limit

New way to periodize


use only free-space Φ_ω , add densities on unit cell walls, enforce QP
fixes 3 problems: avoids blow-up, no lattice sums, no aspect ratio limit



$$\begin{aligned} u &= \mathcal{D}\tau + u_{\text{QP}}[\xi] \\ &\quad \text{as before} \qquad \text{densities } \xi \text{ on } L \text{ and } R \\ \text{BC} \quad u &= -u^i \quad \text{on } \partial\Omega \qquad \text{as before} \end{aligned}$$

New way to periodize

use only free-space Φ_ω , add densities on unit cell walls, enforce QP
fixes 3 problems: avoids blow-up, no lattice sums, no aspect ratio limit



$$\begin{aligned} u &= \mathcal{D}\tau + u_{\text{QP}}[\xi] \\ &\quad \text{as before} \qquad \text{densities } \xi \text{ on } L \text{ and } R \\ \text{BC} \quad u &= -u^i \quad \text{on } \partial\Omega \quad \text{as before} \end{aligned}$$


new condition: vanishing ‘discrepancy’

$$\forall y \quad \begin{cases} f := u_L - \alpha^{-1}u_R = 0 \\ f' := u_{nL} - \alpha^{-1}u_{nR} = 0 \end{cases}$$

2 unknowns $[\tau; \xi]$, 2 conditions \Rightarrow solve 2×2 linear operator system

New way to periodize

use only free-space Φ_ω , add densities on unit cell walls, enforce QP
 fixes 3 problems: avoids blow-up, no lattice sums, no aspect ratio limit



$$u = \mathcal{D}\tau + u_{\text{QP}}[\xi]$$

as before densities ξ on L and R

BC $u = -u^i$ on $\partial\Omega$ as before

new condition: vanishing ‘discrepancy’

$$\forall y \quad \begin{cases} f := u_L - \alpha^{-1}u_R = 0 \\ f' := u_{nL} - \alpha^{-1}u_{nR} = 0 \end{cases}$$


2 unknowns $[\tau; \xi]$, 2 conditions \Rightarrow solve 2×2 linear operator system

Major issues

- (1) How choose rep $u_{\text{QP}}[\xi]$ so effect of ξ on $[f; f']$ is ‘nice’ ? (2nd-kind)
- (2) How handle densities on ∞ -long L, R ? (no decay as $|y| \rightarrow \infty$)

Trick (1): choose a good $u_{QP}[\xi]$ representation


Consider ξ
one SLP:



effect on discrep: $f = (S_{LL} - \alpha^{-1}S_{RL})\sigma$
self-interaction, bad ↗


Trick (1): choose a good $u_{QP}[\xi]$ representation

Consider ξ
one SLP:



effect on discrep: $f = (S_{LL} - \alpha^{-1}S_{RL})\sigma$
self-interaction, bad ↗


Add phased
copy on R :



$$f = (S_{LL} - \alpha^{-1}S_{RL})\sigma + \alpha(S_{LR} - \alpha^{-1}S_{RR})\sigma \\ = (-\alpha^{-1}S_{RL} + \alpha S_{LR})\sigma \quad \text{distant only}$$


Trick (1): choose a good $u_{QP}[\xi]$ representation

Consider ξ
one SLP:



effect on discrep: $f = (S_{LL} - \alpha^{-1}S_{RL})\sigma$
self-interaction, bad ↗


Add phased
copy on R :



$$\begin{aligned} f &= (S_{LL} - \alpha^{-1}S_{RL})\sigma + \alpha(S_{LR} - \alpha^{-1}S_{RR})\sigma \\ &= (-\alpha^{-1}S_{RL} + \alpha S_{LR})\sigma \quad \text{distant only} \\ f' &= (-I - \alpha^{-1}D_{RL}^* + \alpha D_{LR}^*)\sigma \quad I/2's \text{ add} \end{aligned}$$


Trick (1): choose a good $u_{QP}[\xi]$ representation

Consider ξ
one SLP:



effect on discrep: $f = (S_{LL} - \alpha^{-1}S_{RL})\sigma$
self-interaction, bad ↗


Add phased
copy on R :



$$\begin{aligned} f &= (S_{LL} - \alpha^{-1}S_{RL})\sigma + \alpha(S_{LR} - \alpha^{-1}S_{RR})\sigma \\ &= (-\alpha^{-1}S_{RL} + \alpha S_{LR})\sigma \quad \text{distant only} \\ f' &= (-I - \alpha^{-1}D_{RL}^* + \alpha D_{LR}^*)\sigma \quad I/2's \text{ add} \end{aligned}$$

Similarly need to control f via JRs, so...

Add DLP τ
on L, R :



$$\begin{bmatrix} f \\ f' \end{bmatrix} = Q \begin{bmatrix} \tau \\ -\sigma \end{bmatrix} =: Q\xi$$

block operator $Q = I + (\text{interactions of distance } \geq d)$

- If L, R bounded segments: $Q\xi = g$ is 2nd kind, rapidly convergent



Trick (2): handle densities on $y \in (-\infty, \infty)$

Fourier transform in y -direction: work with $\hat{\sigma}, \hat{\tau}, \hat{f}, \hat{f}'$

Trick (2): handle densities on $y \in (-\infty, \infty)$

Fourier transform in y -direction: work with $\hat{\sigma}, \hat{\tau}, \hat{f}, \hat{f}'$, use spectral rep,


$$\Phi_\omega(x, y) = \frac{i}{4\pi} \int_{-\infty}^{\infty} e^{iky} \frac{e^{i\sqrt{\omega^2 - k^2}|x|}}{\sqrt{\omega^2 - k^2}} dk \quad \begin{matrix} \text{exponential tails for } |k| > \omega \\ \text{decay rate prop. to } |x| \end{matrix}$$



Trick (2): handle densities on $y \in (-\infty, \infty)$

Fourier transform in y -direction: work with $\hat{\sigma}, \hat{\tau}, \hat{f}, \hat{f}'$, use spectral rep,

$$\Phi_\omega(x, y) = \frac{i}{4\pi} \int_{-\infty}^{\infty} e^{iky} \frac{e^{i\sqrt{\omega^2 - k^2}|x|}}{\sqrt{\omega^2 - k^2}} dk \quad \begin{matrix} \text{exponential tails for } |k| > \omega \\ \text{decay rate prop. to } |x| \end{matrix}$$




Use FT- y densities on L (or R) wall at x_0 :

$$(\hat{\mathcal{S}}_L \hat{\sigma})(x, y) = \frac{i}{2} \int_{-\infty}^{\infty} e^{iky} \frac{e^{i\sqrt{\omega^2 - k^2}|x-x_0|}}{\sqrt{\omega^2 - k^2}} \hat{\sigma}(k) dk$$

$$(\hat{\mathcal{D}}_L \hat{\tau})(x, y) = \frac{\text{sign}(x - x_0)}{2} \int_{-\infty}^{\infty} e^{iky} e^{i\sqrt{\omega^2 - k^2}|x-x_0|} \hat{\tau}(k) dk$$

- same JRs as before; $\hat{\sigma}(k), \hat{\tau}(k)$ affect only $\hat{f}(k), \hat{f}'(k)$ diagonal in k

Quadrature on Sommerfeld contour




nodes k_j
weights w_j
 $j = 1, \dots, M$

$\text{Re } k$ is periodic trapezoid rule, $\text{Im } k$ is scaled tanh curve

- exponentially convergent as $h \rightarrow 0, K \rightarrow \infty$ beats nodes on real axis

Quadrature on Sommerfeld contour




nodes k_j
weights w_j
 $j = 1, \dots, M$

$\text{Re } k$ is periodic trapezoid rule, $\text{Im } k$ is scaled tanh curve

- exponentially convergent as $h \rightarrow 0, K \rightarrow \infty$ beats nodes on real axis

Unknowns $\hat{\sigma}(k_j), \hat{\tau}(k_j)$; enforce $\hat{f}(k_j) = \hat{f}'(k_j) = 0$: $\hat{Q} = I + \text{block diag.}$

Quadrature on Sommerfeld contour



nodes k_j
weights w_j
 $j = 1, \dots, M$

$\text{Re } k$ is periodic trapezoid rule, $\text{Im } k$ is scaled tanh curve

- exponentially convergent as $h \rightarrow 0, K \rightarrow \infty$ beats nodes on real axis

Unknowns $\hat{\sigma}(k_j), \hat{\tau}(k_j)$; enforce $\hat{f}(k_j) = \hat{f}'(k_j) = 0$: $\hat{Q} = I + \text{block diag.}$


Solve full $(N+2M)$ -by- $(N+2M)$ linear system:

$$\begin{bmatrix} A & B \\ C & \hat{Q} \end{bmatrix} \begin{bmatrix} \tau \\ \hat{\xi} \end{bmatrix} = \begin{bmatrix} -u^i \\ 0 \end{bmatrix} \quad \begin{aligned} \leftarrow \text{BC on } \partial\Omega \\ \leftarrow \text{FT-}y \text{ of discrep.} \end{aligned}$$

- fill B by evaluating $\hat{\mathcal{S}}, \hat{\mathcal{D}}$ Sommerfeld integrals at nodes $\mathbf{y}_j \in \partial\Omega$
- fill C by spectral rep. of each source $\mathbf{y}_j \in \partial\Omega$ at walls L, R

Results



(using Matlab toolbox MPSpack by B-Betcke '09)





$$d = 1.6\lambda \quad N = 70 \quad M = 80 \quad \text{error } 10^{-14} \quad t_{\text{fill}} = 0.13 \text{ s} \quad t_{\text{solve}} = 0.04 \text{ s}$$

Results

(using Matlab toolbox MPSpack by B-Betcke '09)





Exponential convergence:





$$d = 1.6\lambda \quad N = 70 \quad M = 80 \quad \text{error } 10^{-14} \quad t_{\text{fill}} = 0.13 \text{ s} \quad t_{\text{solve}} = 0.04 \text{ s}$$

Results

(using Matlab toolbox MPSpack by B-Betcke '09)




Exponential convergence:



$$d = 1.6\lambda \quad N = 70 \quad M = 80 \quad \text{error } 10^{-14} \quad t_{\text{fill}} = 0.13 \text{ s} \quad t_{\text{solve}} = 0.04 \text{ s}$$

- improved convergence rate by summing 1 or 2 $\partial\Omega$ neighbors directly
- low condition # $\sim 10^2$: solved to 14 digits in 55 GMRES iters.

Results: high aspect ratio



height = 10 periods

if lattice sums were used:
would need >20 neighbor copies of $\partial\Omega$
to be summed directly (slow)

$$d = 1.6\lambda \quad N = 300 \quad M = 250$$

$$\text{error } 10^{-10} \quad t_{\text{fill}} = 1 \text{ s} \quad t_{\text{solve}} = 0.5 \text{ s}$$

- Note, cond. # $\sim 10^6$:
close-to-resonant BVP
(not caused by scheme)


Handling Wood's anomalies ($k_n \rightarrow 0$)

recall k_n are Rayleigh–Bloch y -wavenumbers: $+k_n$ travels upwards, $-k_n$ downwards

Handling Wood's anomalies ($k_n \rightarrow 0$)

recall k_n are Rayleigh–Bloch y -wavenumbers: $+k_n$ travels upwards, $-k_n$ downwards

Sommerfeld FT- y
densities $\hat{\sigma}(k)$, $\hat{\tau}(k)$
have poles at $\pm k_n$:




Why?



Handling Wood's anomalies ($k_n \rightarrow 0$)

recall k_n are Rayleigh–Bloch y -wavenumbers: $+k_n$ travels upwards, $-k_n$ downwards

Sommerfeld FT- y
densities $\hat{\sigma}(k)$, $\hat{\tau}(k)$
have poles at $\pm k_n$:



Why?
$$\frac{i}{2\pi(k_n - k)} \stackrel{\text{FT}}{\leftrightarrow} \begin{cases} e^{ik_n y}, & y > 0 \\ 0, & y < 0 \end{cases} \quad \text{or} \quad \begin{cases} 0, & y > 0 \\ -e^{ik_n y}, & y < 0 \end{cases}$$




interpretation for $k_n > 0$: R–B outgoing above

incoming below



Handling Wood's anomalies ($k_n \rightarrow 0$)

recall k_n are Rayleigh–Bloch y -wavenumbers: $+k_n$ travels upwards, $-k_n$ downwards

Sommerfeld FT- y
densities $\hat{\sigma}(k)$, $\hat{\tau}(k)$
have poles at $\pm k_n$:



Why?
$$\frac{i}{2\pi(k_n - k)} \stackrel{\text{FT}}{\leftrightarrow} \begin{cases} e^{ik_n y}, & y > 0 \\ 0, & y < 0 \end{cases} \quad \text{or} \quad \begin{cases} 0, & y > 0 \\ -e^{ik_n y}, & y < 0 \end{cases}$$



interpretation for $k_n > 0$: R–B outgoing above


incoming below

Numerically: as $k_n \rightarrow 0$, grade nodes geometrically (sinh map) near 0

- works (10^{-9}), but: log blow-up of M , cond. # and $\|\hat{\xi}\|$ diverge (bad)

Well-conditioned handling of Wood's

deform contour to
be safe dist $O(1)$
from all poles:




Now solves BVP with wrong radiation condition!

enforces n^{th} R-B mode incoming below instead of outgoing above

Well-conditioned handling of Wood's

deform contour to
be safe dist $O(1)$
from all poles:



Now solves BVP with wrong radiation condition!

enforces n^{th} R-B mode incoming below instead of outgoing above

The fix:


- add plane-wave $ae^{ik_n x}e^{ik_n y}$, unknown a , to the $u(x, y)$ rep.
- add new linear condition: n^{th} amplitude incoming below = 0
project evaluation of $u(\cdot, -y_0)$ onto n^{th} Fourier mode

matrix now has 1 row and 1 column extra, solve for unknowns $[\tau; \hat{\xi}; a]$


at Wood $k_n = 0$: replace $\{e^{ik_n y}, e^{-ik_n y}\}$ by $\{1, y\}$, enforce “ y ” amplitude = 0

Works: cond. # and error bounded uniformly in params $(\omega, \theta^i) \dots$

Results precisely at Wood's anomaly



Pointwise error:
compared to indep. scheme



10^{-13} unless close to $\partial\Omega$


$$d = 1.6\lambda \quad N = 70 \quad M = 150 \quad \text{error } 10^{-14} \quad t_{\text{fill}} = 0.16 \text{ s} \quad t_{\text{solve}} = 0.1 \text{ s}$$

- cond # $< 10^4$, 150 GMRES iters. for 14 digits accuracy

MOVIES

Similar story for 2D Bloch eigenvalues (taste)

Similar story for 2D Bloch eigenvalues (taste)



Lattice of inclusions, index n in index 1


PDE EVP: u, u_n continuous across $\partial\Omega$

$$(\Delta + n^2\omega^2)u = 0 \text{ in } \Omega$$

$$(\Delta + \omega^2)u = 0 \text{ in } U \setminus \overline{\Omega}$$

u QP in both directions phases (α, β)

Similar story for 2D Bloch eigenvalues (taste)



Lattice of inclusions, index n in index 1


PDE EVP: u, u_n continuous across $\partial\Omega$

$$(\Delta + n^2\omega^2)u = 0 \text{ in } \Omega$$

$$(\Delta + \omega^2)u = 0 \text{ in } U \setminus \overline{\Omega}$$

u QP in both directions phases (α, β)

Photonic crystals: devices with period \approx wavelength of light



- control of propagation: bandgaps, dispersion, density of states
- filters, couplers, integrated optical computing, meta-materials, solar cells

Si, $\lambda = 1.6\mu\text{m}$ (Vlasov)

Need Bloch eigenvalues ('band structure'): are sheets in (ω, α, β)

Periodizing robustly

Single inclusion: use SLP+DLP densities η to rep fields in Ω and $\mathbb{R}^2 \setminus \bar{\Omega}$
integral op. $A\eta = [u^+ - u^-; u_n^+ - u_n^-]$, can make $A = I + \text{cpt}$ (Rokhlin '83)

As before, may periodize by replacing Φ_ω by $\Phi_{\omega,\text{QP}}$, A becomes A_{QP}

Periodizing robustly

Single inclusion: use SLP+DLP densities η to rep fields in Ω and $\mathbb{R}^2 \setminus \bar{\Omega}$
integral op. $A\eta = [u^+ - u^-; u_n^+ - u_n^-]$, can make $A = I + \text{cpt}$ (Rokhlin '83)

As before, may periodize by replacing Φ_ω by $\Phi_{\omega,\text{QP}}$, A becomes A_{QP}

Thm: if A_{QP} exists, $\text{Null } A_{\text{QP}} \neq \{0\} \Leftrightarrow (\omega, \alpha, \beta)$ Bloch eigenvalue

proof: non-physical fields are a swapped-media transmission BVP

Periodizing robustly

Single inclusion: use SLP+DLP densities η to rep fields in Ω and $\mathbb{R}^2 \setminus \bar{\Omega}$
integral op. $A\eta = [u^+ - u^-; u_n^+ - u_n^-]$, can make $A = I + \text{cpt}$ (Rokhlin '83)

As before, may periodize by replacing Φ_ω by $\Phi_{\omega,\text{QP}}$, A becomes A_{QP}

Thm: if A_{QP} exists, $\text{Null } A_{\text{QP}} \neq \{0\} \Leftrightarrow (\omega, \alpha, \beta)$ Bloch eigenvalue

proof: non-physical fields are a swapped-media transmission BVP

... but $\Phi_{\omega,\text{QP}}$ has poles at resonances of *empty* unit cell: scheme fails

Periodizing robustly

Single inclusion: use SLP+DLP densities η to rep fields in Ω and $\mathbb{R}^2 \setminus \bar{\Omega}$
integral op. $A\eta = [u^+ - u^-; u_n^+ - u_n^-]$, can make $A = I + \text{cpt}$ (Rokhlin '83)

As before, may periodize by replacing Φ_ω by $\Phi_{\omega,\text{QP}}$, A becomes A_{QP}

Thm: if A_{QP} exists, $\text{Null } A_{\text{QP}} \neq \{0\} \Leftrightarrow (\omega, \alpha, \beta)$ Bloch eigenvalue

proof: non-physical fields are a swapped-media transmission BVP

... but $\Phi_{\omega,\text{QP}}$ has poles at resonances of *empty* unit cell: scheme fails

Fix: use only free-space Φ_ω , add new densities on cell walls, enforce QP

$$\begin{bmatrix} A & B \\ C & Q \end{bmatrix} \begin{bmatrix} \eta \\ \xi \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad \begin{aligned} &\leftarrow \text{matching cond. on } \partial\Omega \\ &\leftarrow \text{discrep. in QP cond.: } f, f', g, g' \end{aligned}$$

- Robust (no spurious resonances), 2nd-kind sys, rapid convergence


How choose new densities on unit cell walls?

- to control 4 discrepancies (f, f', g, g')

need 4 densities $\xi = [\tau_L; \sigma_L; \tau_B; \sigma_B]$

$$Q = \frac{1}{2}\text{Id} + (\text{self-interactions}) + (\text{other interactions})$$

JRs $\sigma_L \rightarrow u|_L$ $\sigma_L \rightarrow u|_B$




How choose new densities on unit cell walls?

- to control 4 discrepancies (f, f', g, g')
need 4 densities $\xi = [\tau_L; \sigma_L; \tau_B; \sigma_B]$

$$Q = \frac{1}{2}\text{Id} + (\text{self-interactions}) + (\text{other interactions})$$

JRs $\sigma_L \rightarrow u|_L$ $\sigma_L \rightarrow u|_B$




- add phased ghost copies on other 2 walls

recall $f := u|_L - \alpha^{-1}u|_{L+\mathbf{e}_1}$

effect of σ_L on $u_n|_L$
effect of $\alpha\sigma_L$ on $\alpha^{-1}u_n|_{L+\mathbf{e}_1}$

} cancel apart from Id




How choose new densities on unit cell walls?

- to control 4 discrepancies (f, f', g, g')

need 4 densities $\xi = [\tau_L; \sigma_L; \tau_B; \sigma_B]$

$$Q = \frac{1}{2}\text{Id} + (\text{self-interactions}) + (\text{other interactions})$$

JRs	$\sigma_L \rightarrow u _L$	$\sigma_L \rightarrow u _B$
-----	-----------------------------	-----------------------------




- add phased ghost copies on other 2 walls

recall $f := u|_L - \alpha^{-1}u|_{L+\mathbf{e}_1}$

effect of σ_L on $u_n|_L$
 effect of $\alpha\sigma_L$ on $\alpha^{-1}u_n|_{L+\mathbf{e}_1}$

} cancel apart from Id




- add more ‘sticking-out’ ghost images

effect of  on $u_n|_L$

effect of α  on $\alpha^{-1}u_n|_{L+\mathbf{e}_1}$

} cancel apart from Id

\Rightarrow all corner interactions vanish!




- result: $Q = I + (\text{interactions of distance } \geq 1)$

\Rightarrow low rank, rapid convergence: 20-pt Gauss quadr. on $L, B \Rightarrow 10^{-12}$ error

Full Bloch solver scheme

(B-Greengard, JCP, *subm.*)


Finally we add 3×3 phased image copies of densities on $\partial\Omega$, giving:



Full Bloch solver scheme

(B-Greengard, JCP, *subm.*)

Finally we add 3x3 phased image copies of densities on $\partial\Omega$, giving:



- Careful cancellations: B, C, Q have only interactions of distance ≥ 1
- Large dist increases convergence rate, i.e. large c in error = $O(e^{-cN})$


Philosophy: sum neighboring image sources directly, so fields due to remainder of lattice have distant singularities

Results: small inclusion

band structure: simply plot log min sing. val. of M vs $(\omega, k_x, k_y) \dots$

Results: small inclusion

band structure: simply plot log min sing. val. of M vs $(\omega, k_x, k_y) \dots$



0.1 sec per eval
pre-store α, β coeffs
30 sec per
const- ω slice
 24×24 evals

$N = 40$ $M = 20$ (160 unknowns total) err 10^{-9}

MOVIE

Conclusions

- efficient 2nd-kind integral equations for 2D periodic problems
- periodize via small # extra degrees of freedom on cell walls
 - scattering: densities on unbounded walls via Fourier rep.
 - Bloch eigenvalue: kill corner singularities w/ tic-tac-toe
- more robust and flexible than quasi-periodic Greens function:
 - well-behaved at Wood's anomaly or spurious resonances
 - high aspect-ratios, extends simply to 3D, unlike lattice sums

Future:

- multi-layer; 3D; insert fast multipole methods for inclusion ...

code: <http://code.google.com/p/mpspack>
(B-Betcke, SIAM J. Sci. Comp. '10)

funding: NSF DMS-0507614
 DMS-0811005

B-Greengard, J. Comput. Phys., *subm.*
B-Greengard, BIT, *to be subm.*

<http://math.dartmouth.edu/~ahb>
made with: Linux, L^AT_EX, Prosper

EXTRA SLIDES

Main numerical approaches

Time domain

- a) time-stepping on finite-difference grid (FDTD) (e.g. Yee '66)
- low order (inaccurate); close freqs → need large t (inefficient)

Main numerical approaches

Time domain

- a) time-stepping on finite-difference grid (FDTD) (e.g. Yee '66)
low order (inaccurate); close freqs → need large t (inefficient)

Freq domain

- b) multiple-scattering, KKR, cylinders only (McPhedran, Moroz)
- c) Plane-wave method: all in Fourier space (Joannopoulos, Johnson, Sözüer)
discont. dielectric ⇒ Gibbs phenom, slow ($1/N$ or $1/N^2$) convergence
- d) Finite element (FEM) discretization in U (Chew, Dobson, Dossou)
better for discontinuity, N large, meshing complicated

Main numerical approaches

Time domain

- a) time-stepping on finite-difference grid (FDTD) (e.g. Yee '66)
low order (inaccurate); close freqs → need large t (inefficient)

Freq domain

- b) multiple-scattering, KKR, cylinders only (McPhedran, Moroz)
- c) Plane-wave method: all in Fourier space (Joannopoulos, Johnson, Sözüer)
discont. dielectric ⇒ Gibbs phenom, slow ($1/N$ or $1/N^2$) convergence
- d) Finite element (FEM) discretization in U (Chew, Dobson, Dossou)
better for discontinuity, N large, meshing complicated
- e) Integral equations: formulate problem *on* the discontinuity $\partial\Omega$
reduced dimensionality (small N)
high order (quadratures): high accuracy w/ small effort (⇒ sensitivity analysis)
scarcely used for band structure (Yuan '08)

Recast problem on compact domain (torus)

- Bloch wave condition equiv. to quasi-periodic BCs on ∂U


Require vanishing unit cell **discrepancy**:

$$f := u|_L - \alpha^{-1}u|_{L+\mathbf{e}_1} = 0$$

$$f' := u_n|_L - \alpha^{-1}u_n|_{L+\mathbf{e}_1} = 0$$

$$g := u|_B - \beta^{-1}u|_{B+\mathbf{e}_2} = 0$$


$$g' := u_n|_B - \beta^{-1}u_n|_{B+\mathbf{e}_2} = 0$$



Bloch phase parameters $\alpha := e^{i\mathbf{k}\cdot\mathbf{e}_1}$, $\beta := e^{i\mathbf{k}\cdot\mathbf{e}_2}$, $|\alpha| = |\beta| = 1$


- Task: find Bloch eigenvalue triples (ω, k_x, k_y) , i.e. (ω, α, β)

BIE for transmission scattering



Represent $u = u^i + \mathcal{D}\tau + \mathcal{S}\sigma$ outside wavenumber ω
 $u = \mathcal{D}_i\tau + \mathcal{S}_i\sigma$ inside wavenumber $n\omega$

BIE for transmission scattering



Represent $u = u^i + \mathcal{D}\tau + \mathcal{S}\sigma$ outside wavenumber ω
 $u = \mathcal{D}_i\tau + \mathcal{S}_i\sigma$ inside wavenumber $n\omega$

mismatch on $\partial\Omega$: $h := u^+ - u^-$, $h' := u_n^+ - u_n^-$

BCs: mismatch $m := [h; h']$ vanishes, use JRs...

$$\begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} u^i|_{\partial\Omega} \\ u_n^i|_{\partial\Omega} \end{bmatrix} + \underbrace{\begin{bmatrix} I & 0 \\ 0 & I \end{bmatrix}}_A + \underbrace{\begin{bmatrix} D - D_i & S_i - S \\ T - T_i & D_i^T - D^T \end{bmatrix}}_{\eta} \begin{bmatrix} \tau \\ -\sigma \end{bmatrix}$$

block 2nd-kind

A maps densities to their effect on mismatch

- hypersingular part of T cancels, so $A = \text{Id} + \text{compact}$ (Rokhlin '83)
- kernel weakly singular, but exists spectral product quadrature for $f(s) + \log(4 \sin^2 \frac{s}{2})g(s)$, f, g analytic 2π -periodic (Kress '91)

The standard way to periodize

replace kernel $\Phi_\omega(\mathbf{x})$ by $\Phi_{\omega,\text{QP}}(\mathbf{x}) := \sum_{m,n \in \mathbb{Z}} \alpha^m \beta^n \Phi(\mathbf{x} - m\mathbf{e}_1 - n\mathbf{e}_2)$

as before, integral operator A becomes A_{QP}

The standard way to periodize

replace kernel $\Phi_\omega(\mathbf{x})$ by $\Phi_{\omega,\text{QP}}(\mathbf{x}) := \sum_{m,n \in \mathbb{Z}} \alpha^m \beta^n \Phi(\mathbf{x} - m\mathbf{e}_1 - n\mathbf{e}_2)$

as before, integral operator A becomes A_{QP}

- as before, can evaluate via lattice sums, with effort
- seems natural (not yet used in literature)

The standard way to periodize

replace kernel $\Phi_\omega(\mathbf{x})$ by $\Phi_{\omega,\text{QP}}(\mathbf{x}) := \sum_{m,n \in \mathbb{Z}} \alpha^m \beta^n \Phi(\mathbf{x} - m\mathbf{e}_1 - n\mathbf{e}_2)$

as before, integral operator A becomes A_{QP}

- as before, can evaluate via lattice sums, with effort
- seems natural (not yet used in literature)

Theorem (*integral formulation of band structure*) :

If A_{QP} exists, $\text{Nul } A_{\text{QP}} \neq \{0\}$ $\Leftrightarrow (\omega, k_x, k_y)$ is eigenvalue

note: no u^i since eigenvalue problem homogeneous

proof: non-physical fields are a swapped-media transmission BVP

The standard way to periodize

replace kernel $\Phi_\omega(\mathbf{x})$ by $\Phi_{\omega,\text{QP}}(\mathbf{x}) := \sum_{m,n \in \mathbb{Z}} \alpha^m \beta^n \Phi(\mathbf{x} - m\mathbf{e}_1 - n\mathbf{e}_2)$

as before, integral operator A becomes A_{QP}

- as before, can evaluate via lattice sums, with effort
- seems natural (not yet used in literature)

Theorem (*integral formulation of band structure*) :

If A_{QP} exists, $\text{Nul } A_{\text{QP}} \neq \{0\}$ $\Leftrightarrow (\omega, k_x, k_y)$ is eigenvalue

note: no u^i since eigenvalue problem homogeneous

proof: non-physical fields are a swapped-media transmission BVP

Not a robust method: A_{QP} does not exist for certain parameters (ω, k_x, k_y)
since there $\Phi_{\omega,\text{QP}}(\mathbf{x}) \rightarrow \infty, \forall \mathbf{x}$

why...?

Failure at spurious resonances

$\Phi_{\omega,\text{QP}}(\mathbf{x})$ is Helmholtz Greens function in *empty* (index 1) torus

$$= \frac{1}{\text{Vol}(U)} \sum_{\mathbf{q} \in 2\pi\Lambda^*} \frac{e^{i(\mathbf{k}+\mathbf{q}) \cdot \mathbf{x}}}{\omega^2 - |\mathbf{k} + \mathbf{q}|^2} \quad \text{spectral representation on torus}$$


has simple pole wherever (ω, k_x, k_y) is eigenvalue of empty torus...
but physical field u well-behaved here: breakdown is **non-physical!**

Failure at spurious resonances

$\Phi_{\omega, \text{QP}}(\mathbf{x})$ is Helmholtz Greens function in *empty* (index 1) torus

$$= \frac{1}{\text{Vol}(U)} \sum_{\mathbf{q} \in 2\pi\Lambda^*} \frac{e^{i(\mathbf{k} + \mathbf{q}) \cdot \mathbf{x}}}{\omega^2 - |\mathbf{k} + \mathbf{q}|^2} \quad \text{spectral representation on torus}$$

has simple pole wherever (ω, k_x, k_y) is eigenvalue of empty torus...
but physical field u well-behaved here: breakdown is **non-physical!**



Robust way to periodize: back to free-space Φ_ω

represent $u = \mathcal{D}\tau + \mathcal{S}\sigma +$ (densities ξ on walls of U) outside

$$\begin{array}{c} \uparrow \\ \text{can enforce mismatch } m = 0 \end{array} \quad \begin{array}{c} \uparrow \\ \text{can enforce discrepancy } d := [f; f'; g; g'] = 0 \end{array}$$

Robust way to periodize: back to free-space Φ_ω

represent $u = \mathcal{D}\tau + \mathcal{S}\sigma +$ (densities ξ on walls of U) outside

$$\begin{array}{c} \uparrow \\ \text{can enforce mismatch } m = 0 \end{array} \quad \begin{array}{c} \uparrow \\ \text{can enforce discrepancy } d := [f; f'; g; g'] = 0 \end{array}$$

In block operator form

$$\underbrace{\begin{bmatrix} A & B \\ C & Q \end{bmatrix}}_M \begin{bmatrix} \eta \\ \xi \end{bmatrix} = \begin{bmatrix} m \\ d \end{bmatrix}$$

- added extra degrees of freedom (a small #, indep. of complexity of Ω)

Robust way to periodize: back to free-space Φ_ω

represent $u = \mathcal{D}\tau + \mathcal{S}\sigma +$ (densities ξ on walls of U) outside

$$\begin{array}{c} \uparrow \\ \text{can enforce mismatch } m = 0 \end{array} \quad \begin{array}{c} \uparrow \\ \text{can enforce discrepancy } d := [f; f'; g; g'] = 0 \end{array}$$

In block operator form

$$\underbrace{\begin{bmatrix} A & B \\ C & Q \end{bmatrix}}_M \begin{bmatrix} \eta \\ \xi \end{bmatrix} = \begin{bmatrix} m \\ d \end{bmatrix}$$

- added extra degrees of freedom (a small #, indep. of complexity of Ω)
- gain robustness: no matrix element blow-up at spurious resonances


Observe:

$$\text{Nul } M \neq \{0\} \iff (\omega, k_x, k_y) \text{ Bloch eigenvalue}$$

- idea of extra sources of waves not new (*e.g.* Hafner '02)
- what is new: $M = \text{Id} + \text{compact}$ ideal for large-scale, iterative, FMM

Error convergence


$\log_{10} \min \text{ sing. val } M$ for known Bloch eigenvalue (should be zero):



Note: is eigenvalue error
up to $O(1)$ const

$$\omega = 5, \mathbf{k} \approx (-0.39, 2.08)$$

mode:



- spectral (exponential) convergence in inclusion & wall # dofs



Interpolation across the Brillouin zone

How find eigenvalue sheets in the volume $S^1 \times S^1 \times (0, \omega_{\max})$?

Interpolation across the Brillouin zone

How find eigenvalue sheets in the volume $S^1 \times S^1 \times (0, \omega_{\max})$?

E.g. iterative rootfinding on min sing. val. of M :




ω -slice
(α, β fixed)

- slow (20 its per root), unreliable (misses nearby roots)

Interpolation across the Brillouin zone

How find eigenvalue sheets in the volume $S^1 \times S^1 \times (0, \omega_{\max})$?

E.g. iterative rootfinding on min sing. val. of M :



- slow (20 its per root), unreliable (misses nearby roots)

Realize: $M = I + (\text{cpt op-valued analytic func of } \omega, k_x \text{ and } k_y)$

$\det M$ is a **Fredholm determinant**, also analytic


- rootfinding a real-analytic function is nice...

(J. Boyd '02)

Spectral rootfinding of analytic functions

(Boyd '02)

$f : \mathbb{R} \rightarrow \mathbb{R}$, 2π -periodic




Spectral rootfinding of analytic functions

(Boyd '02)

$f : \mathbb{R} \rightarrow \mathbb{R}$, 2π -periodic

use trigonometric poly interpolant



$$f(\theta) \approx \sum_{n=-N}^N c_n e^{in\theta}$$


exponentially convergent in N
width of strip about \mathbb{R} in which f analytic

Spectral rootfinding of analytic functions

(Boyd '02)

$f : \mathbb{R} \rightarrow \mathbb{R}$, 2π -periodic

use trigonometric poly interpolant



$$f(\theta) \approx \sum_{n=-N}^N c_n e^{in\theta}$$

exponentially convergent in N
width of strip about \mathbb{R} in which f analytic


- c_n found by FFT of grid samples $f(\pi n/N)$, $n = 1, \dots, 2N$

Spectral rootfinding of analytic functions

(Boyd '02)

$f : \mathbb{R} \rightarrow \mathbb{R}$, 2π -periodic

use trigonometric poly interpolant



$$f(\theta) \approx \sum_{n=-N}^N c_n e^{in\theta}$$

exponentially convergent in N
width of strip about \mathbb{R} in which f analytic

- c_n found by FFT of grid samples $f(\pi n/N)$, $n = 1, \dots, 2N$

map $e^{i\theta} = z \in \mathbb{C}$

\Rightarrow Laurent $q(z) = \sum_{n=-N}^N c_n z^n$

roots of f lie on $|z| = 1$


has roots *near* $|z| = 1$

Spectral rootfinding of analytic functions

(Boyd '02)

$f : \mathbb{R} \rightarrow \mathbb{R}$, 2π -periodic

use trigonometric poly interpolant



$$f(\theta) \approx \sum_{n=-N}^N c_n e^{in\theta}$$

exponentially convergent in N
width of strip about \mathbb{R} in which f analytic

- c_n found by FFT of grid samples $f(\pi n/N)$, $n = 1, \dots, 2N$

map $e^{i\theta} = z \in \mathbb{C}$

roots of f lie on $|z| = 1$

\Rightarrow Laurent $q(z) = \sum_{n=-N}^N c_n z^n$

has roots *near* $|z| = 1$


“Degree doubling”: $z^N q(z)$ is degree- $2N$ poly, so...

- use Matlab `roots` QR for `eigvals` of companion matrix, $O(N^3)$ but v. stable
- extract the angles θ of roots near unit circle

(Boyd nonlin EVP; Trefethen-Battles '06 `chebfun`)

Rootfinding $\det M$ in the ω direction


$\det M(\omega, k_x, k_y)$ not periodic in ω : map $\omega = \omega_0 + a \cos \theta$ periodic θ
this is Chebyshev interpolation on interval $[\omega_0 - a, \omega_0 + a]$




Fix \mathbf{k} , eval $\det M$ at Cheby pts, get $\omega_j(\mathbf{k})$ in interval
25 evals covers $\omega \in [4, 6]$, i.e. 10-20 evals per root found

Rootfinding $\det M$ in the ω direction


$\det M(\omega, k_x, k_y)$ not periodic in ω : map $\omega = \omega_0 + a \cos \theta$ periodic θ
this is Chebyshev interpolation on interval $[\omega_0 - a, \omega_0 + a]$



Fix \mathbf{k} , eval $\det M$ at Cheby pts, get $\omega_j(\mathbf{k})$ in interval
25 evals covers $\omega \in [4, 6]$, i.e. 10-20 evals per root found




Also analytic in $k_x, k_y \Rightarrow$ interpolate in 3D!



Robust spectrally-accurate bands via small # grid evals
e.g. $25 \times 24 \times 24$ for $\omega \in [4, 6]$ and whole Brillouin zone, error 10^{-8}

Band structure to spectral accuracy



$n=0.3$ inside

$n=1$ outside

large inclusion

eval only 24×24 samples in \mathbf{k}
but contains much finer details

10^{-8} errors, 1 hour on laptop

- Note: eigenvalues $\omega_j(\mathbf{k})$ are **not** analytic!
 - \exists conical (diabolical) points ... interpolates poorly
- like level set method: handle smooth func

movie 1

Software environment (teaser)

Software environment (teaser)

MPSpack: object-oriented 2D PDE toolbox in Matlab (B-Betcke '09)

- implements above & more: Helmholtz, Laplace, scattering
- intuitive interface: curves, domains, basis sets, problems, are **objects**

E.g.

Software environment (teaser)

MPSpack: object-oriented 2D PDE toolbox in Matlab (B-Betcke '09)

- implements above & more: Helmholtz, Laplace, scattering
- intuitive interface: curves, domains, basis sets, problems, are **objects**

E.g.

```
s=segment.radialfunc(100,{@(q) 1+.3*cos(3*q), @(q) -.9*sin(3*q)})  
piecewise analytic curves
```

Software environment (teaser)

MPSpack: object-oriented 2D PDE toolbox in Matlab (B-Betcke '09)

- implements above & more: Helmholtz, Laplace, scattering
- intuitive interface: curves, domains, basis sets, problems, are objects

E.g.

```
s=segment.radialfunc(100,{@(q)1+.3*cos(3*q),@(q)-.9*sin(3*q)})  
piecewise analytic curves
```

```
d = domain([],[],s,-1);
```

make exterior domain using segment

Software environment (teaser)

MPSpack: object-oriented 2D PDE toolbox in Matlab (B-Betcke '09)

- implements above & more: Helmholtz, Laplace, scattering
- intuitive interface: curves, domains, basis sets, problems, are **objects**


E.g.

```
s=segment.radialfunc(100,{@(q) 1+.3*cos(3*q), @(q) -.9*sin(3*q)})
```

piecewise analytic curves

```
d = domain([],[],s,-1);
```

make exterior domain using segment



Software environment (teaser)

MPSpack: object-oriented 2D PDE toolbox in Matlab (B-Betcke '09)

- implements above & more: Helmholtz, Laplace, scattering
- intuitive interface: curves, domains, basis sets, problems, are **objects**

E.g.

```
s=segment.radialfunc(100,{@(q) 1+.3*cos(3*q), @(q) -.9*sin(3*q)})
```


piecewise analytic curves

```
d = domain([],[],s,-1);
```

make exterior domain using segment

```
d.setbc(1,'N',[]);
```

Neumann BCs on exterior



Software environment (teaser)

MPSpack: object-oriented 2D PDE toolbox in Matlab (B-Betcke '09)

- implements above & more: Helmholtz, Laplace, scattering
- intuitive interface: curves, domains, basis sets, problems, are **objects**

E.g.

```
s=segment.radialfunc(100,{@(q) 1+.3*cos(3*q), @(q) -.9*sin(3*q)})
```

piecewise analytic curves

```
d = domain([],[],s,-1);
```


make exterior domain using segment

```
d.setbc(1, 'N', []);
```

Neumann BCs on exterior

```
d.addmfsbasis(s, 200, struct('tau',0.05));
```

choose basis set for solution



Software environment (teaser)

MPSpack: object-oriented 2D PDE toolbox in Matlab (B-Betcke '09)

- implements above & more: Helmholtz, Laplace, scattering
- intuitive interface: curves, domains, basis sets, problems, are **objects**


E.g.

```
s=segment.radialfunc(100,{@(q) 1+.3*cos(3*q), @(q) -.9*sin(3*q)})  
piecewise analytic curves
```

```
d = domain([],[],s,-1);  
make exterior domain using segment
```

```
d.setbc(1, 'N', []);  
Neumann BCs on exterior
```

```
d.addmfsbasis(s, 200, struct('tau',0.05));  
choose basis set for solution
```



```
p = scattering(d, []);
```

make a scattering problem from domain d

```
p = scattering(d, []);
```

make a scattering problem from domain d

```
p.setoverallwavenumber(30); p.setincidentwave(pi/6);
```

```
p = scattering(d, []);
```

make a scattering problem from domain d

```
p.setoverallwavenumber(30); p.setincidentwave(pi/6);
```

```
p.solvecoeffs; p.bcresidualnorm
```

fills matrix, solves in 0.1 sec, L^2 error 6×10^{-9}

```
p = scattering(d, []);
```


make a scattering problem from domain d

```
p.setoverallwavenumber(30); p.setincidentwave(pi/6);
```

```
p.solvecoeffs; p.bcresidualnorm
```

fills matrix, solves in 0.1 sec, L^2 error 6×10^{-9}

```
p.showfullfield(struct('bb', [-2 2 -2 2]));
```



```
p = scattering(d, []);
```


make a scattering problem from domain d

```
p.setoverallwavenumber(30); p.setincidentwave(pi/6);
```

```
p.solvecoeffs; p.bcresidualnorm
```

fills matrix, solves in 0.1 sec, L^2 error 6×10^{-9}

```
p.showfullfield(struct('bb', [-2 2 -2 2]));
```



- was easy case: 8 lines (could have done in 80 lines of Matlab)
- multiple (sub)domains: basis, quadrature, bookkeeping hidden
e.g. dielectric band structure still only 20 lines of code
- human-readable, rapid to code, sensible defaults (you can change)

To do: document periodic BVPs, meshing, Dirichlet EVP, ...