

Improper Integrals

01/10/2007

Improper Integrals

- If the interval of integration of f is infinite or f has an infinite discontinuity in $[a, b]$, then the definite integral of f is called an *improper integral*.

Type I: Infinite intervals

- If $\int_a^t f(x)dx$ exists for every number $t \geq a$, then

$$\int_a^\infty f(x)dx = \lim_{t \rightarrow \infty} \int_a^t f(x)dx,$$

provided this limit exists (as a finite number).

- If $\int_t^b f(x)dx$ exists for every number $t \leq b$, then

$$\int_{-\infty}^b f(x)dx = \lim_{t \rightarrow -\infty} \int_t^b f(x)dx$$

provided this limit exists (as a finite number).

- The improper intervals $\int_a^\infty f(x)dx$ and $\int_{-\infty}^b f(x)dx$ are called **convergent** if the corresponding limit exists and **divergent** if the limit does not exist.

- If $\int_t^b f(x)dx$ exists for every number $t \leq b$, then

$$\int_{-\infty}^b f(x)dx = \lim_{t \rightarrow -\infty} \int_t^b f(x)dx$$

provided this limit exists (as a finite number).

- The improper intervals $\int_a^\infty f(x)dx$ and $\int_{-\infty}^b f(x)dx$ are called **convergent** if the corresponding limit exists and **divergent** if the limit does not exist.
- If both $\int_a^\infty f(x)dx$ and $\int_{-\infty}^b f(x)dx$ are convergent, then we define

$$\int_{-\infty}^\infty f(x)dx = \int_{-\infty}^a f(x)dx + \int_a^\infty f(x)dx.$$

Examples

- Determine whether the following integrals are convergent or divergent:

- $\int_1^{\infty} \frac{1}{x^2} dx$

Examples

- Determine whether the following integrals are convergent or divergent:

- $\int_1^{\infty} \frac{1}{x^2} dx$

- $\int_1^{\infty} \frac{1}{x} dx$

Examples

- Determine whether the following integrals are convergent or divergent:

- $\int_1^{\infty} \frac{1}{x^2} dx$

- $\int_1^{\infty} \frac{1}{x} dx$

- $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$

Examples

- Determine whether the following integrals are convergent or divergent:
- $\int_1^{\infty} \frac{1}{x^2} dx$
- $\int_1^{\infty} \frac{1}{x} dx$
- $\int_{-\infty}^{\infty} \frac{1}{1+x^2} dx$
- Is the area under the curve $y = \frac{\ln x}{x^2}$ from $x = 1$ to $x = \infty$ finite? If so, what is its value?

- For what values of p is the integral

$$\int_1^{\infty} \frac{1}{x^p} dx$$

convergent?