

Functions of Several Variables

Lecture 21

November 6, 2006

Functions of two variables

Functions of two variables

- A function of two variables is a rule that assigns to each ordered pair of real numbers (x, y) in a set D a unique real number denoted by $f(x, y)$.

Functions of two variables

- A function of two variables is a rule that assigns to each ordered pair of real numbers (x, y) in a set D a unique real number denoted by $f(x, y)$.
- The set D is the domain of f and its range is the set of values that f takes on.

Functions of two variables

- A function of two variables is a rule that assigns to each ordered pair of real numbers (x, y) in a set D a unique real number denoted by $f(x, y)$.
- The set D is the domain of f and its range is the set of values that f takes on.
- We also write $z = f(x, y)$

Functions of two variables

- A function of two variables is a rule that assigns to each ordered pair of real numbers (x, y) in a set D a unique real number denoted by $f(x, y)$.
- The set D is the domain of f and its range is the set of values that f takes on.
- We also write $z = f(x, y)$
- The variables x and y are **independent variables** and z is the **dependent variable**.

Examples

Examples

- Find the domain of the function

$$f(x, y) = \frac{2x + 3y}{x^2 + y^2 - 9}$$

Examples

- Find the domain of the function

$$f(x, y) = \frac{2x + 3y}{x^2 + y^2 - 9}$$

- Find the domain and range of

$$f(x, y) = \sqrt{4 - x^2 - y^2}$$

Definition

- If f is a function of two variables with domain D , then the graph of f is the set of all points $(x, y, z) \in \mathbb{R}^3$ such that $z = f(x, y)$ and (x, y) is in D .

Example

Example

- A **linear function** is a function

$$f(x) = ax + by + c$$

Example

- A **linear function** is a function

$$f(x) = ax + by + c$$

- The graph of such a function is a plane.


Example

- $f(x, y) = \sin(x) + \sin(y)$

Examples

Example

- $f(x, y) = \sin(x) + \sin(y)$




Example

- $f(x, y) = (x^2 + y^2)e^{-x^2 - y^2}$

Example

- $f(x, y) = (x^2 + y^2)e^{-x^2 - y^2}$



The Cobb-Douglas production function


Example

- $P(L, K) = bL^\alpha K^{1-\alpha}$

The Cobb-Douglas production function

Example

- $P(L, K) = bL^\alpha K^{1-\alpha}$
- $P(L, K) = 1.01L^{0.75}K^{0.25}$



Definition


- The **level curves** of a function f of two variables are the curves with equations $f(x, y) = k$, where k is constant.

Example

- $f(x, y) = \sin(x) + \sin(y)$

Example


- $f(x, y) = \sin(x) + \sin(y)$



Examples

Example


- $f(x, y) = (x^2 + y^2)e^{-x^2 - y^2}$



The Cobb-Douglas production function

Example

- $P(L, K) = 1.01L^{0.75}K^{0.25}$



Definition

Definition

- We say that a function $f(x, y)$ has limit L as (x, y) approaches a point (a, b) and we write

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L$$

if we can make the values of $f(x, y)$ as close to L as we like by taking the point (x, y) sufficiently close to the point (a, b) , but not equal to (a, b) .

Definition


- We say that a function $f(x, y)$ has limit L as (x, y) approaches a point (a, b) and we write

$$\lim_{(x,y) \rightarrow (a,b)} f(x, y) = L$$

if we can make the values of $f(x, y)$ as close to L as we like by taking the point (x, y) sufficiently close to the point (a, b) , but not equal to (a, b) .

- We write also $f(x, y) \rightarrow L$ as $(x, y) \rightarrow (a, b)$ and

$$\lim_{x \rightarrow a, y \rightarrow b} f(x, y) = L$$



- If $f(x, y) \rightarrow L_1$ as $(x, y) \rightarrow (a, b)$ along a path C_1 and $f(x, y) \rightarrow L_2$ as $(x, y) \rightarrow (a, b)$ along a path C_2 , where $L_1 \neq L_2$, then $\lim_{(x,y) \rightarrow (a,b)} f(x, y)$ does not exist.

- If $f(x, y) \rightarrow L_1$ as $(x, y) \rightarrow (a, b)$ along a path C_1 and $f(x, y) \rightarrow L_2$ as $(x, y) \rightarrow (a, b)$ along a path C_2 , where $L_1 \neq L_2$, then $\lim_{(x,y) \rightarrow (a,b)} f(x, y)$ does not exist.
- **Example:** Show that

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 - y^2}{x^2 + y^2}$$

does not exist.

Definition

Definition

- A function f of two variables is called **continuous at** (a, b) if

$$\lim_{(x,y) \rightarrow (a,b)} f(x,y) = f(a,b)$$

Definition

- A function f of two variables is called **continuous at** (a, b) if

$$\lim_{(x,y) \rightarrow (a,b)} f(x,y) = f(a,b)$$

- **Examples:** polynomials, rational, trigonometric, exponential, logarithmic functions are continuous on their domain.

Examples

Examples

- Find the limit

$$\lim_{(x,y) \rightarrow (0,0)} \frac{2x^2 - 4y}{\sqrt{2x^2 - 4y + 1} - 1}$$

Examples

- Find the limit

$$\lim_{(x,y) \rightarrow (0,0)} \frac{2x^2 - 4y}{\sqrt{2x^2 - 4y + 1} - 1}$$

- Find the largest set on which the function

$$\frac{2xy}{9 - x^2 - y^2}$$

is continuous.