## Math 13. Multivariable Calculus. Written Homework 9.

Due on Wednesday, 5/29/13.

You may leave this homework in the boxes outside of Kemeny 108 by 1:45 pm on Wednesday. Please write problems 1-3 on separate pages from problems 4-6 and turn them in in the corresponding columns.

- 1. (Chapter 16.7, #29) Let  $\mathbf{F} = \langle x, 2y, 3z \rangle$ , and let S be the cube with vertices  $(\pm 1, \pm 1, \pm 1)$  with positive orientation. Evaluate  $\iint_S \mathbf{F} \cdot d\mathbf{S}$ .
- 2. (Chapter 16.7, #49) Let **F** be an inverse square vector field (that is,  $\mathbf{F}(\mathbf{r}) = c\mathbf{r}/|\mathbf{r}|^3$  for some constant c, where  $\mathbf{r} = \langle x, y, z \rangle$ ). Show that the flux of **F** across a sphere S centered at the origin is independent of the radius of S.
- 3. (Chapter 16.9, #18) Let  $\mathbf{F} = \langle z \tan^{-1}(y^2), z^3 \ln(x^2+1), z \rangle$ . Find the flux of  $\mathbf{F}$  across the part of the paraboloid  $x^2 + y^2 + z = 2$  that lies above the plane z = 1 and is oriented upwards.
- 4. (Chapter 16.9, #24) Use the Divergence Theorem to evaluate

$$\iint_{S} (2x + 2y + z^2) \, dS,$$

where S is the sphere  $x^2 + y^2 + z^2 = 1$ .

- 5. (Chapter 16.8, #17) A particle moves along line segments from the origin to the points (1,0,0), (1,2,1), (0,2,1), and then back to the origin under the influence of the force field  $\mathbf{F} = \langle z^2, 2xy, 4y^2 \rangle$ . Find the work done in two separate ways: (a) by directly calculating this line integral, and (b) by using Stokes' Theorem with a suitable choice of surface S.
- 6. (Chapter 16.8, #18) Evaluate

$$\int_{C} (y + \sin x) \, dx + (z^{2} + \cos y) \, dy + x^{3} \, dz,$$

where C is the curve  $\mathbf{r}(t) = \langle \sin t, \cos t, \sin 2t \rangle$ ,  $0 \le t \le 2\pi$ . (Hint: Observe that C lies on the surface z = 2xy.)