

Intro

Logic  
Insufficient data  
Boolean algebra  
Probability notation

Justifications

Betting strategy  
Subjective belief  
Objective frequency  
Many worlds

Bayes

Derivation  
WHy use Bayes?  
Caveat

Expectation

Definition  
Properties

Invariance

Puzzles

# Reasoning under uncertainty

David “davidad” Dalrymple  
davidad@alum.mit.edu

August 20, 2017  
<http://espr.cf>

Intro

Logic  
Insufficient data  
Boolean algebra  
Probability notation

Justifications

Betting strategy  
Subjective belief  
Objective frequency  
Many worlds

Bayes

Derivation  
WHy use Bayes?  
Caveat

Expectation

Definition  
Properties

Invariance

Puzzles

# Reasoning under uncertainty (Foundations of probability theory)

David “davidad” Dalrymple  
davidad@alum.mit.edu

August 20, 2017  
<http://espr.cf>

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

# ① All synapses are either electrical or chemical

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- 1 All synapses are either electrical or chemical
- 2 All chemical synapses are intercellular

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① All synapses are either electrical or chemical
- ② All chemical synapses are intercellular
- ③ If a synapse is not intercellular, it is not an electrical synapse

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① All synapses are either electrical or chemical
- ② All chemical synapses are intercellular
- ③ If a synapse is not intercellular, it is not an electrical synapse
- ④ Are all synapses intercellular?

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① All synapses are either electrical or chemical
- ② All chemical synapses are intercellular
- ③ If a synapse is not intercellular, it is not an electrical synapse
- ④ Are all synapses intercellular (yes or no)?



## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- 1 3 boxes are labeled “Red balls,” “Green balls,” and “Red and green balls.”

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- 1 3 boxes are labeled “Red balls,” “Green balls,” and “Red and green balls.”
- 2 All three labels are incorrect;

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- 1 3 boxes are labeled “Red balls,” “Green balls,” and “Red and green balls.”
- 2 All three labels are incorrect; each belongs on a different box.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- 1 3 boxes are labeled “Red balls,” “Green balls,” and “Red and green balls.”
- 2 All three labels are incorrect; each belongs on a different box.
- 3 You may ask me to take one ball from one box.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- 1 3 boxes are labeled “Red balls,” “Green balls,” and “Red and green balls.”
- 2 All three labels are incorrect; each belongs on a different box.
- 3 You may ask me to take one ball from one box.
- 4 How can you label the boxes correctly?

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- 1 I have a full deck of 52 playing cards, and take the top card.



## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- 1 I have a full deck of 52 playing cards, and take the top card.
- 2 Is it a queen?

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles



<http://xkcd.com/1448>; reference is to “The Last Question” by Isaac Asimov

## Intro

## Logic

## Insufficient data

## Boolean algebra

## Probability notation

## Justifications

## Betting strategy

## Subjective belief

## Objective frequency

## Many worlds

## Bayes

## Derivation

## WHy use Bayes?

## Caveat

## Expectation

## Definition

## Properties

## Invariance

## Puzzles

Different framings of this question:

- ① **Betting strategy**
- ② **Subjective belief**
- ③ **Objective frequency**
- ④ **Many (deterministic) worlds**

## Intro

## Logic

## Insufficient data

## Boolean algebra

## Probability notation

## Justifications

## Betting strategy

## Subjective belief

## Objective frequency

## Many worlds

## Bayes

## Derivation

## WHy use Bayes?

## Caveat

## Expectation

## Definition

## Properties

## Invariance

## Puzzles

Different framings of this question:

- ① **Betting strategy**
  - You're required to set prices for bets. How can you not let an equally-informed opponent guarantee a profit off you?
- ② **Subjective belief**
- ③ **Objective frequency**
- ④ **Many (deterministic) worlds**

## Different framings of this question:

### ① Betting strategy

- You're required to set prices for bets. How can you not let an equally-informed opponent guarantee a profit off you?

### ② Subjective belief

- Assign quantitative “degrees of belief” instead of truth values, as an extension of logic.

### ③ Objective frequency

### ④ Many (deterministic) worlds

## Different framings of this question:

### ① Betting strategy

- You're required to set prices for bets. How can you not let an equally-informed opponent guarantee a profit off you?

### ② Subjective belief

- Assign quantitative “degrees of belief” instead of truth values, as an extension of logic.

### ③ Objective frequency

- Consider hypothetically repeating a random experiment (infinitely) many times, and counting the number of outcomes where a proposition is true.

### ④ Many (deterministic) worlds

## Different framings of this question:

### ① Betting strategy

- You're required to set prices for bets. How can you not let an equally-informed opponent guarantee a profit off you?

### ② Subjective belief

- Assign quantitative “degrees of belief” instead of truth values, as an extension of logic.

### ③ Objective frequency

- Consider hypothetically repeating a random experiment (infinitely) many times, and counting the number of outcomes where a proposition is true.

### ④ Many (deterministic) worlds

- Any given world would have enough information for deduction, but we don't know which world we're in. Count the number of worlds where a proposition is true.

Different framings of this question:

### ① Betting strategy

- You're required to set prices for bets. How can you not let an equally-informed opponent guarantee a profit off you?

### ② Subjective belief

- Assign quantitative “degrees of belief” instead of truth values, as an extension of logic.

### ③ Objective frequency

- Consider hypothetically repeating a random experiment (infinitely) many times, and counting the number of outcomes where a proposition is true.

### ④ Many (deterministic) worlds

- Any given world would have enough information for deduction, but we don't know which world we're in. Count the number of worlds where a proposition is true.



Different framings of this question:

### ① Betting strategy

- You're required to set prices for bets. How can you not let an equally-informed opponent guarantee a profit off you?

### ② Subjective belief

- Assign quantitative “degrees of belief” instead of truth values, as an extension of logic.

### ③ Objective frequency

- Consider hypothetically repeating a random experiment (infinitely) many times, and counting the number of outcomes where a proposition is true.

### ④ Many (deterministic) worlds

- Any given world would have enough information for deduction, but we don't know which world we're in. Count the number of worlds where a proposition is true.

Big takeaway: these framings **all justify equivalent formal theories**, i.e. probability!

Different framings of this question:

### ① Betting strategy

- You're required to set prices for bets. How can you not let an equally-informed opponent guarantee a profit off you?

### ② Subjective belief

- Assign quantitative “degrees of belief” instead of truth values, as an extension of logic.

### ③ Objective frequency

- Consider hypothetically repeating a random experiment (infinitely) many times, and counting the number of outcomes where a proposition is true.

### ④ Many (deterministic) worlds

- Any given world would have enough information for deduction, but we don't know which world we're in. Count the number of worlds where a proposition is true.

Big takeaway: these framings **all justify equivalent formal theories**, i.e. probability!

Note: there are other framings that justify different theories, like Dempster-Shafer theory, but this concordance is still surprising.

## Intro

## Logic

## Insufficient data

## Boolean algebra

## Probability notation

## Justifications

## Betting strategy

## Subjective belief

## Objective frequency

## Many worlds

## Bayes

## Derivation

## WHy use Bayes?

## Caveat

## Expectation

## Definition

## Properties

## Invariance

## Puzzles

Logic		Algebra		Sets	
conjunction	$\wedge$	multiplication	$\cdot$	intersection	$\cap$
disjunction	$\vee$	addition	$+$	union	$\cup$
negation	$\neg$	negation	$\neg$	complement	$\Omega \setminus$

## Intro

## Logic

## Insufficient data

## Boolean algebra

## Probability notation

## Justifications

## Betting strategy

## Subjective belief

## Objective frequency

## Many worlds

## Bayes

## Derivation

## WHy use Bayes?

## Caveat

## Expectation

## Definition

## Properties

## Invariance

## Puzzles

$\top$  means true;  $\perp$  means false

$$A \vee \perp = A$$

$$A \wedge \top = A$$

$$A \vee B = B \vee A$$

$$A \wedge B = B \wedge A$$

$$A \vee (B \wedge C) = (A \vee B) \wedge (A \vee C) \quad A \wedge (B \vee C) = (A \wedge B) \vee (A \wedge C)$$

$$A \vee \neg A = \top$$

$$A \wedge \neg A = \perp$$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- $\mathbb{P}(A|B)$  is the “conditional probability” of a proposition (or “event”)  $A$  given certain knowledge of another proposition  $B$ .

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- $\mathbb{P}(A|B)$  is the “conditional probability” of a proposition (or “event”)  $A$  given certain knowledge of another proposition  $B$ .
- $\mathbb{P}(A)$  is the “unconditional probability” of a proposition, often interpreted as  $\mathbb{P}(A|\Omega)$ , where  $\Omega$  is interpreted as a set of assumptions considered to be unconditional.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- $\mathbb{P}(A|B)$  is the “conditional probability” of a proposition (or “event”)  $A$  given certain knowledge of another proposition  $B$ .
- $\mathbb{P}(A)$  is the “unconditional probability” of a proposition, often interpreted as  $\mathbb{P}(A|\Omega)$ , where  $\Omega$  is interpreted as a set of assumptions considered to be unconditional.
- Comma is sometimes used for  $\wedge$ , e.g.  $\mathbb{P}(A, B|\Omega) := \mathbb{P}(A \wedge B|\Omega)$ .

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

In each framing/justification, I'll state the specific laws of probability, then go over the argument.



## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

1 (Nonnegativity)  $\mathbb{P}(A|\Omega) \geq 0$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① (Nonnegativity)  $\mathbb{P}(A|\Omega) \geq 0$
- ② (Tautology)  $\mathbb{P}(A|\Omega) = 1$  if  $A$  is a logical truth given  $\Omega$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① (Nonnegativity)  $\mathbb{P}(A|\Omega) \geq 0$
- ② (Tautology)  $\mathbb{P}(A|\Omega) = 1$  if  $A$  is a logical truth given  $\Omega$
- ③ (Countable additivity) For arbitrary disjoint  $A_i$ ,  
$$\mathbb{P}\left(\bigvee_{i=1}^{\infty} A_i \middle| \Omega\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i|\Omega)$$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① (Nonnegativity)  $\mathbb{P}(A|\Omega) \geq 0$
- ② (Tautology)  $\mathbb{P}(A|\Omega) = 1$  if  $A$  is a logical truth given  $\Omega$
- ③ (Countable additivity) For arbitrary disjoint  $A_i$ ,  
$$\mathbb{P}\left(\bigvee_{i=1}^{\infty} A_i \middle| \Omega\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i|\Omega)$$
- ④ (Conditional) 
$$\mathbb{P}(A|B, \Omega) = \frac{\mathbb{P}(A \wedge B|\Omega)}{\mathbb{P}(B|\Omega)}$$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

The rules of the game:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

The rules of the game:

- 1 Agent must state  $p(A)$  for all  $A$ , such that Agent would buy or sell arbitrarily (possibly fractionally) many bets on  $A$  with probability  $p(A)$

Intro

Logic

Insufficient data

Boolean algebra

Probability notation

Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

Bayes

Derivation

WHy use Bayes?

Caveat

Expectation

Definition

Properties

Invariance

Puzzles

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

The rules of the game:

- 1 Agent must state  $p(A)$  for all  $A$ , such that Agent would buy or sell arbitrarily (possibly fractionally) many bets on  $A$  with probability  $p(A)$
- 2 Bookie may then buy or sell Agent's bets on any  $As$



A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

The rules of the game:

- 1 Agent must state  $p(A)$  for all  $A$ , such that Agent would buy or sell arbitrarily (possibly fractionally) many bets on  $A$  with probability  $p(A)$
- 2 Bookie may then buy or sell Agent's bets on any  $A$ s
- 3 Finally, an observation is made and the bets are settled

Intro

Logic

Insufficient data

Boolean algebra

Probability notation

Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

Bayes

Derivation

WHy use Bayes?

Caveat

Expectation

Definition

Properties

Invariance

Puzzles

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

The rules of the game:

- 1 Agent must state  $p(A)$  for all  $A$ , such that Agent would buy or sell arbitrarily (possibly fractionally) many bets on  $A$  with probability  $p(A)$
- 2 Bookie may then buy or sell Agent's bets on any  $A$ s
- 3 Finally, an observation is made and the bets are settled

The goal as Agent is to set your probabilities in such a way that Bookie cannot *guarantee* that they will make a profit from you.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

For reference:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

Suppose that Agent's  $\mathbb{P}(A|\Omega) \not\equiv 0$ .

How can you guarantee a profit as Bookie?

For reference:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

Suppose that Agent's  $\mathbb{P}(A|\Omega) \not\geq 0$ .

How can you guarantee a profit as Bookie?

- Bookie buys Agent's bet for  $\mathbb{P}(A|\Omega)$  dollars, a strictly negative amount,

## Intro

## Logic

## Insufficient data

## Boolean algebra

## Probability notation

## Justifications

## Betting strategy

## Subjective belief

## Objective frequency

## Many worlds

## Bayes

## Derivation

## WHy use Bayes?

## Caveat

## Expectation

## Definition

## Properties

## Invariance

## Puzzles

For reference:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

Suppose that Agent's  $\mathbb{P}(A|\Omega) \not\geq 0$ .

How can you guarantee a profit as Bookie?

- Bookie buys Agent's bet for  $\mathbb{P}(A|\Omega)$  dollars, a strictly negative amount, which means effectively Agent must pay Bookie a strictly positive amount.

## Intro

## Logic

## Insufficient data

## Boolean algebra

## Probability notation

## Justifications

## Betting strategy

## Subjective belief

## Objective frequency

## Many worlds

## Bayes

## Derivation

## WHy use Bayes?

## Caveat

## Expectation

## Definition

## Properties

## Invariance

## Puzzles

For reference:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

Suppose that Agent's  $\mathbb{P}(A|\Omega) \not\geq 0$ .

How can you guarantee a profit as Bookie?

- Bookie buys Agent's bet for  $\mathbb{P}(A|\Omega)$  dollars, a strictly negative amount, which means effectively Agent must pay Bookie a strictly positive amount.
  - If  $A$  turns out to be true, Agent must pay Bookie an additional dollar, but in either case Bookie has made a profit.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

For reference:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

Suppose that Agent's  $\mathbb{P}(\text{tautology}|\Omega) > 1$ .

How can you guarantee a profit as Bookie?

For reference:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

Suppose that Agent's  $\mathbb{P}(\text{tautology}|\Omega) > 1$ .

How can you guarantee a profit as Bookie?

- Bookie sells this bet to Agent. Agent must pay strictly more than 1 dollar up front. Since it's a tautology, Bookie must return 1 dollar, but still makes a strictly positive profit.



## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

For reference:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

Suppose that Agent's  $\mathbb{P}(\text{tautology}|\Omega) < 1$ .

How can you guarantee a profit as Bookie?

## Intro

## Logic

## Insufficient data

## Boolean algebra

## Probability notation

## Justifications

## Betting strategy

## Subjective belief

## Objective frequency

## Many worlds

## Bayes

## Derivation

## WHy use Bayes?

## Caveat

## Expectation

## Definition

## Properties

## Invariance

## Puzzles

For reference:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

Suppose that Agent's  $\mathbb{P}(\text{tautology}|\Omega) < 1$ .

How can you guarantee a profit as Bookie?

- Bookie buys Agent's bet for less than 1 dollar, and Agent must pay 1 dollar.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

For reference:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

Suppose that Agent's  $\mathbb{P}(A \vee B|\Omega) > \mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$  for disjoint  $A$  and  $B$ .  
How can you guarantee a profit as Bookie?

## Intro

## Logic

## Insufficient data

## Boolean algebra

## Probability notation

## Justifications

## Betting strategy

## Subjective belief

## Objective frequency

## Many worlds

## Bayes

## Derivation

## WHy use Bayes?

## Caveat

## Expectation

## Definition

## Properties

## Invariance

## Puzzles

For reference:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

Suppose that Agent's  $\mathbb{P}(A \vee B|\Omega) > \mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$  for disjoint  $A$  and  $B$ .  
How can you guarantee a profit as Bookie?

- Bookie buys Agent's bets for  $A$  and  $B$  at a cost of  $\mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$ .  
Bookie then sells Agent a bet for  $A \vee B$  and Agent pays  $\mathbb{P}(A \vee B|\Omega)$ .

For reference:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

Suppose that Agent's  $\mathbb{P}(A \vee B|\Omega) > \mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$  for disjoint  $A$  and  $B$ .  
How can you guarantee a profit as Bookie?

- Bookie buys Agent's bets for  $A$  and  $B$  at a cost of  $\mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$ . Bookie then sells Agent a bet for  $A \vee B$  and Agent pays  $\mathbb{P}(A \vee B|\Omega)$ .
- Bookie has made a profit, and if  $A$  and  $B$  both turn out to be false then no further money changes hands, but if either do turn out to be true then Bookie and Agent owe each other a dollar.

## Intro

## Logic

## Insufficient data

## Boolean algebra

## Probability notation

## Justifications

## Betting strategy

## Subjective belief

## Objective frequency

## Many worlds

## Bayes

## Derivation

## WHy use Bayes?

## Caveat

## Expectation

## Definition

## Properties

## Invariance

## Puzzles

For reference:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

Suppose that Agent's  $\mathbb{P}(A \vee B|\Omega) > \mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$  for disjoint  $A$  and  $B$ .  
How can you guarantee a profit as Bookie?

- Bookie buys Agent's bets for  $A$  and  $B$  at a cost of  $\mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$ . Bookie then sells Agent a bet for  $A \vee B$  and Agent pays  $\mathbb{P}(A \vee B|\Omega)$ .
- Bookie has made a profit, and if  $A$  and  $B$  both turn out to be false then no further money changes hands, but if either do turn out to be true then Bookie and Agent owe each other a dollar.
  - Why do  $A$  and  $B$  have to be disjoint?

For reference:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

Suppose that Agent's  $\mathbb{P}(A \vee B|\Omega) > \mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$  for disjoint  $A$  and  $B$ .  
How can you guarantee a profit as Bookie?

- Bookie buys Agent's bets for  $A$  and  $B$  at a cost of  $\mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$ .  
Bookie then sells Agent a bet for  $A \vee B$  and Agent pays  $\mathbb{P}(A \vee B|\Omega)$ .
- Bookie has made a profit, and if  $A$  and  $B$  both turn out to be false then no further money changes hands, but if either do turn out to be true then Bookie and Agent owe each other a dollar.
  - Why do  $A$  and  $B$  have to be disjoint?
- What if Agent's  $\mathbb{P}(A \vee B|\Omega) < \mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$ ?

For reference:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

Suppose that Agent's  $\mathbb{P}(A \vee B|\Omega) > \mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$  for disjoint  $A$  and  $B$ .  
How can you guarantee a profit as Bookie?

- Bookie buys Agent's bets for  $A$  and  $B$  at a cost of  $\mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$ . Bookie then sells Agent a bet for  $A \vee B$  and Agent pays  $\mathbb{P}(A \vee B|\Omega)$ .
- Bookie has made a profit, and if  $A$  and  $B$  both turn out to be false then no further money changes hands, but if either do turn out to be true then Bookie and Agent owe each other a dollar.
  - Why do  $A$  and  $B$  have to be disjoint?
- What if Agent's  $\mathbb{P}(A \vee B|\Omega) < \mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$ ?
  - Swap buying with selling.



For reference:

A **bet** on  $A$  with probability  $p$  is this contract: buyer pays  $\$p$  up front to seller; if  $A$  is observed to be true, seller pays  $\$1$  to buyer (else, nothing)

Suppose that Agent's  $\mathbb{P}(A \vee B|\Omega) > \mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$  for disjoint  $A$  and  $B$ . How can you guarantee a profit as Bookie?

- Bookie buys Agent's bets for  $A$  and  $B$  at a cost of  $\mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$ . Bookie then sells Agent a bet for  $A \vee B$  and Agent pays  $\mathbb{P}(A \vee B|\Omega)$ .
- Bookie has made a profit, and if  $A$  and  $B$  both turn out to be false then no further money changes hands, but if either do turn out to be true then Bookie and Agent owe each other a dollar.
  - Why do  $A$  and  $B$  have to be disjoint?
- What if Agent's  $\mathbb{P}(A \vee B|\Omega) < \mathbb{P}(A|\Omega) + \mathbb{P}(B|\Omega)$ ?
  - Swap buying with selling.
- This can be generalized to arbitrarily many disjoint events.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

Suppose that if evidence  $B$  is observed, Agent will update to a  $\mathbb{P}(A|B, \Omega) \neq \frac{\mathbb{P}(A \wedge B|\Omega)}{\mathbb{P}(B|\Omega)}$ .  
How can you guarantee a profit as Bookie?

Suppose that if evidence  $B$  is observed, Agent will update to a  $\mathbb{P}(A|B, \Omega) < \frac{\mathbb{P}(A \wedge B|\Omega)}{\mathbb{P}(B|\Omega)}$ .

How can you guarantee a profit as Bookie?

- Bet 1. Bookie sells Agent a bet for  $A \wedge B$ ; Agent pays  $\mathbb{P}(A \wedge B|\Omega)$ .
- Bet 2. Bookie sells Agent  $\frac{\mathbb{P}(A \wedge B|\Omega)}{\mathbb{P}(B|\Omega)}$  of a bet for  $\neg B$ ; Agent pays  $\mathbb{P}(A \wedge B|\Omega) \frac{\mathbb{P}(\neg B|\Omega)}{\mathbb{P}(B|\Omega)}$ .
- Bet 3. Bookie sells Agent  $\frac{\mathbb{P}(A \wedge B|\Omega)}{\mathbb{P}(B|\Omega)} - \mathbb{P}(A|B, \Omega)$  of a bet for  $B$ ; Agent pays  $\mathbb{P}(A \wedge B|\Omega) - \mathbb{P}(A|B, \Omega) \mathbb{P}(B|\Omega)$ .
- If  $B$  is false, then all the bets settle. Bookie pays back  $\frac{\mathbb{P}(A \wedge B|\Omega)}{\mathbb{P}(B|\Omega)}$  for Bet 2. Due to countable additivity, Bets 1 and 2 exactly cancel out and the price for Bet 3 remains as profit.
- If  $B$  is true, then Bets 2 and 3 are settled, with Bookie paying back  $\frac{\mathbb{P}(A \wedge B|\Omega)}{\mathbb{P}(B|\Omega)} - \mathbb{P}(A|B, \Omega)$  for Bet 3.
- Bet 4. Bookie then buys from Agent a bet for  $A$  at Agent's new probability of  $\mathbb{P}(A|B, \Omega)$ .
- At this point, whether or not  $A$  turns out to be true, Bookie and Agent owe each other a net 0 dollars, and Bookie has made a net profit.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① (Nonnegativity)  $\mathbb{P}(A|\Omega) \geq 0$
- ② (Tautology)  $\mathbb{P}(A|\Omega) = 1$  if  $A$  is a logical truth given  $\Omega$
- ③ (Countable additivity) For arbitrary disjoint  $A_i$ ,  
$$\mathbb{P}\left(\bigvee_{i=1}^{\infty} A_i \middle| \Omega\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i|\Omega)$$
- ④ (Conditional) 
$$\mathbb{P}(A|B, \Omega) = \frac{\mathbb{P}(A \wedge B|\Omega)}{\mathbb{P}(B|\Omega)}$$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

① (Bounding)  $0 = \mathbb{P}(\perp|\Omega) \leq \mathbb{P}(A|\Omega) \leq \mathbb{P}(\Omega|\Omega) = 1$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① (Bounding)  $0 = \mathbb{P}(\perp|\Omega) \leq \mathbb{P}(A|\Omega) \leq \mathbb{P}(\Omega|\Omega) = 1$
- ② (Negation)  $\mathbb{P}(A|\Omega) = 1 - \mathbb{P}(\neg A|\Omega)$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① (Bounding)  $0 = \mathbb{P}(\perp|\Omega) \leq \mathbb{P}(A|\Omega) \leq \mathbb{P}(\Omega|\Omega) = 1$
- ② (Negation)  $\mathbb{P}(A|\Omega) = 1 - \mathbb{P}(\neg A|\Omega)$
- ③ (Product rule)  $\mathbb{P}(A \wedge B|\Omega) = \mathbb{P}(A|B, \Omega) \cdot \mathbb{P}(B|\Omega)$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① (Bounding)  $0 = \mathbb{P}(\perp|\Omega) \leq \mathbb{P}(A|\Omega) \leq \mathbb{P}(\Omega|\Omega) = 1$
- ② (Negation)  $\mathbb{P}(A|\Omega) = 1 - \mathbb{P}(\neg A|\Omega)$
- ③ (Product rule)  $\mathbb{P}(A \wedge B|\Omega) = \mathbb{P}(A|B, \Omega) \cdot \mathbb{P}(B|\Omega)$
- ④ (Countable additivity) For arbitrary disjoint  $A_i$ ,  
$$\mathbb{P}\left(\bigvee_{i=1}^{\infty} A_i \middle| \Omega\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i|\Omega)$$



## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- 1  $\mathbb{P}(A|B)$  is a real number, and  $A$  and  $B$  are elements of a Boolean algebra

## Intro

Logic  
Insufficient data  
Boolean algebra  
Probability notation

## Justifications

Betting strategy  
Subjective belief  
Objective frequency  
Many worlds

## Bayes

Derivation  
WHy use Bayes?  
Caveat

## Expectation

Definition  
Properties

## Invariance

## Puzzles

- ①  $\mathbb{P}(A|B)$  is a real number, and  $A$  and  $B$  are elements of a Boolean algebra
- Technically, a complete Boolean algebra

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ①  $\mathbb{P}(A|B)$  is a real number, and  $A$  and  $B$  are elements of a Boolean algebra
- Technically, a complete Boolean algebra, which essentially means that countably infinite disjunctions (and/or conjunctions) are allowed.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ①  $\mathbb{P}(A|B)$  is a real number, and  $A$  and  $B$  are elements of a Boolean algebra
- Technically, a complete Boolean algebra, which essentially means that countably infinite disjunctions (and/or conjunctions) are allowed.
  - Any (incomplete) Boolean algebra (or any topology!) generates a canonical complete Boolean algebra.

## Intro

## Logic

## Insufficient data

## Boolean algebra

## Probability notation

## Justifications

## Betting strategy

## Subjective belief

## Objective frequency

## Many worlds

## Bayes

## Derivation

## WHy use Bayes?

## Caveat

## Expectation

## Definition

## Properties

## Invariance

## Puzzles

- ①  $\mathbb{P}(A|B)$  is a real number, and  $A$  and  $B$  are elements of a Boolean algebra
  - Technically, a complete Boolean algebra, which essentially means that countably infinite disjunctions (and/or conjunctions) are allowed.
  - Any (incomplete) Boolean algebra (or any topology!) generates a canonical complete Boolean algebra.
- ②  $A_1 \subseteq A_2 \subseteq A_3 \subseteq \cdots$  such that  $A_i \nearrow A$  implies  $\mathbb{P}(A_i|\Omega) \nearrow \mathbb{P}(A|\Omega)$ .

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ①  $\mathbb{P}(A|B)$  is a real number, and  $A$  and  $B$  are elements of a Boolean algebra
  - Technically, a complete Boolean algebra, which essentially means that countably infinite disjunctions (and/or conjunctions) are allowed.
  - Any (incomplete) Boolean algebra (or any topology!) generates a canonical complete Boolean algebra.
- ②  $A_1 \subseteq A_2 \subseteq A_3 \subseteq \cdots$  such that  $A_i \nearrow A$  implies  $\mathbb{P}(A_i|\Omega) \nearrow \mathbb{P}(A|\Omega)$ .
- ③  $\mathbb{P}(A \wedge B|\Omega) = F[\mathbb{P}(A|B, \Omega), \mathbb{P}(B|\Omega)]$  for some  $F$ .

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ①  $\mathbb{P}(A|B)$  is a real number, and  $A$  and  $B$  are elements of a Boolean algebra
  - Technically, a complete Boolean algebra, which essentially means that countably infinite disjunctions (and/or conjunctions) are allowed.
  - Any (incomplete) Boolean algebra (or any topology!) generates a canonical complete Boolean algebra.
- ②  $A_1 \subseteq A_2 \subseteq A_3 \subseteq \dots$  such that  $A_i \nearrow A$  implies  $\mathbb{P}(A_i|\Omega) \nearrow \mathbb{P}(A|\Omega)$ .
- ③  $\mathbb{P}(A \wedge B|\Omega) = F[\mathbb{P}(A|B, \Omega), \mathbb{P}(B|\Omega)]$  for some  $F$ .
- ④  $\mathbb{P}(\neg A|\Omega) = N[\mathbb{P}(A|\Omega)]$  for some  $N$ .

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Cox “proved” something like this in 1946, known as Cox’s Theorem.



## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Cox “proved” something like this in 1946, known as Cox’s Theorem.
- It turns out Cox’s Theorem is wrong!

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Cox “proved” something like this in 1946, known as Cox’s Theorem.
- It turns out Cox’s Theorem is wrong!
  - Factoid of the hour: “Cox’s Theorem is false; Halpern in 1999 provided a counterexample”

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Cox “proved” something like this in 1946, known as Cox’s Theorem.
- It turns out Cox’s Theorem is wrong!
  - Factoid of the hour: “Cox’s Theorem is false; Halpern in 1999 provided a counterexample”
  - The version on Wikipedia is also wrong

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Cox “proved” something like this in 1946, known as Cox’s Theorem.
- It turns out Cox’s Theorem is wrong!
  - Factoid of the hour: “Cox’s Theorem is false; Halpern in 1999 provided a counterexample”
  - The version on Wikipedia is also wrong
- But the laws of probability I showed here *do* follow from the premises I showed here, in a modern proof by Terenin and Draper

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① (Bounding)  $0 = \mathbb{P}(\perp|\Omega) \leq \mathbb{P}(A|\Omega) \leq \mathbb{P}(\Omega|\Omega) = 1$
- ② (Negation)  $\mathbb{P}(A|\Omega) = 1 - \mathbb{P}(\neg A|\Omega)$
- ③ (Product rule)  $\mathbb{P}(A \wedge B|\Omega) = \mathbb{P}(A|B, \Omega) \cdot \mathbb{P}(B|\Omega)$
- ④ (Countable additivity) For arbitrary disjoint  $A_i$ ,  
$$\mathbb{P}\left(\bigvee_{i=1}^{\infty} A_i \middle| \Omega\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i|\Omega)$$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

1 (Nonnegativity)  $\mathbb{P}(A) \geq 0$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

① (Nonnegativity)  $\mathbb{P}(A) \geq 0$

② (Normalization)  $\mathbb{P}(\Omega) = 1$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① (Nonnegativity)  $\mathbb{P}(A) \geq 0$
- ② (Normalization)  $\mathbb{P}(\Omega) = 1$
- ③ (Countable additivity) For arbitrary disjoint  $A_i$ ,  
$$\mathbb{P}\left(\bigvee_{i=1}^{\infty} A_i \middle| \Omega\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i | \Omega)$$



## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① (Nonnegativity)  $\mathbb{P}(A) \geq 0$
- ② (Normalization)  $\mathbb{P}(\Omega) = 1$
- ③ (Countable additivity) For arbitrary disjoint  $A_i$ ,  
$$\mathbb{P}\left(\bigvee_{i=1}^{\infty} A_i \middle| \Omega\right) B = \sum_{i=1}^{\infty} \mathbb{P}(A_i | \Omega)$$
- ④ (Conditional)  $\mathbb{P}(A|B) = \frac{\mathbb{P}(A \wedge B)}{\mathbb{P}(B)}$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Consider set of outcomes of (infinitely) repeated independent experiments

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Consider set of outcomes of (infinitely) repeated independent experiments
- “Probability”  $\mathbb{P}(A)$  is defined as the ratio  $N_A/N$  of outcomes where  $A$  is true ( $N_A$ ) to total number of experiments ( $N$ ).

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Nonnegativity

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Nonnegativity
  - Both the total number of experiments, and the number of outcomes in which the proposition is true, cannot be negative, therefore their ratio cannot be negative.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Nonnegativity
  - Both the total number of experiments, and the number of outcomes in which the proposition is true, cannot be negative, therefore their ratio cannot be negative.
- Normalization

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Nonnegativity
  - Both the total number of experiments, and the number of outcomes in which the proposition is true, cannot be negative, therefore their ratio cannot be negative.
- Normalization
  - The number of experiments in which an experiment was performed is identical to the total number of experiments, so their ratio is 1.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Nonnegativity
  - Both the total number of experiments, and the number of outcomes in which the proposition is true, cannot be negative, therefore their ratio cannot be negative.
- Normalization
  - The number of experiments in which an experiment was performed is identical to the total number of experiments, so their ratio is 1.
- Countable additivity



## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Nonnegativity
  - Both the total number of experiments, and the number of outcomes in which the proposition is true, cannot be negative, therefore their ratio cannot be negative.
- Normalization
  - The number of experiments in which an experiment was performed is identical to the total number of experiments, so their ratio is 1.
- Countable additivity
  - The number of outcomes in which one of many mutually exclusive alternatives is true is precisely the number of outcomes in which at least one is true.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Conditional

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Conditional
  - We can justify this by the principle of rejection:

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Conditional

- We can justify this by the principle of rejection: to perform an experiment “conditional” on the knowledge of  $B$  is simply to reject (exclude from consideration) every experiment in which  $B$  is false.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Conditional

- We can justify this by the principle of rejection: to perform an experiment “conditional” on the knowledge of  $B$  is simply to reject (exclude from consideration) every experiment in which  $B$  is false.
- Thus, the (effective) total number of experiments is the number of experiments in which  $B$  is true, and the (effective) total number of experiments in which  $A$  is true is those in which both  $A$  and  $B$  are true.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Conditional

- We can justify this by the principle of rejection: to perform an experiment “conditional” on the knowledge of  $B$  is simply to reject (exclude from consideration) every experiment in which  $B$  is false.
- Thus, the (effective) total number of experiments is the number of experiments in which  $B$  is true, and the (effective) total number of experiments in which  $A$  is true is those in which both  $A$  and  $B$  are true.
- So

$$\mathbb{P}(A|B) = \frac{N_{A \wedge B}}{N_B} = \frac{N_{A \wedge B}/N}{N_B/N} = \frac{\mathbb{P}(A \wedge B)}{\mathbb{P}(B)}$$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Instead of considering a sequence of independent experiments, we can just as well consider an un-ordered set of possible worlds.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Instead of considering a sequence of independent experiments, we can just as well consider an un-ordered set of possible worlds.
- The same arguments as for objective frequency go through.



## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Instead of considering a sequence of independent experiments, we can just as well consider an un-ordered set of possible worlds.
- The same arguments as for objective frequency go through.
- You may prefer one or the other philosophically.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Instead of considering a sequence of independent experiments, we can just as well consider an un-ordered set of possible worlds.
- The same arguments as for objective frequency go through.
- You may prefer one or the other philosophically.
- Bernoulli even used both: one layer of uncertainty due to ignorance, and one due to randomness.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- Instead of considering a sequence of independent experiments, we can just as well consider an un-ordered set of possible worlds.
- The same arguments as for objective frequency go through.
- You may prefer one or the other philosophically.
- Bernoulli even used both: one layer of uncertainty due to ignorance, and one due to randomness.
- Formally, they behave the same and blend together.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- If we take the objective-frequency or many-worlds interpretations, we call the experimental outcomes or possible worlds as elements of a **sample space**, which is notated  $\Omega$  and is an object of set theory.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- If we take the objective-frequency or many-worlds interpretations, we call the experimental outcomes or possible worlds as elements of a **sample space**, which is notated  $\Omega$  and is an object of set theory.
- The subjective-belief and betting-strategy frameworks are sufficiently abstract that they don't come with a concrete sample space, but a canonical one can be constructed (via the Loomis-Sikorski representation theorem).

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

Using the **Product Rule**:  $\mathbb{P}(A \wedge B|\Omega) = \mathbb{P}(A|B, \Omega) \cdot \mathbb{P}(B|\Omega)$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

Using the **Product Rule**:  $\mathbb{P}(A \wedge B|\Omega) = \mathbb{P}(A|B, \Omega) \cdot \mathbb{P}(B|\Omega)$

$$\mathbb{P}(A|B, \Omega) \cdot \mathbb{P}(B|\Omega) = P(A \wedge B) = P(B \wedge A) = \mathbb{P}(B|A, \Omega) \cdot \mathbb{P}(A|\Omega)$$

$$\mathbb{P}(A|B, \Omega) \cdot \mathbb{P}(B|\Omega) = \mathbb{P}(B|A, \Omega) \cdot \mathbb{P}(A|\Omega)$$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

Using the **Product Rule**:  $\mathbb{P}(A \wedge B|\Omega) = \mathbb{P}(A|B, \Omega) \cdot \mathbb{P}(B|\Omega)$

$$\mathbb{P}(A|B, \Omega) \cdot \mathbb{P}(B|\Omega) = P(A \wedge B) = P(B \wedge A) = \mathbb{P}(B|A, \Omega) \cdot \mathbb{P}(A|\Omega)$$

$$\mathbb{P}(A|B, \Omega) \cdot \mathbb{P}(B|\Omega) = \mathbb{P}(B|A, \Omega) \cdot \mathbb{P}(A|\Omega)$$

$$\text{Bayes: } \mathbb{P}(A|B, \Omega) = \frac{\mathbb{P}(B|A, \Omega) \cdot \mathbb{P}(A|\Omega)}{\mathbb{P}(B|\Omega)}$$



## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

Using the **Product Rule**:  $\mathbb{P}(A \wedge B|\Omega) = \mathbb{P}(A|B, \Omega) \cdot \mathbb{P}(B|\Omega)$

$$\mathbb{P}(A|B, \Omega) \cdot \mathbb{P}(B|\Omega) = P(A \wedge B) = P(B \wedge A) = \mathbb{P}(B|A, \Omega) \cdot \mathbb{P}(A|\Omega)$$

$$\mathbb{P}(A|B, \Omega) \cdot \mathbb{P}(B|\Omega) = \mathbb{P}(B|A, \Omega) \cdot \mathbb{P}(A|\Omega)$$

$$\text{Bayes: } \mathbb{P}(A|B, \Omega) = \frac{\mathbb{P}(B|A, \Omega) \cdot \mathbb{P}(A|\Omega)}{\mathbb{P}(B|\Omega)}$$

Why is this useful?

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

Suppose you're a geologist examining a rock. You can measure its density but want to know its composition.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

Suppose you're a geologist examining a rock. You can measure its density but want to know its composition. Let's set propositions

$A$  = the rock contains iron.

$B$  = the rock's density is 3,160 kg/m<sup>3</sup>.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

Suppose you're a geologist examining a rock. You can measure its density but want to know its composition. Let's set propositions

$A$  = the rock contains iron.

$B$  = the rock's density is  $3,160 \text{ kg/m}^3$ .

We say that  $\mathbb{P}(A|B, \Omega)$  is a **discriminative model** (which is usually what you want), and  $\mathbb{P}(B|A, \Omega)$  is a **generative model** (which is usually easier to make scientifically).

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

Suppose you're a geologist examining a rock. You can measure its density but want to know its composition. Let's set propositions

$A$  = the rock contains iron.

$B$  = the rock's density is 3,160 kg/m<sup>3</sup>.

We say that  $\mathbb{P}(A|B, \Omega)$  is a **discriminative model** (which is usually what you want), and  $\mathbb{P}(B|A, \Omega)$  is a **generative model** (which is usually easier to make scientifically).

Bayes lets you convert between the two.

$$\text{Bayes: } \mathbb{P}(A|B, \Omega) = \frac{\mathbb{P}(B|A, \Omega) \cdot \mathbb{P}(A|\Omega)}{\mathbb{P}(B|\Omega)}$$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

 $A$  = the rock contains iron. $B$  = the rock's density is 3,160 kg/m<sup>3</sup>.

$$\text{Bayes: } \mathbb{P}(A|B, \Omega) = \frac{\mathbb{P}(B|A, \Omega) \cdot \mathbb{P}(A|\Omega)}{\mathbb{P}(B|\Omega)}$$

In the cases where we want to use Bayes,  $\mathbb{P}(B|\Omega)$  is usually impossible to compute, because one must consider *all possible hypotheses* for how  $B$  might be true, and we're bounded agents.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

$A$  = the rock contains iron.

$B$  = the rock's density is 3,160 kg/m<sup>3</sup>.

$$\text{Bayes: } \mathbb{P}(A|B, \Omega) = \frac{\mathbb{P}(B|A, \Omega) \cdot \mathbb{P}(A|\Omega)}{\mathbb{P}(B|\Omega)}$$

In the cases where we want to use Bayes,  $\mathbb{P}(B|\Omega)$  is usually impossible to compute, because one must consider *all possible hypotheses* for how  $B$  might be true, and we're bounded agents. Instead, we generally use Bayes to *compare* hypotheses.

$$\frac{P(A_1|B)}{P(A_2|B)} = \frac{P(B|A_1) \cdot P(A_1)/P(B)}{P(B|A_2) \cdot P(A_2)/P(B)}$$

$$\frac{P(A_1|B)}{P(A_2|B)} = \frac{P(B|A_1)}{P(B|A_2)} \cdot \frac{P(A_1)}{P(A_2)}$$

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

$A$  = the rock contains iron.

$B$  = the rock's density is 3,160 kg/m<sup>3</sup>.

$$\text{Bayes: } \mathbb{P}(A|B, \Omega) = \frac{\mathbb{P}(B|A, \Omega) \cdot \mathbb{P}(A|\Omega)}{\mathbb{P}(B|\Omega)}$$

In the cases where we want to use Bayes,  $\mathbb{P}(B|\Omega)$  is usually impossible to compute, because one must consider *all possible hypotheses* for how  $B$  might be true, and we're bounded agents. Instead, we generally use Bayes to *compare* hypotheses.

$$\frac{P(A_1|B)}{P(A_2|B)} = \frac{P(B|A_1) \cdot P(A_1)/P(B)}{P(B|A_2) \cdot P(A_2)/P(B)}$$

$$\frac{P(A_1|B)}{P(A_2|B)} = \frac{P(B|A_1)}{P(B|A_2)} \cdot \frac{P(A_1)}{P(A_2)}$$

Generating good candidate hypotheses is much of the difficulty in most real inference problems.



## Intro

Logic  
Insufficient data  
Boolean algebra  
Probability notation

## Justifications

Betting strategy  
Subjective belief  
Objective frequency  
Many worlds

## Bayes

Derivation  
WHy use Bayes?  
Caveat

## Expectation

Definition  
Properties

## Invariance

## Puzzles

- If  $V$  is a real-valued function defined on the sample space, the **expected value**  $\mathbb{E}(V)$  is defined as the Lebesgue integral  $\int_{\Omega} V(x) d\mathbb{P}(x|\Omega)$ .

## Intro

Logic  
Insufficient data  
Boolean algebra  
Probability notation

## Justifications

Betting strategy  
Subjective belief  
Objective frequency  
Many worlds

## Bayes

Derivation  
WHy use Bayes?  
Caveat

## Expectation

Definition  
Properties

## Invariance

## Puzzles

- If  $V$  is a real-valued function defined on the sample space, the **expected value**  $\mathbb{E}(V)$  is defined as the Lebesgue integral  $\int_{\Omega} V(x) d\mathbb{P}(x|\Omega)$ .
- In cases where a discrete summation  $\sum_{x \in \Omega} V(x) \cdot \mathbb{P}(x|\Omega)$  applies, this agrees with the Lebesgue integral.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- If  $V$  is a real-valued function defined on the sample space, the **expected value**  $\mathbb{E}(V)$  is defined as the Lebesgue integral  $\int_{\Omega} V(x) d\mathbb{P}(x|\Omega)$ .
- In cases where a discrete summation  $\sum_{x \in \Omega} V(x) \cdot \mathbb{P}(x|\Omega)$  applies, this agrees with the Lebesgue integral.
  - In such a case,  $\mathbb{P}(x|\Omega)$  is known as a **probability mass function**.

## Intro

Logic  
Insufficient data  
Boolean algebra  
Probability notation

## Justifications

Betting strategy  
Subjective belief  
Objective frequency  
Many worlds

## Bayes

Derivation  
WHy use Bayes?  
Caveat

## Expectation

Definition  
Properties

## Invariance

## Puzzles

- If  $V$  is a real-valued function defined on the sample space, the **expected value**  $\mathbb{E}(V)$  is defined as the Lebesgue integral  $\int_{\Omega} V(x) d\mathbb{P}(x|\Omega)$ .
- In cases where a discrete summation  $\sum_{x \in \Omega} V(x) \cdot \mathbb{P}(x|\Omega)$  applies, this agrees with the Lebesgue integral.
  - In such a case,  $\mathbb{P}(x|\Omega)$  is known as a **probability mass function**.
- Also, in cases where a Riemann integral  $\int_{\Omega} V(x) \cdot \text{pdf}(x) dx$  applies (i.e. where  $\Omega \subseteq \mathbb{R}^n$ ), this agrees with the Lebesgue integral as well.

## Intro

Logic  
Insufficient data  
Boolean algebra  
Probability notation

## Justifications

Betting strategy  
Subjective belief  
Objective frequency  
Many worlds

## Bayes

Derivation  
WHy use Bayes?  
Caveat

## Expectation

Definition  
Properties

## Invariance

## Puzzles

- If  $V$  is a real-valued function defined on the sample space, the **expected value**  $\mathbb{E}(V)$  is defined as the Lebesgue integral  $\int_{\Omega} V(x) d\mathbb{P}(x|\Omega)$ .
- In cases where a discrete summation  $\sum_{x \in \Omega} V(x) \cdot \mathbb{P}(x|\Omega)$  applies, this agrees with the Lebesgue integral.
  - In such a case,  $\mathbb{P}(x|\Omega)$  is known as a **probability mass function**.
- Also, in cases where a Riemann integral  $\int_{\Omega} V(x) \cdot \text{pdf}(x) dx$  applies (i.e. where  $\Omega \subseteq \mathbb{R}^n$ ), this agrees with the Lebesgue integral as well.
  - $\text{pdf}(x) = \frac{d\mathbb{P}(\omega \leq x|\Omega)}{dx}$  is known as a **probability density function**.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- If  $V$  is a real-valued function defined on the sample space, the **expected value**  $\mathbb{E}(V)$  is defined as the Lebesgue integral  $\int_{\Omega} V(x) d\mathbb{P}(x|\Omega)$ .
- In cases where a discrete summation  $\sum_{x \in \Omega} V(x) \cdot \mathbb{P}(x|\Omega)$  applies, this agrees with the Lebesgue integral.
  - In such a case,  $\mathbb{P}(x|\Omega)$  is known as a **probability mass function**.
- Also, in cases where a Riemann integral  $\int_{\Omega} V(x) \cdot \text{pdf}(x) dx$  applies (i.e. where  $\Omega \subseteq \mathbb{R}^n$ ), this agrees with the Lebesgue integral as well.
  - $\text{pdf}(x) = \frac{d\mathbb{P}(\omega \leq x|\Omega)}{dx}$  is known as a **probability density function**.
  - $\mathbb{P}(\omega \leq x|\Omega)$  itself, also a function of  $x$ , is known as a **cumulative distribution function** (or simply **distribution function**), and uniquely specifies the probability of all events depending on  $\omega$ .

## Intro

Logic  
Insufficient data  
Boolean algebra  
Probability notation

## Justifications

Betting strategy  
Subjective belief  
Objective frequency  
Many worlds

## Bayes

Derivation  
WHy use Bayes?  
Caveat

## Expectation

Definition  
Properties

## Invariance

## Puzzles

- The expected value of an indicator variable  $1_A$  (defined to be 1 if  $A$  is true and 0 otherwise) is  $\mathbb{E}(1_A) = \mathbb{P}(A|\Omega)$ .

## Intro

Logic  
Insufficient data  
Boolean algebra  
Probability notation

## Justifications

Betting strategy  
Subjective belief  
Objective frequency  
Many worlds

## Bayes

Derivation  
WHy use Bayes?  
Caveat

## Expectation

Definition  
Properties

## Invariance

## Puzzles

- The expected value of an indicator variable  $1_A$  (defined to be 1 if  $A$  is true and 0 otherwise) is  $\mathbb{E}(1_A) = \mathbb{P}(A|\Omega)$ .
- The most important property of expectation is linearity:

$$\mathbb{E}(X + Y) = \mathbb{E}(X) + \mathbb{E}(Y)$$

$$\mathbb{E}(a \cdot X) = a \cdot \mathbb{E}(X)$$



## Intro

Logic  
Insufficient data  
Boolean algebra  
Probability notation

## Justifications

Betting strategy  
Subjective belief  
Objective frequency  
Many worlds

## Bayes

Derivation  
WHy use Bayes?  
Caveat

## Expectation

Definition  
Properties

## Invariance

## Puzzles

- The expected value of an indicator variable  $1_A$  (defined to be 1 if  $A$  is true and 0 otherwise) is  $\mathbb{E}(1_A) = \mathbb{P}(A|\Omega)$ .
- The most important property of expectation is linearity:

$$\mathbb{E}(X + Y) = \mathbb{E}(X) + \mathbb{E}(Y)$$

$$\mathbb{E}(a \cdot X) = a \cdot \mathbb{E}(X)$$

These follow from the linearity of the Lebesgue integral.

## Intro

Logic  
Insufficient data  
Boolean algebra  
Probability notation

## Justifications

Betting strategy  
Subjective belief  
Objective frequency  
Many worlds

## Bayes

Derivation  
WHy use Bayes?  
Caveat

## Expectation

Definition  
Properties

## Invariance

## Puzzles

- If you can assume probability mass functions are invariant under arbitrary permutations of a finite set of events, then the probabilities of those events are **uniform**, i.e. equal.

## Intro

Logic  
Insufficient data  
Boolean algebra  
Probability notation

## Justifications

Betting strategy  
Subjective belief  
Objective frequency  
Many worlds

## Bayes

Derivation  
WHy use Bayes?  
Caveat

## Expectation

Definition  
Properties

## Invariance

## Puzzles

- If you can assume probability mass functions are invariant under arbitrary permutations of a finite set of events, then the probabilities of those events are **uniform**, i.e. equal.
- From countable additivity, they must equal  $1/N$  (where  $N$  is the number of events that can be permuted in such a way).

## Intro

Logic  
Insufficient data  
Boolean algebra  
Probability notation

## Justifications

Betting strategy  
Subjective belief  
Objective frequency  
Many worlds

## Bayes

Derivation  
WHy use Bayes?  
Caveat

## Expectation

Definition  
Properties

## Invariance

## Puzzles

- If you can assume probability mass functions are invariant under arbitrary permutations of a finite set of events, then the probabilities of those events are **uniform**, i.e. equal.
- From countable additivity, they must equal  $1/N$  (where  $N$  is the number of events that can be permuted in such a way).
  - Thus we obtain probability  $1/2$  for a “fair” (symmetric) coin landing on either side,  $1/6$  for a “fair” (symmetric) cubical die landing on any face, etc.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- If you can assume probability mass functions are invariant under arbitrary permutations of a finite set of events, then the probabilities of those events are **uniform**, i.e. equal.
- From countable additivity, they must equal  $1/N$  (where  $N$  is the number of events that can be permuted in such a way).
  - Thus we obtain probability  $1/2$  for a “fair” (symmetric) coin landing on either side,  $1/6$  for a “fair” (symmetric) cubical die landing on any face, etc.
- Similar arguments can be applied to density functions in continuous cases, usually to justify uniform density over an interval (scaled so that the integral of the density is 1, as required by the laws of probability).

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- 1 Are conditional probabilities probabilities of implications?
  - Either prove that  $\mathbb{P}(B|A, \Omega) = \mathbb{P}(A \rightarrow B|\Omega)$ , or find a counterexample.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① Are conditional probabilities probabilities of implications?
  - Either prove that  $\mathbb{P}(B|A, \Omega) = \mathbb{P}(A \rightarrow B|\Omega)$ , or find a counterexample.
- ② From each system of "laws of probability", prove all the laws of all the other systems.
  - Note: there's a lot of overlapping laws, which are trivially provable from each other.

## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① Are conditional probabilities probabilities of implications?
  - Either prove that  $\mathbb{P}(B|A, \Omega) = \mathbb{P}(A \rightarrow B|\Omega)$ , or find a counterexample.
- ② From each system of "laws of probability", prove all the laws of all the other systems.
  - Note: there's a lot of overlapping laws, which are trivially provable from each other.
- ③ A meter-stick is cut at a uniformly sampled point along its length. What is the expected value of the larger piece?



## Intro

Logic

Insufficient data

Boolean algebra

Probability notation

## Justifications

Betting strategy

Subjective belief

Objective frequency

Many worlds

## Bayes

Derivation

WHy use Bayes?

Caveat

## Expectation

Definition

Properties

## Invariance

## Puzzles

- ① Are conditional probabilities probabilities of implications?
  - Either prove that  $\mathbb{P}(B|A, \Omega) = \mathbb{P}(A \rightarrow B|\Omega)$ , or find a counterexample.
- ② From each system of "laws of probability", prove all the laws of all the other systems.
  - Note: there's a lot of overlapping laws, which are trivially provable from each other.
- ③ A meter-stick is cut at a uniformly sampled point along its length. What is the expected value of the larger piece?
  - Bonus puzzle: if the stick is instead cut simultaneously at two uniformly sampled points, what is the expected value of the largest piece?