

Teilmenge



irreflexiv $A \not\subseteq A$

reflexiv $A \subseteq A$

$\subseteq$; lösle $\{ \} + \in$

$$M = \{ \{1\} ; 2 ; \{1,2\} ; \{\{3\}\} \}$$

$\{1\} \subset M$	$\Rightarrow$	$1 \in M$	$\{$ falsch
$\{2\} \subset M$	$\Rightarrow$	$2 \in M$	$\checkmark$
$\{\{1\}\} \subset M$	$\Rightarrow$	$\{1\} \in M$	$\checkmark$

$$L \cap \overline{N \cap M}$$

$$L \cap \overline{\{4,5,9\} \cap \{1,2,4,7\}}$$

$$L \cap \overline{\{4\}} \rightarrow \Omega \setminus \{4\}$$

$$\underline{\{1,2,4,7\}} \cap \underline{\{1,2,3,5,6,7,8,9\}}$$

$$\{1,2,7\}$$

de Morgan

$$L \cap (\overline{N} \cup \overline{M}) \left. \begin{array}{l} \text{dopp} \\ \text{Nega.} \end{array} \right\}$$

$$L \cap (\overline{N} \cup M)$$

$$L \cap (\{1,2,3,6,7,8\} \cup \{3,5,6,8,9\})$$

$$L \cap \{1,2,3,5,6,7,8,9\}$$

$$\{1,2,7\}$$

$$\mu_4 \cap (\mu_2 \setminus \mu_3)$$

↓

$$[-1; 2[\cap (\mathbb{N} \setminus \{-1, 1, 2\})$$

$$\{0; 1\} \overset{\substack{\uparrow \\ \text{clé: } \epsilon_2}}{\cap} \underbrace{(\mathbb{N} \setminus \{1, 2\})}_{\{0; 3; 4; \dots\}} = \{0\}$$

$$\mu_3 \setminus (\mu_1 \setminus \mu_2)$$

$$\{-1, 1, 2\} \setminus (\mathbb{Z} \setminus \underbrace{\mathbb{N}}_{\mathbb{Z}_0^+})$$

$$\{-1, 1, 2\} \setminus \mathbb{Z}^- = \{1, 2\}$$