THE MASTERMIND COACHING CLASSES

CBSE / RBSE • NURSERY – CLASS XII • JAIPUR

Road No. 5, VKIA, Jaipur — 9887258679

ANSWER KEY & MARKING SCHEME SET 1

Chapter 4: **Quadratic Equations** | Class 10 Maths | Maximum Marks: 40

SECTION — A (Objective Type) — Answers — 10 × 1 = 10

Q.	Correct Option	Brief Reasoning
1	**(a)** $x^2 + 3x + 2 = 0$	Degree must be 2
2	**(d)** Yes	LHS $= 0$ at $x = 2$
3	**(b)** -3	Compare with $ax^2 + bx + c = 0$
4	**(a)** 2 and 3	Put each factor equal to zero
5	**(c)** 1	$D = b^2 - 4ac$
6	**(a)** two distinct real roots	$D = b^2 - 4ac = 1$
7	**(a)** $x^2 - 5x + 6 = 0$	Degree must be 2
8	**(d)** No	LHS $= 2$ at $x = 4$
9	**(d)** 7	Compare with $ax^2 + bx + c = 0$
10	**(a)** -1 and 4	Put each factor equal to zero

SECTION — B (Very Short Answer) — Answers — 4 × 2 = 8

Q.	Expected Answer / Value Points	Marks
11	Put $x = 2$: LHS $= 0$ (1); equals RHS 0, so it is a root (1).	1+1
12	$x - 4 = 0$ or $x - 5 = 0$ (1); $x = 4$ or $x = 5$ (1).	1+1
13	$D = (-6)^2 - 4(1)(8) = 4$ (1); two distinct real roots (1).	1+1
14	Put $x = 3$: LHS $= -2$ (1); not equal to 0, so it is not a root (1).	1+1

SECTION — C (Short Answer) — Answers — 4 × 3 = 12

Q.	Expected Answer / Value Points	Marks
15	Split the middle term: $-5 = -3 - 2$ because $ac = 6$ (1); regroup and factorise (1); roots $x = 2, 3$ (1).	1+1+1
16	$a = 2$, $b = -4$, $c = 3$ (1); $D = -8$ (1); no real roots (1).	1+1+1
17	Let integers be x and $x + 1$: $x(x + 1) = 20$ (1); $x^2 + x - 20 = 0$ (1); $(x - 4)(x + 5) = 0$, $x = 4$ (1); integers 4 and 5 (1).	1+1+1+1
	OR: Split the middle term: $-4 = -5 + 1$ because $ac = -5$ (1); regroup and factorise (1); roots $x = -1, 5$ (1).	1+1+1
18	$a = 1$, $b = -8$, $c = 16$ (1); $D = 0$ (1); two equal real roots (1).	1+1+1
	OR: Let integers be x and $x + 1$: $x(x + 1) = 42$ (1); $x^2 + x - 42 = 0$ (1); $(x - 6)(x + 7) = 0$, $x = 6$ (1); integers 6 and 7 (1).	1+1+1+1

SECTION — D (Long Answer) — Answers — 2 × 5 = 10

Q.	Expected Answer / Value Points	Marks
19	Split the middle term: $-7 = -6 - 1$ because $ac = 6$ (1); regroup and factorise (1); roots $x = \frac{1}{2}, 3$ (1).	1+1+1
	OR: Breadth x, length $x + 5$: $x(x + 5) = 66$ (1); $x^2 + 5x - 66 = 0$ (1); $(x - 6)(x + 11) = 0$ (1); $x = 6$ m (breadth), length 11 m (reject negative) (2).	1+1+1+2
20	Split the middle term: $-1 = -3 + 2$ because $ac = -6$ (1); regroup and factorise (1); roots $x = -\frac{2}{3}, 1$ (1).	1+1+1
	OR: Breadth x, length $x + 4$: $x(x + 4) = 96$ (1); $x^2 + 4x - 96 = 0$ (1); $(x - 8)(x + 12) = 0$ (1); $x = 8$ m (breadth), length 12 m (reject negative) (2).	1+1+1+2

Evaluator's note: Award step marks as shown; accept any correct alternative method. In internal-choice questions, evaluate only the option attempted. Deduct up to ½ mark for missing units or figures where required.

THE MASTERMIND COACHING CLASSES

CBSE / RBSE • NURSERY – CLASS XII • JAIPUR

Road No. 5, VKIA, Jaipur — 9887258679

ANSWER KEY & MARKING SCHEME SET 2

Chapter 4: **Quadratic Equations** | Class 10 Maths | Maximum Marks: 40

SECTION — A (Objective Type) — Answers — 10 × 1 = 10

Q.	Correct Option	Brief Reasoning
1	**(a)** 1	$D = b^2 - 4ac$
2	**(d)** two equal real roots	$D = b^2 - 4ac = 0$
3	**(b)** $x^2 + 4x + 1 = 0$	Degree must be 2
4	**(d)** Yes	LHS $= 0$ at $x = 2$
5	**(b)** -4	Compare with $ax^2 + bx + c = 0$
6	**(a)** 3 and 5	Put each factor equal to zero
7	**(b)** 0	$D = b^2 - 4ac$
8	**(a)** no real roots	$D = b^2 - 4ac = -11$
9	**(c)** $x^2 + 7x - 2 = 0$	Degree must be 2
10	**(c)** Yes	LHS $= 0$ at $x = -1$

SECTION — B (Very Short Answer) — Answers — 4 × 2 = 8

Q.	Expected Answer / Value Points	Marks
11	$x + 3 = 0$ or $x + 4 = 0$ (1); $x = -3$ or $x = -4$ (1).	1+1
12	$D = (-4)^2 - 4(1)(4) = 0$ (1); two equal real roots (1).	1+1
13	Put $x = 1$: LHS $= 0$ (1); equals RHS 0, so it is a root (1).	1+1
14	$x - 2 = 0$ or $x + 5 = 0$ (1); $x = 2$ or $x = -5$ (1).	1+1

SECTION — C (Short Answer) — Answers — 4 × 3 = 12

Q.	Expected Answer / Value Points	Marks
15	Split the middle term: $-10 = -6 - 4$ because $ac = 24$ (1); regroup and factorise (1); roots $x = 4, 6$ (1).	1+1+1
16	$a = 3$, $b = 5$, $c = -2$ (1); $D = 49$ (1); two distinct real roots (1).	1+1+1
17	Let integers be x and $x + 1$: $x(x + 1) = 72$ (1); $x^2 + x - 72 = 0$ (1); $(x - 8)(x + 9) = 0$, $x = 8$ (1); integers 8 and 9 (1).	1+1+1+1
	OR: Split the middle term: $-4 = -7 + 3$ because $ac = -21$ (1); regroup and factorise (1); roots $x = -3, 7$ (1).	1+1+1
18	$a = 1$, $b = 6$, $c = 10$ (1); $D = -4$ (1); no real roots (1).	1+1+1
	OR: Let integers be x and $x + 1$: $x(x + 1) = 132$ (1); $x^2 + x - 132 = 0$ (1); $(x - 11)(x + 12) = 0$, $x = 11$ (1); integers 11 and 12 (1).	1+1+1+1

SECTION — D (Long Answer) — Answers — 2 × 5 = 10

Q.	Expected Answer / Value Points	Marks
19	Split the middle term: $1 = -3 + 4$ because $ac = -12$ (1); regroup and factorise (1); roots $x = \frac{3}{2}, -2$ (1).	1+1+1
	OR: Breadth x, length $x + 3$: $x(x + 3) = 130$ (1); $x^2 + 3x - 130 = 0$ (1); $(x - 10)(x + 13) = 0$ (1); $x = 10$ m (breadth), length 13 m (reject negative) (2).	1+1+1+2
20	Split the middle term: $-7 = -6 - 1$ because $ac = 6$ (1); regroup and factorise (1); roots $x = \frac{1}{3}, 2$ (1).	1+1+1
	OR: Breadth x, length $x + 6$: $x(x + 6) = 91$ (1); $x^2 + 6x - 91 = 0$ (1); $(x - 7)(x + 13) = 0$ (1); $x = 7$ m (breadth), length 13 m (reject negative) (2).	1+1+1+2

Evaluator's note: Award step marks as shown; accept any correct alternative method. In internal-choice questions, evaluate only the option attempted. Deduct up to ½ mark for missing units or figures where required.

THE MASTERMIND COACHING CLASSES

CBSE / RBSE • NURSERY – CLASS XII • JAIPUR

Road No. 5, VKIA, Jaipur — 9887258679

ANSWER KEY & MARKING SCHEME SET 3

Chapter 4: **Quadratic Equations** | Class 10 Maths | Maximum Marks: 40

SECTION — A (Objective Type) — Answers — 10 × 1 = 10

Q.	Correct Option	Brief Reasoning
1	**(d)** 6	Compare with $ax^2+bx+c=0$
2	**(d)** -2 and 6	Put each factor equal to zero
3	**(b)** 16	$D=b^2-4ac$
4	**(b)** two equal real roots	$D=b^2-4ac=0$
5	**(c)** $x^2+2x+9=0$	Degree must be 2
6	**(a)** No	LHS $=2$ at $x=1$
7	**(b)** -8	Compare with $ax^2+bx+c=0$
8	**(d)** 1 and 7	Put each factor equal to zero
9	**(a)** -11	$D=b^2-4ac$
10	**(d)** $x^2-1x+3=0$	Degree must be 2

SECTION — B (Very Short Answer) — Answers — 4 × 2 = 8

Q.	Expected Answer / Value Points	Marks
11	$D=(1)^2-4(2)(3)=-23$ (1); no real roots (1).	1+1
12	Put $x=-4$: LHS $=0$ (1); equals RHS 0, so it is a root (1).	1+1
13	$x-6=0$ or $x+1=0$ (1); $x=6$ or $x=-1$ (1).	1+1
14	$D=(-7)^2-4(3)(2)=25$ (1); two distinct real roots (1).	1+1

SECTION — C (Short Answer) — Answers — 4 × 3 = 12

Q.	Expected Answer / Value Points	Marks
15	Split the middle term: $-9=-8-1$ because $ac=8$ (1); regroup and factorise (1); roots $x=1,8$ (1).	1+1+1
16	$a=5, b=-10, c=5$ (1); $D=0$ (1); two equal real roots (1).	1+1+1
17	Let integers be x and $x+1$: $x(x+1)=182$ (1); $x^2+x-182=0$ (1); $(x-13)(x+14)=0$, $x=13$ (1); integers 13 and 14 (1).	1+1+1+1
	OR: Split the middle term: $6=2+4$ because $ac=8$ (1); regroup and factorise (1); roots $x=-4,-2$ (1).	1+1+1
18	$a=2, b=7, c=3$ (1); $D=25$ (1); two distinct real roots (1).	1+1+1
	OR: Let integers be x and $x+1$: $x(x+1)=240$ (1); $x^2+x-240=0$ (1); $(x-15)(x+16)=0$, $x=15$ (1); integers 15 and 16 (1).	1+1+1+1

SECTION — D (Long Answer) — Answers — 2 × 5 = 10

Q.	Expected Answer / Value Points	Marks
19	Split the middle term: $-5=-8+3$ because $ac=-24$ (1); regroup and factorise (1); roots $x=-\frac{3}{4}, 2$ (1).	1+1+1
	OR: Breadth x, length $x+8$: $x(x+8)=153$ (1); $x^2+8x-153=0$ (1); $(x-9)(x+17)=0$ (1); $x=9$ m (breadth), length 17 m (reject negative) (2).	1+1+1+2
20	Split the middle term: $-3=-5+2$ because $ac=-10$ (1); regroup and factorise (1); roots $x=\frac{5}{2}, -1$ (1).	1+1+1
	OR: Breadth x, length $x+5$: $x(x+5)=204$ (1); $x^2+5x-204=0$ (1); $(x-12)(x+17)=0$ (1); $x=12$ m (breadth), length 17 m (reject negative) (2).	1+1+1+2

Evaluator's note: Award step marks as shown; accept any correct alternative method. In internal-choice questions, evaluate only the option attempted. Deduct up to ½ mark for missing units or figures where required.