THE MASTERMIND COACHING CLASSES

CBSE / RBSE • NURSERY – CLASS XII • JAIPUR

Road No. 5, VKIA, Jaipur — 9887258679

ANSWER KEY & MARKING SCHEME SET 1

Chapter 8: **Introduction to Trigonometry** | Class 10 Maths | Maximum Marks: 40

SECTION — A (Objective Type) — Answers — 10 × 1 = 10

Q.	Correct Option	Brief Reasoning
1	**(a)** \frac{4}{5}	Hypotenuse is 5
2	**(c)** \frac{12}{5}	Find the perpendicular
3	**(b)** 1	Substitute the standard values
4	**(d)** 1	$\tan(90^\circ - \theta) = \cot\theta$
5	**(a)** 1	$\cos(90^\circ - \theta) = \sin\theta$
6	**(d)** 4	$k^2(\sin^2 + \cos^2)$
7	**(a)** \frac{3}{5}	Reciprocal
8	**(d)** \frac{12}{13}	Hypotenuse is 13
9	**(b)** \frac{15}{8}	Find the perpendicular
10	**(c)** $\frac{8}{3}$	Substitute the standard values

SECTION — B (Very Short Answer) — Answers — 4 × 2 = 8

Q.	Expected Answer / Value Points	Marks
11	Hypotenuse $= 5$ (1); $\sec A = \frac{5}{3}$ (1).	1+1
12	$\cos 65^\circ = \sin 25^\circ$ and $\cot 50^\circ = \tan 40^\circ$ (2); value $= 1 + 1 = 2$ (1).	2+1
13	$\cot 69^\circ = \tan 21^\circ$ (1); value $= 1$ (1).	1+1
14	Hypotenuse $= 13$ (1); $\sec A = \frac{13}{5}$ (1).	1+1

SECTION — C (Short Answer) — Answers — 4 × 3 = 12

Q.	Expected Answer / Value Points	Marks
15	Hypotenuse $= \sqrt{12^2 + 5^2} = 13$ (1); $\sin\theta = \frac{12}{13}$, $\cos\theta = \frac{5}{13}$ (1); $\sec\theta = \frac{13}{5}$ (1).	1+1+1
16	$AC = \sqrt{7^2 + 24^2} = 25$ cm (1); $\sin A = \frac{BC}{AC} = \frac{24}{25}$, $\cos A = \frac{AB}{AC} = \frac{7}{25}$ (1); $\tan A = \frac{BC}{AB} = \frac{24}{7}$ (1).	1+1+1
17	$(1 - \cos^2\theta) = \sin^2\theta$ (1); $\sin^2\theta \cdot \csc^2\theta$ (1) $= 1$ (1).	1+1+1
	OR: Hypotenuse $= \sqrt{15^2 + 8^2} = 17$ (1); $\sin\theta = \frac{15}{17}$, $\cos\theta = \frac{8}{17}$ (1); $\sec\theta = \frac{17}{8}$ (1).	1+1+1
18	$AC = \sqrt{20^2 + 21^2} = 29$ cm (1); $\sin A = \frac{BC}{AC} = \frac{21}{29}$, $\cos A = \frac{AB}{AC} = \frac{20}{29}$ (1); $\tan A = \frac{BC}{AB} = \frac{21}{20}$ (1).	1+1+1
	OR: LHS $= \dfrac{\sin^2\theta + \cos^2\theta}{\sin\theta\cos\theta}$ (2); $= \dfrac{1}{\sin\theta\cos\theta} = \sec\theta\csc\theta$ (1).	2+1

SECTION — D (Long Answer) — Answers — 2 × 5 = 10

Q.	Expected Answer / Value Points	Marks
19	LHS $= \dfrac{\sin^2\theta + (1+\cos\theta)^2}{\sin\theta(1+\cos\theta)}$ (1); $= \dfrac{\sin^2\theta + 1 + 2\cos\theta + \cos^2\theta}{\sin\theta(1+\cos\theta)} = \dfrac{2 + 2\cos\theta}{\sin\theta(1+\cos\theta)}$ (2); $= \dfrac{2}{\sin\theta} = 2\csc\theta$ = RHS (2).	5
	OR: Divide numerator and denominator by $\cos A$ (1): $\dfrac{4\tan A + 3}{4\tan A + 3}$ (1); $= \dfrac{4(\frac{4}{3}) + 3}{4(\frac{4}{3}) + 3} = 1$ (2); verified with $\sin A = \frac{4}{5}$, $\cos A = \frac{3}{5}$ (1).	1+1+2+1
20	LHS $= \left(\dfrac{1 - \cos\theta}{\sin\theta}\right)^2$ (2); $= \dfrac{(1-\cos\theta)^2}{1 - \cos^2\theta}$ (1); $= \dfrac{(1-\cos\theta)^2}{(1-\cos\theta)(1+\cos\theta)} = \dfrac{1-\cos\theta}{1+\cos\theta}$ (2).	5
	OR: Divide numerator and denominator by $\cos A$ (1): $\dfrac{5\tan A + 2}{3\tan A + 2}$ (1); $= \dfrac{5(\frac{12}{5}) + 2}{3(\frac{12}{5}) + 2} = \frac{35}{23}$ (2); verified with $\sin A = \frac{12}{13}$, $\cos A = \frac{5}{13}$ (1).	1+1+2+1

Evaluator's note: Award step marks as shown; accept any correct alternative method. In internal-choice questions, evaluate only the option attempted. Deduct up to ½ mark for missing units or figures where required.

THE MASTERMIND COACHING CLASSES

CBSE / RBSE • NURSERY – CLASS XII • JAIPUR

Road No. 5, VKIA, Jaipur — 9887258679

ANSWER KEY & MARKING SCHEME SET 2

Chapter 8: **Introduction to Trigonometry** | Class 10 Maths | Maximum Marks: 40

SECTION — A (Objective Type) — Answers — 10 × 1 = 10

Q.	Correct Option	Brief Reasoning
1	**(b)** 1	$\tan(90^\circ-\theta)=\cot\theta$
2	**(b)** 1	$\cos(90^\circ-\theta)=\sin\theta$
3	**(d)** 9	$k^2(\sin^2+\cos^2)$
4	**(b)** \frac{5}{13}	Reciprocal
5	**(d)** \frac{15}{17}	Hypotenuse is 17
6	**(c)** \frac{24}{7}	Find the perpendicular
7	**(c)** 1	Substitute the standard values
8	**(b)** 1	$\tan(90^\circ-\theta)=\cot\theta$
9	**(c)** 1	$\cos(90^\circ-\theta)=\sin\theta$
10	**(b)** 16	$k^2(\sin^2+\cos^2)$

SECTION — B (Very Short Answer) — Answers — 4 × 2 = 8

Q.	Expected Answer / Value Points	Marks
11	$\cos 72^\circ=\sin 18^\circ$ and $\cot 55^\circ=\tan 35^\circ$ (2); value $=1+1=2$ (1).	2+1
12	$\cot 57^\circ=\tan 33^\circ$ (1); value $=1$ (1).	1+1
13	Hypotenuse $=17$ (1); $\sec A=\frac{17}{8}$ (1).	1+1
14	$\cos 58^\circ=\sin 32^\circ$ and $\cot 40^\circ=\tan 50^\circ$ (2); value $=1+1=2$ (1).	2+1

SECTION — C (Short Answer) — Answers — 4 × 3 = 12

Q.	Expected Answer / Value Points	Marks
15	Hypotenuse $=\sqrt{24^2+7^2}=25$ (1); $\sin\theta=\frac{24}{25}$, $\cos\theta=\frac{7}{25}$ (1); $\sec\theta=\frac{25}{7}$ (1).	1+1+1
16	$AC=\sqrt{9^2+40^2}=41$ cm (1); $\sin A=\frac{BC}{AC}=\frac{40}{41}$, $\cos A=\frac{AB}{AC}=\frac{9}{41}$ (1); $\tan A=\frac{BC}{AB}=\frac{40}{9}$ (1).	1+1+1
17	Expand: $2\sin^2\theta+2\cos^2\theta$ (2); $=2$ (1).	2+1
	OR: Hypotenuse $=\sqrt{21^2+20^2}=29$ (1); $\sin\theta=\frac{21}{29}$, $\cos\theta=\frac{20}{29}$ (1); $\sec\theta=\frac{29}{20}$ (1).	1+1+1
18	$AC=\sqrt{6^2+8^2}=10$ cm (1); $\sin A=\frac{BC}{AC}=\frac{4}{5}$, $\cos A=\frac{AB}{AC}=\frac{3}{5}$ (1); $\tan A=\frac{BC}{AB}=\frac{4}{3}$ (1).	1+1+1
	OR: LHS $=\dfrac{1}{\cos^2\theta}+\dfrac{1}{\sin^2\theta}=\dfrac{\sin^2\theta+\cos^2\theta}{\sin^2\theta\cos^2\theta}$ (2); $=\sec^2\theta\csc^2\theta$ (1).	2+1

SECTION — D (Long Answer) — Answers — 2 × 5 = 10

Q.	Expected Answer / Value Points	Marks
19	Write in $\sin,\cos$: LHS $=\dfrac{\sin^2\theta}{\cos\theta(\sin\theta-\cos\theta)}+\dfrac{\cos^2\theta}{\sin\theta(\cos\theta-\sin\theta)}$ (2); $=\dfrac{\sin^3\theta-\cos^3\theta}{\sin\theta\cos\theta(\sin\theta-\cos\theta)}$ (1); $=\dfrac{\sin^2\theta+\sin\theta\cos\theta+\cos^2\theta}{\sin\theta\cos\theta}=1+\sec\theta\csc\theta$ (2).	5
	OR: Divide numerator and denominator by $\cos A$ (1): $\dfrac{2\tan A+3}{5\tan A+1}$ (1); $=\dfrac{2(\frac{15}{8})+3}{5(\frac{15}{8})+1}=\frac{54}{83}$ (2); verified with $\sin A=\frac{15}{17}$, $\cos A=\frac{8}{17}$ (1).	1+1+2+1
20	Expand: $\sin^2\theta+2+\csc^2\theta+\cos^2\theta+2+\sec^2\theta$ (2); $=5+\csc^2\theta+\sec^2\theta$ (1); $=5+(1+\cot^2\theta)+(1+\tan^2\theta)=7+\tan^2\theta+\cot^2\theta$ (2).	5
	OR: Divide numerator and denominator by $\cos A$ (1): $\dfrac{3\tan A+1}{2\tan A+1}$ (1); $=\dfrac{3(\frac{24}{7})+1}{2(\frac{24}{7})+1}=\frac{79}{55}$ (2); verified with $\sin A=\frac{24}{25}$, $\cos A=\frac{7}{25}$ (1).	1+1+2+1

Evaluator's note: Award step marks as shown; accept any correct alternative method. In internal-choice questions, evaluate only the option attempted. Deduct up to ½ mark for missing units or figures where required.

THE MASTERMIND COACHING CLASSES

CBSE / RBSE • NURSERY – CLASS XII • JAIPUR

Road No. 5, VKIA, Jaipur — 9887258679

ANSWER KEY & MARKING SCHEME SET 3

Chapter 8: **Introduction to Trigonometry** | Class 10 Maths | Maximum Marks: 40

SECTION — A (Objective Type) — Answers — 10 × 1 = 10

Q.	Correct Option	Brief Reasoning
1	**(a)** \frac{8}{17}	Reciprocal
2	**(a)** \frac{24}{25}	Hypotenuse is 25
3	**(d)** \frac{21}{20}	Find the perpendicular
4	**(a)** $\frac{9}{4}$	Substitute the standard values
5	**(c)** 1	$\tan(90^\circ - \theta) = \cot\theta$
6	**(b)** 1	$\cos(90^\circ - \theta) = \sin\theta$
7	**(b)** 25	$k^2(\sin^2 + \cos^2)$
8	**(d)** \frac{21}{29}	Hypotenuse is 29
9	**(b)** \frac{40}{9}	Find the perpendicular
10	**(b)** 1	Substitute the standard values

SECTION — B (Very Short Answer) — Answers — 4 × 2 = 8

Q.	Expected Answer / Value Points	Marks
11	$\cot 43^\circ = \tan 47^\circ$ (1); value $= 1$ (1).	1+1
12	Hypotenuse $= 25$ (1); $\sec A = \frac{25}{7}$ (1).	1+1
13	$\cos 76^\circ = \sin 14^\circ$ and $\cot 29^\circ = \tan 61^\circ$ (2); value $= 1 + 1 = 2$ (1).	2+1
14	$\cot 32^\circ = \tan 58^\circ$ (1); value $= 1$ (1).	1+1

SECTION — C (Short Answer) — Answers — 4 × 3 = 12

Q.	Expected Answer / Value Points	Marks
15	Hypotenuse $= \sqrt{40^2 + 9^2} = 41$ (1); $\sin\theta = \frac{40}{41}$, $\cos\theta = \frac{9}{41}$ (1); $\sec\theta = \frac{41}{9}$ (1).	1+1+1
16	$AC = \sqrt{9^2 + 12^2} = 15$ cm (1); $\sin A = \frac{BC}{AC} = \frac{4}{5}$, $\cos A = \frac{AB}{AC} = \frac{3}{5}$ (1); $\tan A = \frac{BC}{AB} = \frac{4}{3}$ (1).	1+1+1
17	$1 + \tan^2 A = \sec^2 A$ (1); LHS $= (1 - \tan^2 A)\cos^2 A$ (1); $= \cos^2 A - \sin^2 A$ (1).	1+1+1
	OR: Hypotenuse $= \sqrt{8^2 + 6^2} = 10$ (1); $\sin\theta = \frac{4}{5}$, $\cos\theta = \frac{3}{5}$ (1); $\sec\theta = \frac{5}{3}$ (1).	1+1+1
18	$AC = \sqrt{12^2 + 16^2} = 20$ cm (1); $\sin A = \frac{BC}{AC} = \frac{4}{5}$, $\cos A = \frac{AB}{AC} = \frac{3}{5}$ (1); $\tan A = \frac{BC}{AB} = \frac{4}{3}$ (1).	1+1+1
	OR: $(1 + \tan^2 A) = \sec^2 A$ (1); $\sec^2 A \sin^2 A = \dfrac{\sin^2 A}{\cos^2 A}$ (1) $= \tan^2 A$ (1).	1+1+1

SECTION — D (Long Answer) — Answers — 2 × 5 = 10

Q.	Expected Answer / Value Points	Marks
19	In sin, cos: $\left(\dfrac{\sin A + \cos A - 1}{\sin A}\right)\left(\dfrac{\cos A + \sin A + 1}{\cos A}\right)$ (2); numerator $= (\sin A + \cos A)^2 - 1 = 2\sin A\cos A$ (2); so value $= 2$ (1).	5
	OR: Divide numerator and denominator by $\cos A$ (1): $\dfrac{5\tan A + 4}{5\tan A + 2}$ (1); $= \dfrac{5(\frac{21}{20}) + 4}{5(\frac{21}{20}) + 2} = \frac{37}{29}$ (2); verified with $\sin A = \frac{21}{29}$, $\cos A = \frac{20}{29}$ (1).	1+1+2+1
20	$\dfrac{1}{\cos A}(1 - \sin A) \cdot \dfrac{1 + \sin A}{\cos A}$ (2); $= \dfrac{1 - \sin^2 A}{\cos^2 A}$ (2); $= 1$ (1).	5
	OR: Divide numerator and denominator by $\cos A$ (1): $\dfrac{7\tan A + 2}{4\tan A + 1}$ (1); $= \dfrac{7(\frac{40}{9}) + 2}{4(\frac{40}{9}) + 1} = \frac{298}{169}$ (2); verified with $\sin A = \frac{40}{41}$, $\cos A = \frac{9}{41}$ (1).	1+1+2+1

Evaluator's note: Award step marks as shown; accept any correct alternative method. In internal-choice questions, evaluate only the option attempted. Deduct up to ½ mark for missing units or figures where required.