

Exercise sheet 2

05. May 2023

Exercise 1

Consider the ODE

$$\begin{aligned}\frac{d}{dt}x(t) &= \lambda x(t) \\ x(t_0) &= x_0,\end{aligned}$$

for some $\lambda \in \mathbb{C}$. Set $t_n = t_0 + n\Delta t$ for some $\Delta t > 0, n = 1, 2, \dots$. Consider the following methods to propagate from t_n to t_{n+1} :

- Euler method:

$$x_{n+1} = x_n + \Delta t \lambda x_n$$

- Implicit Euler method:

$$x_{n+1} = x_n + \Delta t \lambda x_{n+1}$$

Determine the regions of stability for both methods. Assume that λ lies in the left half-plane, i.e., it has a negative real part. Under what conditions on Δt is the implicit Euler method stable? Assume that λ is purely imaginary, i.e., it has a zero real part. Is Euler's method a good choice for solving the Cauchy problem defined above?

Exercise 2

Consider a p -th order accurate summation by part operator given by a grid \underline{x} , a first derivative operator D and a mass matrix M .

- Assume that M is diagonal and show that $\underline{1}^T M \underline{x}^{2q} = \frac{1}{2q+1}(x_{\max}^{2q+1} - x_{\min}^{2q+1})$ for $q \in \{1, \dots, p-1\}$.

Remark: This shows that M induces a quadrature rule that integrates monomials of even degrees $\leq 2p-1$ exactly. It, hence, completes the proof of theorem 1.15 in the lecture notes.

- Assuming that M is not necessary diagonal, show that $\underline{1}^T M$ still yields a quadrature rule that is at least $(p - 1)$ -th order accurate.

Exercise 3

Consider the problem

$$\begin{aligned}\partial_t u(t, x) + a \partial_x u(t, x) &= 0 && \text{in } (0, T) \times [0, 2] \\ u(0, x) &= \sin(\pi x) && \text{for } x \in [-1, 1] \\ u(t, 0) &= -\sin(\pi t) && \text{for } x \in [-1, 1]\end{aligned}$$

for some $a = 1/2$. Solve this equation using the SBP operator from example 1.10 and a third or fourth order Runge-Kuta method. Conduct convergence studies where you consider as a metric the L^2 error, i.e.,

$$\|u - u_{\text{approximate}}\|_{L^2} = \left(\int_0^2 (u(x) - u_{\text{approximate}}(x))^2 dx \right)^{1/2}.$$

In the convergence studies differentiate between the transient region $\text{TR} = \{x \in [0, 2] \mid x > at\}$ and the asymptotic region $\text{AR} = [0, 2] \setminus \text{TR}$. Repeat the convergence study for different values of $a > 0$ and a higher-order SBP operator.

Hint: To compute the error remember that the mass matrix M induces a quadrature rule. You can impose the boundary condition either weakly or strongly. An example of a weak imposition is given in the lecture notes.