

FE5222 Advanced Derivative Pricing

Wu Lei

Risk Management Institute, National University of Singapore

rmiwul@nus.edu.sg

November 8, 2019

Overview

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

1 Introduction

2 Dupire's Equation

3 Understanding Local Volatility

4 How to Build Local Volatility Surface

5 Implied Binomial Tree

Local Volatility Model

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

The assumption of constant volatility in BSM model is inconsistent with market observations.

Is there a BSM-like model that can price European options in the market consistently?

Local Volatility Model

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

- Dupire (1994) developed a local volatility model for continuous time and showed that there exists a unique risk neutral diffusion process that is consistent with European option prices.
- Derman & Iraj Kani (1994) developed a tree model which is consistent with market prices for European options.

Local Volatility Model

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation


Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

1. A unique diffusion process

If we restrict ourselves to diffusions, there is a unique risk-neutral (drift equal to the short-term rate) process for the spot which is compatible with European option prices:



Source: Dupire (1994)

Local Volatility Model

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

In LVM, the instantaneous volatility is a deterministic function of t and S_t

$$\frac{dS_t}{S_t} = rdt + \sigma(t, S_t)dW_t$$

Local Volatility Model

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

In LVM, the pricing PDE is

$$V_t + rSV_s + \frac{1}{2}\sigma^2(t, S)S^2V_{SS} - rV = 0 \quad (1)$$

Dupire's Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

In this section, we will discuss

- Kolmogorov Backward Equation
- Kolmogorov Forward Equation
- Dupire's Equation

Kolmogorov Backward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Consider the stochastic differential equation

$$dX_t = \mu(t, X_t)dt + \sigma(t, X_t)dW_t$$

Kolmogorov Backward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Fix $T > t$, let $p(t, T, x, y)$ be the transition probability density for the solution to this equation.

It is the probability density function of X_T if we solve the equation with initial condition $X_t = x$.

Kolmogorov Backward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Theorem

The transition density function $p(t, T, x, y)$ satisfies the Kolmogorov backward equation

$$p_t(t, T, x, y) + \mu(t, x)p_x(t, T, x, y) + \frac{1}{2}\sigma^2(t, x)p_{xx}(t, T, x, y) = 0$$

Kolmogorov Backward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

In the following proof, we will need the concept of a smooth function and compact support.

- A smooth function is a function that has derivatives of all orders.
- The support of a function f is

$$\text{supp } f = \overline{\{x : f(x) \neq 0\}}$$

- A compact set in \mathbb{R}^n is a closed and bounded set.

Kolmogorov Backward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

One example for a smooth function with compact support is the so called *bump function* defined as

$$f(x) = \begin{cases} e^{-\frac{1}{1-x^2}} & \forall -1 < x < 1 \\ 0 & \text{otherwise} \end{cases}$$

Kolmogorov Backward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Lemma

Let f be an integrable function such that

$$\int f(x)h(x)dx = 0$$

for all smooth and compact function h . Then $f(x) = 0$ for (almost surely) all x .

Remark: In fact, smoothness is not necessary for this lemma. However we will need smoothness for the derivation of Komogorov Forward Equation.

Kolmogorov Backward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

For any smooth function $h(x)$ with compact support, let

$$\begin{aligned} g(t, x) &= \mathbb{E}[h(X_T)] \\ &= \int h(y)p(t, T, x, y)dy \end{aligned}$$



Kolmogorov Backward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Taking partial derivatives w.r.t. t and x respectively, we have

$$g_t(t, x) = \int h(y) p_t(t, T, x, y) dy \quad (2)$$

$$g_x(t, x) = \int h(y) p_x(t, T, x, y) dy \quad (3)$$

$$g_{xx}(t, x) = \int h(y) p_{xx}(t, T, x, y) dy \quad (4)$$



Kolmogorov Backward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

From Feynman-Kac Theorem, we have

$$g_t(t, x) + \mu(t, x)g_x(t, x) + \frac{1}{2}\sigma^2(t, x)g_{xx}(t, x) = 0$$

Replacing Equation (2), (3) and (4) into the above equation, we have

$$\int h(y)p_t dy + \mu(t, x) \int h(y)p_x dy + \frac{1}{2}\sigma^2(t, x) \int h(y)p_{xx} dy = 0$$



Kolmogorov Backward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Hence

$$\int h(y) \left[p_t + \mu(t, x) p_x + \frac{1}{2} \sigma^2(t, x) p_{xx} \right] dy = 0$$



Kolmogorov Backward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Since this holds for all smooth function h with compact support, we must have

$$p_t + \mu(t, x)p_x + \frac{1}{2}\sigma^2(t, x)p_{xx} = 0 \quad \text{Q.E.D.} \quad (5)$$



Kolmogorov Forward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Theorem

The transition density function $p(t, T, x, y)$ satisfies the Kolmogorov forward equation

$$\frac{\partial p}{\partial T} + \frac{\partial}{\partial y} (\mu(T, y)p) - \frac{1}{2} \frac{\partial^2}{\partial y^2} (\sigma^2(T, y)p) = 0$$

Remark: It is also called Fokker-Planck equation.

Kolmogorov Forward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Let h be a smooth function with compact support. By Ito's Lemma, we have

$$\begin{aligned} dh(X_s) &= h_x dX_s + \frac{1}{2} h_{xx} d[X, X](s) \\ &= \left[\mu(s, X_s) h_x(X_s) + \frac{1}{2} \sigma^2(s, X_s) h_{xx}(X_s) \right] ds \quad (6) \\ &\quad + h_x(X_s) \sigma(s, X_s) dW_s \end{aligned}$$



Kolmogorov Forward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Consider the process X that starts from t with initial condition $X(t) = x$.

Integrating Equation (6) from t to T , we obtain

$$h(X_T) = h(X_t) + I_1 + I_2$$

where

$$I_1 = \int_t^T \left[\mu(s, X_s) h_x(X_s) + \frac{1}{2} \sigma^2(s, X_s) h_{xx}(X_s) \right] ds$$

and

$$I_2 = \int_t^T h_x(X_s) \sigma(s, X_s) dW_s$$

Kolmogorov Forward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Since I_2 is an Ito's integral whose mean is zero, we have

$$\mathbb{E}[h(X_T)] = h(x) + \mathbb{E}[I_1] \quad (7)$$



Kolmogorov Forward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

The LHS of Equation (7) is

$$\mathbb{E}[h(X_T)] = \int_{-\infty}^{\infty} h(y)p(t, T, x, y)dy$$



Kolmogorov Forward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

$$\begin{aligned} & \mathbb{E}[I_1] \\ &= \mathbb{E} \left[\int_t^T [\mu(s, X_s) h_x(X_s) + \frac{1}{2} \sigma^2(s, X_s) h_{xx}(X_s)] ds \right] \\ &= \int_t^T \mathbb{E} [\mu(s, X_s) h_x(X_s) + \frac{1}{2} \sigma^2(s, X_s) h_{xx}(X_s)] ds \\ &= \int_t^T \int_{-\infty}^{\infty} p(t, s, x, y) [\mu(s, y) h_x(y) + \frac{1}{2} \sigma^2(s, y) h_{xx}(y)] dy ds \\ &= \int_t^T \left[\int_{-\infty}^{\infty} p(t, s, x, y) \mu(s, y) h_x(y) dy \right] ds \\ &\quad + \frac{1}{2} \int_t^T \left[\int_{-\infty}^{\infty} p(t, s, x, y) \sigma^2(s, y) h_{xx}(y) dy \right] ds \end{aligned}$$



Kolmogorov Forward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Now we evaluate

$$\int_{-\infty}^{\infty} p(t, s, x, y) \mu(s, y) h_x(y) dy$$

and

$$\int_{-\infty}^{\infty} p(t, s, x, y) \sigma^2(s, y) h_{xx}(y) dy$$



Kolmogorov Forward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Integrating by parts and using the fact that $h(x)$, $h_x(x)$ and $h_{xx}(x)$ vanish when $|x|$ is large enough, we have

$$\begin{aligned} & \int_{-\infty}^{\infty} p(t, s, x, y) \mu(s, y) h_x(y) dy \\ = & - \int_{-\infty}^{\infty} \frac{\partial}{\partial y} (p(t, s, x, y) \mu(s, y)) h(y) dy \end{aligned} \quad (8)$$

and

$$\begin{aligned} & \int_{-\infty}^{\infty} p(t, s, x, y) \sigma^2(s, y) h_{xx}(y) dy \\ = & \int_{-\infty}^{\infty} \frac{\partial^2}{\partial y^2} (p(t, s, x, y) \sigma^2(s, y)) h(y) dy \end{aligned} \quad (9)$$



Kolmogorov Forward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Hence

$$\begin{aligned} & \mathbb{E}[I_1] \\ = & - \int_t^T \left[\int_{-\infty}^{\infty} \frac{\partial}{\partial y} (p(t, s, x, y) \mu(s, y)) h(y) dy \right] ds \\ & + \frac{1}{2} \int_t^T \left[\int_{-\infty}^{\infty} \frac{\partial^2}{\partial y^2} (p(t, s, x, y) \sigma^2(s, y)) h(y) dy \right] ds \end{aligned}$$



Kolmogorov Forward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Substituting these to Equation (7), we obtain

$$\begin{aligned} & \int_{-\infty}^{\infty} h(y) p(t, T, x, y) dy \\ = & h(x) - \int_t^T \left[\int_{-\infty}^{\infty} \frac{\partial}{\partial y} (p(t, s, x, y) \mu(s, y)) h(y) dy \right] ds \\ & + \frac{1}{2} \int_t^T \left[\int_{-\infty}^{\infty} \frac{\partial^2}{\partial y^2} (p(t, s, x, y) \sigma^2(s, y)) h(y) dy \right] ds \end{aligned}$$



Kolmogorov Forward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Taking derivative w.r.t. to T , we have

$$\begin{aligned} & \int_{-\infty}^{\infty} h(y) \frac{\partial}{\partial T} p(t, T, x, y) dy \\ = & - \int_{-\infty}^{\infty} \frac{\partial}{\partial y} (p(t, T, x, y) \mu(T, y)) h(y) dy \\ & + \frac{1}{2} \int_{-\infty}^{\infty} \frac{\partial^2}{\partial y^2} (p(t, T, x, y) \sigma^2(T, y)) h(y) dy \end{aligned}$$



Kolmogorov Forward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Re-arranging it, we have

$$\int_{-\infty}^{\infty} h(y) \left[\frac{\partial p}{\partial T} + \frac{\partial}{\partial y} (\mu(T, y)p) - \frac{1}{2} \frac{\partial^2}{\partial y^2} (\sigma^2(T, y)p) \right] dy = 0$$

which implies

$$\frac{\partial p}{\partial T} + \frac{\partial}{\partial y} (\mu(T, y)p) - \frac{1}{2} \frac{\partial^2}{\partial y^2} (\sigma^2(T, y)p) = 0 \quad \text{Q.E.D.}$$



Kolmogorov Backward/Forward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Kolmogorov Backward Equation

$$p_t(t, T, x, y) + \mu(t, x)p_x(t, T, x, y) + \frac{1}{2}\sigma^2(t, x)p_{xx}(t, T, x, y) = 0$$

Kolmogorov Forward Equation

$$\frac{\partial p}{\partial T} + \frac{\partial}{\partial y} (\mu(T, y)p) - \frac{1}{2} \frac{\partial^2}{\partial y^2} (\sigma^2(T, y)p) = 0$$

Kolmogorov Backward/Forward Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

- Kolmogorov backward equation
Fix T , it is an PDE of initial condition $X_t = x$.
- Kolmogorov forward equation
Fix initial condition $X(t) = x$ and it is an PDE w.r.t. T
and $X_T = y$.

Dupire's Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Consider the SDE

$$\frac{dS_t}{S_t} = rdt + \sigma(t, S_t)dW_t$$

where r is a constant and W_t is a standard Brownina motion.
Let $C(T, K)$ be the price of a call option with expiry T and strike K , given $S(0) = S_0$.

Dupire's Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Theorem

Let $C(T, K)$ be the price of a call option with strike K and expiry T . Then the following so called Dupire's equation holds

$$C_T + rKC_K - \frac{1}{2}\sigma^2(T, K)K^2C_{KK} = 0$$

Dupire's Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Remark: Dupire's equation is often written as

$$\sigma^2(T, K) = \frac{C_T + rKC_K(T, K)}{\frac{1}{2}K^2C_{KK}(T, K)} \quad (10)$$

The RHS of Equation (10) can be used to define the notion of local volatility.

Dupire's Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

- Dupire's equation is a forward equation. It is PDE for the option price with different expiry T and strike K .
- On the contrary, the PDE Equation (1) is the option price with fixed expiry T and strike K , but different time t and spot price S_t .
- In practice, Dupire's equation is often used for model calibration and the PDE Equation (1) is used for pricing.

Dupire's Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Let $p(0, T, S_0, y)$ be the transition density function for the stock price process that starts at time $t = 0$ with S_0 . For notational simplicity, we write it as $p(T, y)$.

The call price is

$$C(T, K) = e^{-rT} \int (y - K)^+ p(T, y) dy$$

Taking partial derivative w.r.t. T , we have

$$\begin{aligned} & C_T(T, K) \\ = & -rC(T, K) + e^{-rT} \int (y - K)^+ \frac{\partial p}{\partial T} dy \end{aligned}$$



Dupire's Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

By Kolmogorov forward equation

$$= \frac{\partial p}{\partial T} (ryp(T, y)) + \frac{1}{2} \frac{\partial^2}{\partial y^2} (\sigma^2(T, y) y^2 p(T, y))$$

Hence

$$\begin{aligned} & \int (y - K)^+ \frac{\partial p}{\partial T} dy \\ = & - \int (y - K)^+ \frac{\partial}{\partial y} (ryp(T, y)) dy \\ & + \frac{1}{2} \int (y - K)^+ \frac{\partial^2}{\partial y^2} (\sigma^2(T, y) y^2 p(T, y)) dy \end{aligned}$$



Dupire's Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Integrating by parts, we have

$$\begin{aligned} & \int (y - K)^+ \frac{\partial}{\partial y} (ryp(T, y)) dy \\ &= \int_K^\infty (y - K) \frac{\partial}{\partial y} (ryp(T, y)) dy \\ &= -r \int_K^\infty yp(T, y) dy \end{aligned}$$

with the assumption

$$\lim_{y \rightarrow \infty} (y - k)yp(T, y) = 0$$



Dupire's Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Similarly

$$\begin{aligned} & \int (y - K)^+ \frac{\partial^2}{\partial y^2} (\sigma^2(T, y) y^2 p(T, y)) dy \\ &= \int_K^\infty (y - K) \frac{\partial^2}{\partial y^2} (\sigma^2(T, y) y^2 p(T, y)) dy \\ &= \sigma^2(T, K) K^2 p(T, K) \end{aligned}$$

with the assumption

$$\lim_{y \rightarrow \infty} (y - K) \frac{\partial}{\partial y} (\sigma^2(T, y) y^2 p(T, y))$$

and

$$\lim_{y \rightarrow \infty} \sigma^2(T, y) y^2 p(T, y) = 0$$



Dupire's Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

Hence

$$\begin{aligned} & \int (y - K)^+ \frac{\partial}{\partial T} p(T, y) dy \\ = & r \int_K^\infty y p(T, y) dy + \frac{1}{2} \sigma^2(T, K) K^2 p(T, K) \end{aligned}$$



Dupire's Equation

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Proof.

It follows that

$$\begin{aligned} & C_T(T, K) \\ = & -rC(T, K) + e^{-rT} \int (y - K)^+ \frac{\partial}{\partial T} p dy \\ = & -rC(T, K) + re^{-rT} \int_K^\infty yp(T, y) dy \\ & + \frac{1}{2} e^{-rT} \sigma^2(T, K) K^2 p(T, K) \\ = & rKe^{-rT} \int_K^\infty p(T, y) dy + \frac{1}{2} e^{-rT} \sigma^2(T, K) K^2 p(T, K) \\ = & -rKC_K(T, K) + \frac{1}{2} \sigma^2(T, K) K^2 C_{KK}(T, K) \end{aligned}$$

where in the last equality we use the identities for implied risk-neutral probability density. Q.E.D. □

Understanding Local Volatility

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

In this section we will look at some facts and properties of local volatility.

- Local variance as the conditional expectation of instantaneous variance
- Local volatility in terms of implied volatility
- Implied variance as the average of local variance over the life of option when there is no skew.

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Theorem (Tanaka-Meyer Formula)

Let X_t be an Ito process such that

$$dX_t = \alpha(t, X_t)dt + \beta(t, X_t)dW_t$$

and K is a real number. Then

$$\begin{aligned} & (X_t - K)^+ \\ &= (X_0 - K)^+ + \int_0^t H_K(X_s) dX_s + \frac{1}{2} \int_0^t \delta_K(X_s) d[X, X](s) \end{aligned}$$

where $H_K(\cdot)$ is the Heaviside function and $\delta_K(\cdot)$ is the Dirac delta function.

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

In differential form, we have

$$d(X_t - K)^+ = H_K(X_t)dX_t + \frac{1}{2}\delta_K(X_t)d[X, X](t)$$

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

If we let $f(x) = (x - K)^+$, then

$$f'(x) = H_K(x)$$

and

$$f''(x) = \delta_K(x)$$

Hence Tanaka-Meyer formula is

$$df(X_t) = f'(X_t)dX_t + \frac{1}{2}f''(X_t)d[X, X](t)$$

which is a generalization of Ito's formula.

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Now we assume the stock price follows

$$\frac{dS_t}{S_t} = rdt + \sigma(t, \omega)dW_t$$

where σ is an arbitrary adapted-process.

We want to investigate how the instantaneous volatility $\sigma(t, \omega)$ is related to local volatility as defined in Equation (10).

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

The price for the call option with expiry T and strike K is

$$C(T, K) = e^{-rT} \mathbb{E} [(S_T - K)^+]$$

Taking derivative w.r.t. T we have

$$C_T = -rC + e^{-rT} \frac{\partial}{\partial T} \mathbb{E} [(S_T - K)^+] \quad (11)$$

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

To evaluate

$$\frac{\partial}{\partial T} \mathbb{E} [(S_T - K)^+]$$

we will use Tanaka-Meyer formula.

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Applying Tanaka-Meyer formula, we have

$$\begin{aligned} & d(S_T - K)^+ \\ &= H_K(S_T) dS_T + \frac{1}{2} \delta_K(S_T) d[S, S](T) \\ &= \left[H_K(S_T) r S_T + \frac{1}{2} S_T^2 \sigma^2(T, \omega) \delta_K(S_T) \right] dT \\ &\quad + H_K(S_T) \sigma(T, \omega) dW_T \end{aligned}$$

Taking expectation on both sides, we have

$$\begin{aligned} & \mathbb{E} [d(S_T - K)^+] \\ &= \mathbb{E} \left[H_K(S_T) r S_T + \frac{1}{2} S_T^2 \sigma^2(T, \omega) \delta_K(S_T) \right] dT \\ &\quad + \mathbb{E} [H_K(S_T) \sigma(T, \omega) dW_T] \end{aligned}$$

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Since $H_K(S_T)$ and $\sigma(T, \omega)$ are \mathcal{F}_T -measurable, using iterated property of conditional expectation, we have

$$\begin{aligned} & \mathbb{E} [H_K(S_T) \sigma(T, \omega) dW_T] \\ &= \mathbb{E} [\mathbb{E} [H_K(S_T) \sigma(T, \omega) dW_T | \mathcal{F}_T]] \\ &= \mathbb{E} [H_K(S_T) \sigma(T, \omega) \mathbb{E} [dW_T | \mathcal{F}_T]] \\ &= 0 \end{aligned}$$

where in the last equality we use the fact that

$$\mathbb{E} [dW_T | \mathcal{F}_T] = 0$$

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Hence

$$\begin{aligned} & \mathbb{E} [d(S_T - K)^+] \\ = & \mathbb{E} \left[H_K(S_T) r S_T + \frac{1}{2} S_T^2 \sigma^2(T, \omega) \delta_K(S_T) \right] dT \end{aligned}$$

Dividing both sides by dT , we have

$$\begin{aligned} & \frac{\partial}{\partial T} \mathbb{E} [(S_T - K)^+] \\ = & \mathbb{E} \left[H_K(S_T) r S_T + \frac{1}{2} S_T^2 \sigma^2(T, \omega) \delta_K(S_T) \right] \\ = & r \mathbb{E} [H_K(S_T) S_T] + \frac{1}{2} \mathbb{E} [S_T^2 \sigma^2(T, \omega) \delta_K(S_T)] \end{aligned} \tag{12}$$

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

To evaluate $\mathbb{E}[H_K(S_T)S_T]$, we notice that

$$\begin{aligned} & \mathbb{E}[H_K(S_T)S_T] \\ = & \mathbb{E}[(S_T - K)H_K(S_T)] + K\mathbb{E}[H_K(S_T)] \end{aligned} \quad (13)$$

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Note that

$$\begin{aligned}\mathbb{E}[(S_T - K)H_K(S_T)] &= \mathbb{E}[(S_T - K)^+] \\ &= e^{rT} C\end{aligned}\quad (14)$$

Since

$$\begin{aligned}\frac{\partial}{\partial K} \mathbb{E}[(S_T - K)^+] &= \mathbb{E}\left[\frac{\partial(S_T - K)^+}{\partial K}\right] \\ &= -\mathbb{E}[H_K(S_T)]\end{aligned}$$

we have

$$\begin{aligned}\mathbb{E}[H_K(S_T)] &= -\frac{\partial}{\partial K} \mathbb{E}[(S_T - K)^+] \\ &= -e^{rT} C_K\end{aligned}\quad (15)$$

Local Variance as Conditional Expectation of Instantaneous

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Substituting Equation (14) and (15) into Equation (13), we have

$$\mathbb{E}[H_K(S_T)S_T] = e^{rT} (C - KC_K) \quad (16)$$

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Now we evaluate $\mathbb{E} [S_T^2 \sigma^2(T, \omega) \delta_K(S_T)]$.

Using tower property of conditional expectation, we have

$$\begin{aligned} & \mathbb{E} [S_T^2 \sigma^2(T, \omega) \delta_K(S_T)] \\ &= \mathbb{E} [\mathbb{E} [S_T^2 \sigma^2(T, \omega) \delta_K(S_T) | S_T]] \\ &= \mathbb{E} [\mathbb{E} [\sigma^2(T, \omega) | S_T] S_T^2 \delta_K(S_T)] \end{aligned}$$

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Let $p(x)$ be the probability density function of S_T , then

$$\begin{aligned} & \mathbb{E} [\mathbb{E} [\sigma^2(T, \omega) | S_T] S_T^2 \delta_K(S_T)] \\ &= \int \mathbb{E} [\sigma^2(T, \omega) | S_T = x] x^2 \delta_K(x) p(x) dx \\ &= K^2 p(K) \mathbb{E} [\sigma^2(T, \omega) | S_T = K] \end{aligned}$$

Note that in the last equality we use the following property of Dirac delta function

$$\int f(x) \delta_K(x) dx = f(K)$$

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Since

$$\begin{aligned} p(K) &= \frac{\partial^2}{\partial K^2} \mathbb{E} [(S_T - K)^+] \\ &= e^{rT} C_{KK} \end{aligned}$$

we have

$$\begin{aligned} &\mathbb{E} [S_T^2 \sigma^2(T, \omega) \delta_K(S_T)] \\ &= e^{rT} K^2 C_{KK} \mathbb{E} [\sigma^2(T, \omega) | S_T = K] \end{aligned} \tag{17}$$

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Substituting Equation (16) and Equation (17) into Equation (12), we have

$$\begin{aligned} & \frac{\partial}{\partial T} \mathbb{E} [(S_T - K)^+] \\ = & e^{rT} \left(r(C - KC_K) + \frac{1}{2} K^2 C_{KK} \mathbb{E} [\sigma^2(T, \omega) | S_T = K] \right) \end{aligned} \quad (18)$$

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Substituting the above equation into Equation (11), we have

$$C_T = -rKC_K + \frac{1}{2}K^2C_{KK}\mathbb{E}[\sigma^2(T, \omega)|S_T = K]$$

which implies

$$\mathbb{E}[\sigma^2(T, \omega)|S_T = K] = \frac{C_T + rKC_K}{\frac{1}{2}K^2C_{KK}} \quad (19)$$

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Comparing Equation (19) with Equation (10), we can see that local variance is the risk-neutral expectation of the instantaneous variance conditional on the final stock price S_T being equal to strike K .

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

We can see that in general the solution for Equation (19) is not unique, there are two instantaneous volatility process $\sigma = \sigma'$ such that

$$\mathbb{E} [\sigma(T, \omega)^2 | S_T = K] = \mathbb{E} [\sigma'(T, \omega)^2 | S_T = K]$$

Knowing the vanilla option prices is not enough to find σ in general.

Local Variance as Conditional Expectation of Instantaneous Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

However if we restrict ourselves to the case $\sigma(t, S_t)$ as a deterministic function of t and S_t , we can uniquely determine $\sigma(t, S_t)$ from vanilla option prices from Dupire's equation.

Local Volatility from Implied Volatility

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

In practice we often work in terms of implied volatilities as opposed to price. In the following we will derive the local volatility in terms of implied volatilities.

Local Volatility from Implied Volatility

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Recall that Dupire's equation

$$\sigma(T, K)^2 = \frac{C_T + rKC_K}{\frac{1}{2}K^2 C_{KK}}$$

We assume

$$C(T, K) = C_{BSM}(T, K, \Sigma(T, K))$$

where C_{BSM} is the BSM pricing formula and $\Sigma(T, K)$ is the implied volatility.

Local Volatility from Implied Volatility

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

The numerator is

$$C_T + rKC_K = \frac{\partial C_{BSM}}{\partial T} + \frac{\partial C_{BSM}}{\partial \Sigma} \frac{\partial \Sigma}{\partial T} + rK \left(\frac{\partial C_{BSM}}{\partial K} + \frac{\partial C_{BSM}}{\partial \Sigma} \frac{\partial \Sigma}{\partial K} \right) \quad (20)$$

Local Volatility from Implied Volatility

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

From

$$\frac{\partial C_{BSM}}{\partial T} = \frac{S\Sigma\phi(d_1)}{2\sqrt{T}} + re^{-rT}K\Phi(d_2)$$

$$\frac{\partial C_{BSM}}{\partial K} = -e^{-rT}\Phi(d_2)$$

and

$$\frac{\partial C_{BSM}}{\partial \Sigma} = S\sqrt{T}\phi(d_1)$$

we have

$$\frac{\partial C_{BSM}}{\partial T} + rK\frac{\partial C_{BSM}}{\partial K} = \frac{\partial C_{BSM}}{\partial \Sigma} \frac{\Sigma}{2T}$$

Local Volatility from Implied Volatility

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Substituting this into Equation (20), the numerator becomes

$$\frac{\partial C_{BSM}}{\partial \Sigma} \left(\frac{\Sigma}{2T} + \frac{\partial \Sigma}{\partial T} + rK \frac{\partial \Sigma}{\partial K} \right) \quad (21)$$

Local Volatility from Implied Volatility

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

For the denominator, we have

$$= \frac{C_{KK}}{\frac{\partial^2 C_{BSM}}{\partial K^2}} + 2 \frac{\frac{\partial^2 C_{BSM}}{\partial K \partial \Sigma}}{\frac{\partial^2 C_{BSM}}{\partial \Sigma^2}} \frac{\partial \Sigma}{\partial K} + \frac{C_{BSM}}{\frac{\partial^2 C_{BSM}}{\partial \Sigma^2}} \left(\frac{\partial \Sigma}{\partial K} \right)^2 + \frac{C_{BSM}}{\frac{\partial^2 C_{BSM}}{\partial \Sigma^2}} \frac{\partial^2 \Sigma}{\partial K^2}$$

Local Volatility from Implied Volatility

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

From

$$\frac{\partial^2 C_{BSM}}{\partial K^2} = \frac{\partial C_{BSM}}{\partial \Sigma} \frac{1}{\Sigma T K^2}$$

$$\frac{\partial^2 C_{BSM}}{\partial K \partial \Sigma} = \frac{\partial C_{BSM}}{\partial \Sigma} \frac{d_1}{\Sigma \sqrt{T} K}$$

and

$$\frac{\partial^2 C_{BSM}}{\partial \Sigma^2} = \frac{\partial C_{BSM}}{\partial \Sigma} \frac{d_1 d_2}{\Sigma}$$

we have

$$C_{KK} = \frac{C_{BSM}}{\partial \Sigma} \left(\frac{1}{\Sigma T K^2} + \frac{d_1}{\Sigma \sqrt{T} K} \frac{\partial \Sigma}{\partial K} + \frac{d_1 d_2}{\Sigma} \left(\frac{\partial \Sigma}{\partial K} \right)^2 + \frac{\partial^2 \Sigma}{\partial K^2} \right) \quad (22)$$

Local Volatility from Implied Volatility

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Substituting Equation (21) and (22) into Dupire's Equation we have

$$\sigma^2(T, K) = \frac{\frac{\Sigma}{2T} + \frac{\partial \Sigma}{\partial T} + rK \frac{\partial \Sigma}{\partial K}}{\frac{1}{2} K^2 \left(\frac{1}{\Sigma T K^2} + \frac{d_1}{\Sigma \sqrt{T} K} \frac{\partial \Sigma}{\partial K} + \frac{d_1 d_2}{\Sigma} \left(\frac{\partial \Sigma}{\partial K} \right)^2 + \frac{\partial^2 \Sigma}{\partial K^2} \right)} \quad (23)$$

Implied Variance as the Average of Local Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Implied volatility is often interpreted as the market expectation of the average of volatility throughout the life of an option. This is in general not true. However it can be justified when there is no skew.

Implied Variance as the Average of Local Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Assume that there is no skew, i.e., $\Sigma(T, K)$ does not depend on K . In this case, the local volatility σ does not depend on K either. From Equation (23), we have

$$\sigma^2(T) = \Sigma^2 + 2T\Sigma \frac{\partial \Sigma}{\partial T}$$

which implies

$$\Sigma^2(T) = \frac{1}{T} \int_0^T \sigma^2(t) dt$$

Implied Variance as the Average of Local Variance

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

This shows

- implied variance as the average of instantaneous variance for the life of option
- implied volatility is a global measure of volatility
- local volatility is a local measure of volatility for a particular pair of T and K .

Features of Local Volatility Model

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Local volatility model

- 1 A model to value European options in the market consistently.
- 2 A snapshot of current market rather than a model for the dynamics of volatility.
- 3 Smiles in local volatility model tend to flat as time advances.

Model Calibration

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

In practice the models we build usually come with parameters. The process of choosing model parameters to fit market prices (or volatility) is called calibration.

Model Calibration

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

For a local volatility model, calibration is about finding the local volatility function $\sigma(t, S_t)$.

Two common approaches

- Parametric Approach
- Non-parametric Approach
 - Match prices

$$C_T + rKC_K - \frac{1}{2}\sigma^2(T, K)K^2C_{KK} = 0 \quad (24)$$

- Match implied volatilities

$$\sigma^2(T, K) = \frac{\frac{\Sigma}{2T} + \frac{\partial \Sigma}{\partial T} + rK \frac{\partial \Sigma}{\partial K}}{\frac{1}{2}K^2 \left(\frac{1}{\Sigma T K^2} + \frac{d_1}{\Sigma \sqrt{T} K} \frac{\partial \Sigma}{\partial K} + \frac{d_1 d_2}{\Sigma} \left(\frac{\partial \Sigma}{\partial K} \right)^2 + \frac{\partial^2 \Sigma}{\partial K^2} \right)} \quad (25)$$

Parametric Approach

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

In the parametric approach, we assume the local volatility surface takes a particular functional form.

Examples

1 CEV model

$$\sigma(t, S_t) = \alpha S_t^{\beta-1}$$

2 Quadratic local volatility

$$\sigma(t, S_t) = \gamma(t) (a + bS_t + cS_t^2)$$

Parametric Approach

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

We will use a particular functional form for local volatility surface to show the whole process of model calibration.

We assume the local volatility function takes the form

$$\sigma(t, S_t) = a(t) + b(t)x + c(t)x^2$$

where x is the log-moneyness

$$x = \ln \left(\frac{S_t}{S_0} \right)$$

and $a(t)$, $b(t)$ and $c(t)$ are time-dependent functions.

Parametric Approach

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Suppose that we can obtain option prices for expiries $T_1 < T_2 < \dots < T_m$. And for each expiry T_i , there are n options with strikes $K_{i,j}, j = 1, \dots, n$.

K_{j+2}			$C(i, j+2)$	$C(i+1, j+2)$
K_{j+1}			$C(i, j+1)$	$C(i+1, j+1)$
K_j			$C(i, j)$	$C(i+1, j)$
K_{j-1}			$C(i, j-1)$	$C(i+1, j-1)$
	T_{i-1}	T_i	T_{i+1}	T_{i+2}

Parametric Approach

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

In practice we usually look at the discretized form of $a(t)$, $b(t)$ and $c(t)$. The common choices are piece-wise constant or piece-wise linear.

1 Piece-wise constant

$$a(t) = a_i, b(t) = b_i, c(t) = c_i, \forall t \in [T_{i-1}, T_i]$$

where a_i , b_i and c_i are constants to be determined

2 Piece-wise linear

$a(t)$, $b(t)$ and $c(t)$ are linear on each interval $[T_{i-1}, T_i]$

Parametric Approach

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

In this example, we assume $a(t)$, $b(t)$ and $c(t)$ are piece-wise constant functions.

The model parameters are a_i , b_i and c_i for $1 \leq i \leq m$.

We use a bootstrap process to solve these parameters.

Parametric Approach

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Suppose a_s , b_s and c_s are known for all $s = 1, \dots, i - 1$.

At i -th step, we solve

$$\min_{a_i, b_i, c_i} \sum_{j=1}^n \left(C^M(T_i, K_j) - C(T_i, K_j) \right)^2 \quad (26)$$

where $C^M(T_i, K_j)$ are model prices by solving Dupire's forward equation (24).

Parametric Approach

Local
Volatility
Model

Wu Lei


Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree



Parametric Approach

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

The minimization problem (26) is usually solved by an numerical iteration schema as illustrated below

Algorithm 1: Finding a_i, b_i and c_i

- 1 Initialize a_i, b_i, c_i
 - 2 **for** $s \leftarrow 1$ **to** S **do**
 - 3 Using Dupire's forward equation to solve $C^M(T_i, K_j)$
 for all $j = 1, \dots, n$
 - 4 **if** *converges* **then**
 - 5 **return** a_i, b_i, c_i
 - 6 try new values for a_i, b_i, c_i
 - 7 **return** a_i, b_i, c_i
-

Parametric Approach

Local
Volatility
Model

Wu Lei

Introduction


Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Once we have solved values for parameter a_i , b_i and c_i , we can move to next layer to solve a_{i+1} , b_{i+1} and c_{i+1} .



Parametric Approach

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

We can repeat this process until we reach the final (m -th) layer.

$$a_1, b_1, c_1 \rightarrow \dots \rightarrow a_m, b_m, c_m$$

Non-parametric Approach

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

The non-parametric approach does not assume a functional form for the local volatility surface. Instead it aims to recover the whole local volatility surface (in practice a fine grid of strikes and expiries).

There are several ways to do this. We present two of them

- Optimization based
- Interpolation/extrapolation based

Non-parametric Approach

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

In the optimization based method, we typically have a grid for strike and expiry space. Let $\sigma_{i,j} = \sigma(T_i, K_j)$ be the local volatility for the point (T_i, K_j) on the grid. The calibration becomes the following minimization problem

$$\min_{\sigma_{i,j}} \sum_{i,j} w_{i,j} \left(C^M(T_i, K_j) - C(T_i, K_j) \right)^2$$

where $w_{i,j}$ are weights.

Note that we attempt to solve all $\sigma_{i,j}$ on one go.

Non-parametric Approach

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

This minimization problem is often ill-posed since the number of parameters $\sigma_{i,j}$ are usually larger than the number of options in the market. Hence the solution is not stable. A small perturbation in the input data will result in very different results.

Non-parametric Approach

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

To stabilize the calibration, regularization methods are usually employed. One popular method is Tikhonov regularization. For example, the second order regularization is

$$\min_{\sigma_{i,j}} \sum_{i,j} w_{i,j} \left(C^M(T_i, K_j) - C(T_i, K_j) \right)^2 + \lambda \left\| \frac{\partial^2 \sigma}{\partial s^2} + \frac{\partial^2 \sigma}{\partial s \partial t} + \frac{\partial^2 \sigma}{\partial t^2} \right\|$$

where $\| \cdot \|$ is usually L_2 norm.

Non-parametric Approach

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Note that in this approach, $\sigma_{i,j}$ is discrete. In case local volatility is needed for a point not on the grid, interpolation or extrapolation will be needed.

Non-parametric Approach

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Another popular non-parametric approach is to use the implied volatility formulation of Dupire's equation (25). In this approach, we interpolate/extrapolate an implied volatility surface. The local volatilities can then be calculated from Equation (25).

$$IVS \implies LVS$$

Non-parametric Approach

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

The implied volatility surface can be constructed in various ways (see for example Borovkova & Permanab (2009)).

Examples:

- Parametric

$$\sigma(x, T) = b_1 + b_2x + b_3x^2 + b_4T + b_5xT$$

where $x = \ln\left(\frac{S_t}{S_0}\right) / T$.

- Semi-parametric

$$\sigma(x, T) = b_{1,T} + b_{2,T}x + b_{3,T}x^2$$

- Non-parametric
Cubic spline and its variant etc.

Comparison Between Different Approaches

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

- Parametric approach usually gives more smooth local volatility surface. However in the case the number of parameters are larger than available market option prices. There may be an identifiability issue.
- The optimization based non-parametric approach is usually a high dimension optimization problem and slow to solve.
- It is easy to implement the implied volatility approach. However the extrapolation becomes difficult for strikes far from ATM. There is no guarantee that the local variance is positive.

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Binomial tree model is often used as a prototype model. How do we incorporate smile in binomial tree?

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction


Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Let $\Delta t = \frac{T}{N}$ and $t_n = n\Delta t$. We build a binomial tree as below



- The node (n, i) corresponds to the i -th stock price level at time t_n
- At the node (n, i) , the (implied risk neutral) probability of the stock price moves up to $S_{n+1,i+1}$ is $p_{n,i}$, and moves down to $S_{n+1,i}$ is $1 - p_{n,i}$.
- The initial price $S_{1,1} = S_0$.

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction


Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Structure of implied binomial tree



Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction


Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Structure of implied binomial tree for interior nodes



Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction


Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

For BSM model, the binomial tree is homogeneous where the price goes up (or down) by a factor of u (or d) and the transition probability is p for all nodes.



$$u > 1, d < 1, ud = 1$$

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction


Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

The implied binomial tree is inhomogeneous as we need to adjust S_u , S_d and p for each node to match market prices of options.



Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

In binomial tree, at each node, we choose stock price S_u , S_d and transition probability p for the next period to match the first two moments of the stock price dynamics.

In implied binomial tree, for each node, we choose S_u , S_d and p to match the first moment and option prices.

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Let $\lambda_{n,i}$ be the risk neutral probability that the stock price reaches to the node (n, i) .

Note that $e^{-rt_n} \lambda_{n,i}$ is the Arrow-Debreu price of a derivative that pays out 1 if the stock price is $S_{n,i}$ at time t_n and zero otherwise.

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction


Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

From



we have

$$\lambda_{n+1,i+1} = \lambda_{n,i+1}(1 - p_{n,i+1}) + \lambda_{n,i}p_{n,i} \quad (27)$$

where $1 \leq i \leq n - 1$ for interior nodes.

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

For boundary nodes, we have

$$\lambda_{n+1,n+1} = \lambda_{n,n} p_{n,n} \quad (28)$$

and

$$\lambda_{n+1,1} = \lambda_{n,1} (1 - p_{n,1}) \quad (29)$$

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Together with the initial condition that $\lambda_{1,1} = 1$, Equation (27), (28) and (29) provide an iterative schema to compute $\lambda_{n,i}$ provided that we know $p_{n,i}$.

We compute $p_{n,i}$ and $S_{n,i}$ by forward induction.

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Suppose we have obtained values of $\lambda_{n,i}$, $p_{n,i}$ and $S_{n,i}$ for all i at n -th level. To move to level $n + 1$, what are the parameters we need to compute?

Implied Binomial Tree

Local
Volatility
Model

Wu Lei


Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree



- Transition probability $p_{n,i}$ for $1 \leq i \leq n$
- $S_{n+1,i}$ for $1 \leq i \leq n+1$.

There are a total of $2n+1$ parameters.

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

We impose two constraints

- Match the first moment
- Match the market prices of call/put options

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

To match the first moment, we must have

$$S_{n,i} = e^{-r\Delta t} (p_{n,i} S_{n+1,i+1} + (1 - p_{n,i}) S_{n+1,i})$$

for $i = 1, \dots, n$.

If we denote $F_{n,i} = e^{r\Delta t} S_{n,i}$, the constraint can be written as

$$F_{n,i} = p_{n,i} S_{n+1,i+1} + (1 - p_{n,i}) S_{n+1,i}$$

Solving the above equation for $p_{n,i}$, we have

$$p_{n,i} = \frac{F_{n,i} - S_{n+1,i}}{S_{n+1,i+1} - S_{n+1,i}} \quad (30)$$

Note that $S_{n+1,i}$ and $S_{n+1,i+1}$ are unknown.

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Let $C(T, K)$ be the undiscounted price of a call option with strike K and expiry T . Using risk neutral pricing formula, we have

$$C(t_{n+1}, K) = \sum_{j=1}^{n+1} \lambda_{n+1,j} (S_{n+1,j} - K)^+$$

Note that the summation is over all the nodes at time t_{n+1}

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Note that the price of a call option with strike $S_{n,i}$ and expiry t_{n+1} can reduce to


$$C(t_{n+1}, S_{n,i}) = \sum_{j=i+1}^{n+1} \lambda_{n+1,j} (S_{n+1,j} - S_{n,i})$$

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

This is illustrated in the figure below



Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Since

$$\lambda_{n+1,j} = p_{n,j-1}\lambda_{n,j} + (1 - p_{n,j})\lambda_{n,j}$$

we have

$$C(t_{n+1}, S_{n,i}) = \sum_{j=i+1}^{n+1} (p_{n,j-1}\lambda_{n,j} + (1 - p_{n,j})\lambda_{n,j}) (S_{n+1,j} - S_{n,i}) \quad (31)$$

with the convention that $\lambda_{n,n+1} = 0$.

Implied Binomial Tree

Local
Volatility
Model

Wu Lei


Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree



Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction


Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

We can reorganize the terms in the summation as below



Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Note that

$$\begin{aligned} & \lambda_{n,j} p_{n,j} (S_{n+1,j+1} - S_{n,i}) + \lambda_{n,j} (1 - p_{n,j}) (S_{n+1,j} - S_{n,i}) \\ = & \lambda_{n,j} [(p_{n,j} S_{n+1,j+1} + (1 - p_{n,j}) S_{n+1,j}) - S_{n,i}] \\ = & \lambda_{n,j} (F_{n,j} - S_{n,i}) \end{aligned}$$

Using this to simplify the Equation (31), we have

$$C(t_{n+1}, S_{n,i}) = \lambda_{n,i} p_{n,i} (S_{n+1,i+1} - S_{n,i}) + \sum_{j=i+1}^n \lambda_{n,j} (F_{n,j} - S_{n,i}) \quad (32)$$

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

Combing it with Equation (30), we can solve $S_{n+1,i+1}$ as a function of $S_{n+1,i}$

$$= \frac{S_{n+1,i+1}}{S_{n+1,i} \left[C(t_{n+1}, S_{n,i}) - \sum_{j=i+1}^n \lambda_{n,j} (F_{n,j} - S_{n,i}) \right] - S_{n,i} \lambda_{n,i} (F_{n,i} - S_{n+1,i})} \\ \left[C(t_{n+1}, S_{n,i}) - \sum_{j=i+1}^n \lambda_{n,j} (F_{n,j} - S_{n,i}) \right] - \lambda_{n,i} (F_{n,i} - S_{n+1,i})$$

Note that $C(t_{n+1}, S_{n,i})$ is obtained from market and $\lambda_{n,j}, (F_{n,j}$ and $S_{n,i}$ are known by induction, hence if we know the value for $S_{n+1,i}$, we can compute $S_{n+1,i+1}$.

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

If we know $S_{n+1,i}$ for some i , then we know the value of $S_{n+1,j}$ for all $j \geq i$

$$S_{n+1,i} \longrightarrow S_{n+1,i+1} \longrightarrow \dots \longrightarrow S_{n+1,n+1}$$

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

We choose i to be around the center

- If $n + 1$ is odd, we choose $i = \frac{n+2}{2}$

$$S_{n+1,i} = S_0$$

- If $n + 1$ is even, we choose $i = \frac{n+1}{2}$ and let

$$S_{n+1,i} = \frac{S_0^2}{S_{n+1,i+1}}$$

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

For the nodes below the center node, we can similarly use put options to compute $S_{n+1,i}$ and $p_{n,i}$, using the formula

$$S_{n+1,i} = \frac{S_{n+1,i+1} [P(t_{n+1}, S_{n,i}) - \sum_{j=i+1}^n \lambda_{n,j} (S_{n,i} - F_{n,j})] - S_{n,i} \lambda_{n,i} (F_{n,i} - S_{n+1,i+1})}{[P(t_{n+1}, S_{n,i}) - \sum_{j=i+1}^n \lambda_{n,j} (S_{n,i} - F_{n,j})] + \lambda_{n,i} (F_{n,i} - S_{n+1,i})}$$

We can solve $S_{n+1,i}$ in the reverse order

$$S_{n+1,i} \longrightarrow S_{n+1,i-1} \longrightarrow \dots \longrightarrow S_{n+1,1}$$

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

To avoid arbitrage, we must have

$$0 \leq p_{n,i} \leq 1$$

Hence the stock price must satisfy

$$F_{n,i} \leq S_{n+1,i+1} \leq F_{n,i+1}$$

In case the resulting prices from the above process don't satisfy this condition, we can may choose the prices as

$$\ln \left(\frac{S_{n+1,i+1}}{S_{n+1,i}} \right) = \ln \left(\frac{S_{n,i}}{S_{n,i-1}} \right)$$

or

$$S_{n+1,i+1} = \frac{F_{n,i} + F_{n,i+1}}{2}$$

This does not guarantee the absence of negative probability.

Implied Binomial Tree

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree

What is the local volatility for the node (n, i) ?

$$\sigma^2(t_n, S_{n,i})\Delta t = p_{n,i}(1 - p_{n,i}) \ln \left(\frac{S_{n+1,i+1}}{S_{n+1,i}} \right)$$

This can be obtained by using the fact

$$\text{Var} [\Delta \ln S(t) | \mathcal{F}_t] \approx \sigma^2(t, S_t)\Delta t$$

References

Local
Volatility
Model

Wu Lei

Introduction

Dupire's
Equation

Understanding
Local
Volatility

How to Build
Local
Volatility
Surface

Implied
Binomial Tree



Dupire (1994)

Pricing with a Smile, *Risk*, 7(1): 18-20



Derman, E., Iraj Kani (1994)

"Riding on a Smile." *RISK*, 7(2): 32-39



Matthias R. Fengler (2005)

Semiparametric Modeling of Implied Volatility, *Springer*



Lorenzo Bergomi (2016)

Stochastic Volatility Modeling, *Chapman & Hall/CRC Financial Mathematics Series*



Svetlana Borovkova, Ferry J. Permanab (2009)

Implied volatility in oil markets, *Computational Statistics and Data Analysis* 53 (2009) 2022-2039

Thank you!