

FE5222 Advanced Derivative Pricing

Wu Lei

Risk Management Institute, National University of Singapore

rmiwul@nus.edu.sg

August 26, 2019

Overview

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

1 Probability

2 Stochastic Process

3 Stochastic Calculus

Probability Space

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

A probability space is a triplet of $(\Omega, \mathcal{F}, \mathbb{P})$, where

- Ω is the sample space,
- \mathcal{F} is a σ -algebra, and
- \mathbb{P} is a probability measure.

Sample Space

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

A sample space Ω is the set of all possible outcomes. Each $\omega \in \Omega$ represents an outcome.

Example

- Toss a coin once, $\Omega = \{H, T\}$

Sample Space

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

A sample space Ω is the set of all possible outcomes. Each $\omega \in \Omega$ represents an outcome.

Example

- Toss a coin once, $\Omega = \{H, T\}$
- Toss a coin twice, $\Omega = \{HH, HT, TH, TT\}$

Sample Space

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

A sample space Ω is the set of all possible outcomes. Each $\omega \in \Omega$ represents an outcome.

Example

- Toss a coin once, $\Omega = \{H, T\}$
- Toss a coin twice, $\Omega = \{HH, HT, TH, TT\}$
- Toss a coin infinite times? $\Omega = [0, 1)$

Definition

A σ -algebra (σ -field) \mathcal{F} is a family of subsets in Ω such that

- $\Omega \in \mathcal{F}$,
- If $E \in \mathcal{F}$, then $E^c \in \mathcal{F}$, and
- If $E_n \in \mathcal{F}$, $n = 1, 2, \dots$, then $\bigcup_{n=1}^{\infty} E_n \in \mathcal{F}$.

Definition

A σ -algebra (σ -field) \mathcal{F} is a family of subsets in Ω such that

- $\Omega \in \mathcal{F}$,
- If $E \in \mathcal{F}$, then $E^c \in \mathcal{F}$, and
- If $E_n \in \mathcal{F}$, $n = 1, 2, \dots$, then $\bigcup_{n=1}^{\infty} E_n \in \mathcal{F}$.

Definition

A set $E \in \mathcal{F}$ is called an event.

Borel σ -algebra on \mathbf{R}^n

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

The Borel σ -algebra $\mathcal{B}(\mathbf{R}^n)$ the smallest σ -algebra containing all open sets in \mathbf{R}^n .

Borel σ -algebra on \mathbf{R}^n

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

The σ -algebra $\mathcal{B}(\mathbf{R}^n)$ includes all subsets in \mathbf{R}^n of our interest.
For example, the Borel σ -algebra $\mathcal{B}(\mathbf{R})$ includes

- Open set
- Close set
- Singleton: $\{a\}$
- Half-open (closed) sets: $(a, b], [a, b)$

σ -algebra

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Since the power set $2^\Omega = \text{all subsets of } \Omega$ is a σ -algebra, why do we need other σ -algebras?

σ -algebra

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Since the power set $2^\Omega = \text{all subsets of } \Omega$ is a σ -algebra, why do we need other σ -algebras?

1 Overcome technical difficulties

The σ -algebra 2^Ω is too big to define a meaningful probability measure in some cases.

Since the power set $2^\Omega = \text{all subsets of } \Omega$ is a σ -algebra, why do we need other σ -algebras?

1 Overcome technical difficulties

The σ -algebra 2^Ω is too big to define a meaningful probability measure in some cases.

2 Model information

More importantly we can use σ -algebra to model information.

σ -algebra

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

The outcome $\omega \in \Omega$ is what has really happened. This may not be observable by a person. Each $E \in \mathcal{F}$ is a set of ω 's in Ω . If our information is \mathcal{F} , then we know whether a particular E has happened or not. We may know one of the ω 's in E has happened, but don't know exactly which ω .

Example

In the example of tossing a coin twice, the sample space is $\Omega = \{HH, HT, TH, TT\}$, we can have the following two different σ -algebras.

- $\mathcal{F}_1 = \{\emptyset, \{HT, HH\}, \{TH, TT\}, \Omega\}$
- $\mathcal{F}_2 = \{\emptyset, \{HT, HH, TH\}, \{TT\}, \Omega\}$

σ -algebra

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

If $\omega = HT$, a person possessing the information \mathcal{F}_1 knows that the event $\{HT, HH\}$ (i.e., the first toss is head) has happened. But he or she does not know whether it is HT or HH . This can happen if this person only observes the first coin toss.

σ -algebra

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Similarly, a person possessing the information \mathcal{F}_2 knows that the event $\{HT, HH, TH\}$ (i.e., at least the one toss is head) has happened. But he or she does not know whether it is HT , HH or TH .

Probability Measure

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

A probability measure \mathbb{P} is a mapping from \mathcal{F} to $[0, 1]$ such that

- $\mathbb{P}(\emptyset) = 0$,
- $\mathbb{P}(\Omega) = 1$, and
- $\mathbb{P}(\cup_{n=1}^{\infty} E_n) = \sum_{n=1}^{\infty} \mathbb{P}(E_n)$, where $E_i \cap E_j = \emptyset \ \forall i \neq j$.

Probability Measure

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

A probability measure \mathbb{P} is a mapping from \mathcal{F} to $[0, 1]$ such that

- $\mathbb{P}(\emptyset) = 0$,
- $\mathbb{P}(\Omega) = 1$, and
- $\mathbb{P}(\cup_{n=1}^{\infty} E_n) = \sum_{n=1}^{\infty} \mathbb{P}(E_n)$, where $E_i \cap E_j = \emptyset \ \forall i \neq j$.

Note that a probability measure is only defined on the σ -algebra. We cannot assign a probability to the set $E \notin \mathcal{F}$.

Almost Surely

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $E \in \mathcal{F}$ and $\mathbb{P}(E) = 1$. Then we say the event E happens almost surely (a.s.).

Note that E does not always happen. The event it does not happen is insignificant in probabilistic sense.

Random Variable

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

A random variable (r.v.) X is a real-valued¹ function on Ω such that

$$X^{-1}(B) \in \mathcal{F}$$

for all $B \in \mathcal{B}(\mathbf{R})$, where

$$X^{-1}(B) = \{\omega \in \Omega : X(\omega) \in B\}$$

¹In some cases, we shall allow X to take ∞ or $-\infty$.

Random Variable

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Note that

- The requirement for $X^{-1}(B)$ to be \mathcal{F} -measurable is to ensure we can assign probability to the events we are interested in. For example, if X is the stock price, we would like to ask what is the probability that X is between 100 and 120 tomorrow. To assign the probability to such an event, we must have

$$\{\omega : 100 \leq X(\omega) \leq 120\} \in \mathcal{F}$$

.

Random Variable

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Note that

- The requirement for $X^{-1}(B)$ to be \mathcal{F} -measurable is to ensure we can assign probability to the events we are interested in. For example, if X is the stock price, we would like to ask what is the probability that X is between 100 and 120 tomorrow. To assign the probability to such an event, we must have

$$\{\omega : 100 \leq X(\omega) \leq 120\} \in \mathcal{F}$$

.

- If X is a random variable, f is a Borel function, then $Y = f(X)$ is also a random variable.

Information from a Random Variable

Mathematical
Foundation



Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Suppose we don't observe $\omega \in \Omega$, but we know the values of the random variable X_1 or X_2 .



Information from a Random Variable

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

A person who knows X_1 can tell whether $\{HH, HT, TH\}$ or $\{TT\}$ happens by observing the values of X_1 . Similarly a person who knows X_2 can tell between the event $\{HH, HT\}$ and $\{TH, TT\}$. Hence X_1 and X_2 convey different information!

Information from a Random Variable

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let X be a random variable, the σ -algebra generated by X is defined as

$$\sigma(X) = \{X^{-1}(B) : B \in \mathcal{B}(\mathbb{R})\}$$

which is the smallest σ -algebra such that X is $\sigma(X)$ -measurable.

Information from a Random Variable

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Note that

- The σ -algebra $\sigma(X)$ is the information contained in the random variable X .

Information from a Random Variable

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Note that

- The σ -algebra $\sigma(X)$ is the information contained in the random variable X .
- In the above example, a person knows the values of both X_1 and X_2 will have more information. For example, if he/she observes $X_1(\omega) = 1$ and $X_2(\omega) = 2$, he/she can immediately deduce that $\omega = TH$.

Information from Random Variables

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $X_\lambda, \lambda \in \Lambda$ be random variables, the σ -algebra generated by X_λ , denoted by $\sigma(X_\lambda, \lambda \in \Lambda)$, is the smallest σ -algebra that contains $\sigma(X_\lambda)$ for all $\lambda \in \Lambda$.

Information from Random Variables

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $X_\lambda, \lambda \in \Lambda$ be random variables, the σ -algebra generated by X_λ , denoted by $\sigma(X_\lambda, \lambda \in \Lambda)$, is the smallest σ -algebra that contains $\sigma(X_\lambda)$ for all $\lambda \in \Lambda$.

$\sigma(X_\lambda, \lambda \in \Lambda)$ can be interpreted as the information contained in all X_λ .

Information from Random Variables

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, $\mathcal{G} \subset \mathcal{F}$ be a sub σ -algebra of \mathcal{F} . X is a random variable. If $\sigma(X) \subset \mathcal{G}$, then we say X is \mathcal{G} -measurable.

Information from Random Variables

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, $\mathcal{G} \subset \mathcal{F}$ be a sub σ -algebra of \mathcal{F} . X is a random variable. If $\sigma(X) \subset \mathcal{G}$, then we say X is \mathcal{G} -measurable.

$\sigma(X)$ is the information contained in X . A person with the knowledge of \mathcal{G} knows whether an event E happens or not for any $E \in \mathcal{G}$. Since $\sigma(X) \subset \mathcal{G}$. The person knows whether $X^{-1}(B)$ happens or not for all $B \in \mathcal{B}(R)$.

Information from Random Variables

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem

Let X and Y be two random variables. If X is $\sigma(Y)$ -measurable (i.e. $\sigma(X) \subset \sigma(Y)$), then there is a Borel function f such that $X(\omega) = f(Y(\omega))$.

Convergence

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $X_n, n = 1, 2, \dots$ be a sequence of random variables, X be a random variable, X_n is said to converge to X almost surely (a.s.) if

$$\lim_{n \rightarrow \infty} X_n(\omega) = X(\omega)$$

for all $\omega \in \Omega$ except on a set $N \subset \Omega$ with $\mathbb{P}(N) = 0$.

Convergence in Probability

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $X_n, n = 1, 2, \dots$ be a sequence of random variables, X be a random variable, X_n is said to converge to X in probability if for any $\delta > 0$,

$$\lim_{n \rightarrow \infty} \mathbb{P}(|X_n(\omega) - X(\omega)| > \delta) = 0$$

Convergence of Expectation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem (Monotone Convergence Theorem)

Let $0 \leq X_1 \leq X_n \leq \dots$ and $\lim_{n \rightarrow \infty} X_n = X$, then

$$\lim_{n \rightarrow \infty} \mathbb{E}(X_n) = \mathbb{E}(X).$$

Note that

- 1 $X(\omega)$ may take value $+\infty$ at some ω
- 2 $\mathbb{E}(X)$ may be $+\infty$

Convergence of Expectation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem (Dominated Convergence Theorem)

Let $|X_n| \leq Y$ for all $n = 1, 2, \dots$, $\lim X_n = X$ and $\mathbb{E}(Y) < \infty$, then

$$\lim \mathbb{E}(X_n) = \mathbb{E}(X).$$

Convergence of Expectation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Pay attention to the conditions under which the two convergence theorems hold!

- For the Monotone Convergence Theorem, X_n are assumed to be non-negative.
- For the Dominated Convergence Theorem, X_n are bounded by an integrable random variable Y .

Change of Measure

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

A probability measure essentially defines a weight for each $\omega \in \Omega$. We can scale the weights provided that they still sum up to 1. This process is called *change of measure*.

Change of Measure

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $Z(\omega)$ be a non-negative random variable such that $\mathbb{E}[Z] = 1$. Define

$$\tilde{\mathbb{P}}(E) = \int_E Z(\omega) d\mathbb{P}(\omega)$$

for all $E \in \mathcal{F}$. Then $\tilde{\mathbb{P}}$ is a probability measure.

Change of Measure

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem

Let $\tilde{\mathbb{E}}[\cdot]$ denote the expectation under $\tilde{\mathbb{P}}$, we have

$$\tilde{\mathbb{E}}[Y] = \mathbb{E}[YZ]$$

provided that both expectations exist.

Change of Measure

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem

Let $\tilde{\mathbb{E}}[\cdot]$ denote the expectation under $\tilde{\mathbb{P}}$, we have

$$\tilde{\mathbb{E}}[Y] = \mathbb{E}[YZ]$$

provided that both expectations exist.

An intuitive proof.

$$\begin{aligned}\tilde{\mathbb{E}}[Y] &= \int Y(\omega) d\tilde{\mathbb{P}} \\ &= \sum_{\omega} Y(\omega) \tilde{\mathbb{P}}(\omega) \\ &= \sum_{\omega} Y(\omega) Z(\omega) \mathbb{P}(\omega) \\ &= \int Y(\omega) Z(\omega) d\mathbb{P}(\omega) \\ &= \mathbb{E}[YZ]\end{aligned}$$



Equivalent Probability Measures

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let \mathbb{P} and $\tilde{\mathbb{P}}$ be two probability measures on (Ω, \mathcal{F}) . They are said to be equivalent if

$$\mathbb{P}(E) = 0 \iff \tilde{\mathbb{P}}(E) = 0$$

or equivalently

$$\mathbb{P}(E) = 1 \iff \tilde{\mathbb{P}}(E) = 1$$

Equivalent Probability Measures

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

In the risk-neutral pricing approach, we price derivatives under the risk-neutral measure $\tilde{\mathbb{P}}$. This will require $\tilde{\mathbb{P}}$ to be equivalent to the real-world measure \mathbb{P} so that no arbitrage in one world implies no arbitrage in the other world.

Equivalent Probability Measures

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

In the risk-neutral pricing approach, we price derivatives under the risk-neutral measure $\tilde{\mathbb{P}}$. This will require $\tilde{\mathbb{P}}$ to be equivalent to the real-world measure \mathbb{P} so that no arbitrage in one world implies no arbitrage in the other world.

No arbitrage in real world

$$\iff \mathbb{P}(\text{arbitrage}) = 0$$

$$\iff \tilde{\mathbb{P}}(\text{arbitrage}) = 0$$

$$\iff \text{No arbitrage in risk-neutral world}$$

Radon-Nikodym

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem

Let \mathbb{P} and $\tilde{\mathbb{P}}$ be two equivalent probability measures on (Ω, \mathcal{F}) , then there exists an a.s. positive random variable Z such that

$$\mathbb{P}(E) = \int Z(\omega) d\tilde{\mathbb{P}}(\omega)$$

for any $E \in \mathcal{F}$. We usually denote Z as $\frac{d\mathbb{P}}{d\tilde{\mathbb{P}}}$.

Independence

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, $\mathcal{G}_i \subset \mathcal{F}, i = 1, \dots, n$, be sub σ -algebras. \mathcal{G}_i are (mutually) independent if

$$\mathbb{P}(A_1 \cap \dots \cap A_n) = \mathbb{P}(A_1) \dots \mathbb{P}(A_n)$$

for all $A_i \in \mathcal{G}_i$.

Independence

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Note that

- Independence is not equivalent to pairwise independence, which requires

$$\mathbb{P}(A_i \cap A_j) = \mathbb{P}(A_i)\mathbb{P}(A_j)$$

for all $A_i \in \mathcal{G}_i$ and $A_j \in \mathcal{G}_j, i \neq j$.

Independence

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Note that

- Independence is not equivalent to pairwise independence, which requires

$$\mathbb{P}(A_i \cap A_j) = \mathbb{P}(A_i)\mathbb{P}(A_j)$$

for all $A_i \in \mathcal{G}_i$ and $A_j \in \mathcal{G}_j, i \neq j$.

- Since we can take $A_i = \Omega$, the definition of independence is equivalent to

$$\mathbb{P}(A_{i_1} \cap \dots \cap A_{i_m}) = \mathbb{P}(A_{i_1}) \dots \mathbb{P}(A_{i_m})$$

for all $A_{i_j} \in \mathcal{G}_{i_j}, j = 1, \dots, m, m \leq n$.

Independence

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Two random variables X and Y are independent if $\sigma(X)$ and $\sigma(Y)$ are independent.

Independence

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem

Let X and Y be two random variables. Their joint moment generating function is

$$M_{X,Y}(s, t) = \mathbb{E}[e^{sX+tY}]$$

Let

$$M_X(s) = \mathbb{E}[e^{sX}]$$

and

$$M_Y(t) = \mathbb{E}[e^{tY}]$$

be the moment generating function for X and Y respectively. Then X and Y are independent if and only if

$$M_{X,Y}(s, t) = M_X(s)M_Y(t)$$

Conditional Expectation


Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus



Conditional Expectation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Suppose we have the information

$\mathcal{F}_1 = \{\emptyset, \Omega, \{HH, HT\}, \{TH, TT\}\}$. Let $\omega = HH$, what will be our estimate of expected values for X ?

Conditional Expectation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Suppose we have the information

$\mathcal{F}_1 = \{\emptyset, \Omega, \{HH, HT\}, \{TH, TT\}\}$. Let $\omega = HH$, what will be our estimate of expected values for X ?

Answer: Since our information is \mathcal{F}_1 , given $\omega = HH$, we can only know that the event $\{HH, HT\}$ has happened. But we don't know whether it is HH or HT . Hence our best estimate of X is $\frac{1+3}{2} = 2$.

Conditional Expectation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

And what if $\omega = TH$?

Conditional Expectation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

And what if $\omega = TH$?

Given our information \mathcal{F}_1 , we know $\omega = TH$ or TT . In either case, the value of X is 1. Hence our estimate will be 1.

Conditional Expectation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

In summary, our estimate of X based on the information \mathcal{F}_1 is

$$Y(\omega) = \begin{cases} 2 & \text{if } \omega \in \{HH, HT\} \\ 1 & \text{if } \omega \in \{TH, TT\} \end{cases}$$

Conditional Expectation


Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus



Conditional Expectation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Note that

- Y is a random variable
- Y is \mathcal{F}_1 -measurable
- Y has the same expectation as X on set $\{HH, HT\}$ and $\{TH, TT\}$.

Conditional Expectation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let X be an integrable random variable, $\mathcal{G} \subset \mathcal{F}$ be a sub σ -algebra. A random variable Y is the conditional expectation of X given \mathcal{G} if

1 Y is \mathcal{G} -measurable

2

$$\int_A Y(\omega) d\mathbb{P}(\omega) = \int_A X(\omega) d\mathbb{P}(\omega)$$

for all $A \in \mathcal{G}$.

We denote $Y = \mathbb{E}[X|\mathcal{G}]$.

Conditional Expectation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Properties

1 If $\mathcal{G} = \{\emptyset, \Omega\}$, then $\mathbb{E}[X|\mathcal{G}] = \mathbb{E}[X]$.

Conditional Expectation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Properties

- 1 If $\mathcal{G} = \{\emptyset, \Omega\}$, then $\mathbb{E}[X|\mathcal{G}] = \mathbb{E}[X]$.
- 2 If Y is \mathcal{G} -measurable, then $\mathbb{E}[YX|\mathcal{G}] = Y\mathbb{E}[X|\mathcal{G}]$.

Conditional Expectation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Properties

- 1 If $\mathcal{G} = \{\emptyset, \Omega\}$, then $\mathbb{E}[X|\mathcal{G}] = \mathbb{E}[X]$.
- 2 If Y is \mathcal{G} -measurable, then $\mathbb{E}[YX|\mathcal{G}] = Y\mathbb{E}[X|\mathcal{G}]$.
- 3 If $\mathcal{H} \subset \mathcal{G}$, then $\mathbb{E}[\mathbb{E}[X|\mathcal{G}]|\mathcal{H}] = \mathbb{E}[X|\mathcal{H}]$

Conditional Expectation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Properties

- 1 If $\mathcal{G} = \{\emptyset, \Omega\}$, then $\mathbb{E}[X|\mathcal{G}] = \mathbb{E}[X]$.
- 2 If Y is \mathcal{G} -measurable, then $\mathbb{E}[YX|\mathcal{G}] = Y\mathbb{E}[X|\mathcal{G}]$.
- 3 If $\mathcal{H} \subset \mathcal{G}$, then $\mathbb{E}[\mathbb{E}[X|\mathcal{G}]|\mathcal{H}] = \mathbb{E}[X|\mathcal{H}]$
- 4 If X is independent of \mathcal{G} , then $\mathbb{E}[X|\mathcal{G}] = \mathbb{E}[X]$

Independence and Conditional Expectation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, $\mathcal{G} \subset \mathcal{F}$ be a sub σ -algebra. Suppose the random variables X_1, \dots, X_m are \mathcal{G} -measurable and Y_1, \dots, Y_n are independent of \mathcal{G} . $f(x_1, \dots, x_m, y_1, \dots, y_n)$ is a Borel function. Define

$$g(x_1, \dots, x_m) = \mathbb{E}[f(x_1, \dots, x_m, Y_1, \dots, Y_n)]$$

Then

$$\mathbb{E}[f(X_1, \dots, X_m, Y_1, \dots, Y_n) | \mathcal{G}] = g(X_1, \dots, X_m)$$

Stochastic Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

A stochastic process $X(t, \omega)$ is a function from $[0, \infty) \times \Omega$ to \mathbf{R}^n .

2

²Strictly speaking we need to impose some sort of measurability in the definition, but we ignore it here.

Stochastic Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

In the definition t is interpreted as time. Fix t , $X(t, \omega)$ is a random variable. Hence a stochastic process $X(t, \omega)$ can be interpreted as random variables that evolve with time. For example, it can be stock price.

Stochastic Process

Mathematical
Foundation


Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

We can also interpret X in another way. Fix ω , $X(\cdot, \omega)$ is a function on $[0, \infty)$. Hence X is a function from Ω to the space of all functions on $[0, \infty)$. This function is called a sample path of X .



Stochastic Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

In the second interpretation, an $\omega \in \Omega$ has been picked. However we don't know exactly ω . Our observation is the value of $X(t, \omega)$ up to time t . There could be many ω 's that will have the same sample path up to time t . Based on our observation, we can differentiate ω 's whose sample paths up to time t are different from what we have observed, but not the others.

Stochastic Process


Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus



Filtration

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

We model the accumulation of information with filtration.

Definition

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, $\{\mathcal{F}_t\}_{t \geq 0}$ be a collection of sub σ -algebra of \mathcal{F} . $\{\mathcal{F}_t\}_{t \geq 0}$ is a filtration if

$$\mathcal{F}_s \subset \mathcal{F}_t, \quad \forall s \leq t$$

Adapted Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $\{\mathcal{F}_t\}_{t \geq 0}$ be a filtration. A stochastic process X is said to be adapted to this filtration if $X(t)$ is \mathcal{F}_t -measurable.

Filtration from a Stochastic Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let X be a stochastic process, we can define

$$\mathcal{F}_t^X = \sigma(X_s, 0 \leq s \leq t)$$

Then

- $\{\mathcal{F}_t^X\}_{t \geq 0}$ is a filtration
- X_t is adapted to $\{\mathcal{F}_t^X\}_{t \geq 0}$

$\{\mathcal{F}_t^X\}_{t \geq 0}$ is called the filtration generated from X .

\mathcal{F}_t^X is the information one has by observing the value of X up to time t .

Martingale

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $\{\mathcal{F}_t\}_{t \geq 0}$ be a filtration on it. A stochastic process $M(t)$ is a martingale if

- $\mathbb{E}[|M(t)|] < \infty$
- $\mathbb{E}[M(t)|\mathcal{F}_s] = M(s)$ for all $s < t$.

Martingale

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $\{\mathcal{F}_t\}_{t \geq 0}$ be a filtration on it. A stochastic process $M(t)$ is a martingale if

- $\mathbb{E}[|M(t)|] < \infty$
- $\mathbb{E}[M(t)|\mathcal{F}_s] = M(s)$ for all $s < t$.

Note that

- In some sense, a martingale represents a fair game

Martingale

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $\{\mathcal{F}_t\}_{t \geq 0}$ be a filtration on it. A stochastic process $M(t)$ is a martingale if

- $\mathbb{E}[|M(t)|] < \infty$
- $\mathbb{E}[M(t)|\mathcal{F}_s] = M(s)$ for all $s < t$.

Note that

- In some sense, a martingale represents a fair game
- $M(t) = \mathbb{E}[X|\mathcal{F}_t]$ is a martingale where X is an integrable random variable

Martingale

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $\{\mathcal{F}_t\}_{t \geq 0}$ be a filtration on it. A stochastic process $M(t)$ is a martingale if

- $\mathbb{E}[|M(t)|] < \infty$
- $\mathbb{E}[M(t)|\mathcal{F}_s] = M(s)$ for all $s < t$.

Note that

- In some sense, a martingale represents a fair game
- $M(t) = \mathbb{E}[X|\mathcal{F}_t]$ is a martingale where X is an integrable random variable
- M is a supermartingale if $\mathbb{E}[M(t)|\mathcal{F}_s] \leq M(s)$

Martingale

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $\{\mathcal{F}_t\}_{t \geq 0}$ be a filtration on it. A stochastic process $M(t)$ is a martingale if

- $\mathbb{E}[|M(t)|] < \infty$
- $\mathbb{E}[M(t)|\mathcal{F}_s] = M(s)$ for all $s < t$.

Note that

- In some sense, a martingale represents a fair game
- $M(t) = \mathbb{E}[X|\mathcal{F}_t]$ is a martingale where X is an integrable random variable
- M is a supermartingale if $\mathbb{E}[M(t)|\mathcal{F}_s] \leq M(s)$
- M is a submartingale if $\mathbb{E}[M(t)|\mathcal{F}_s] \geq M(s)$

Markov Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $\{\mathcal{F}_t\}_{t \geq 0}$ be a filtration on it. A stochastic process $X(t)$ is a Markov process if for any Borel function f and $s < t$, there is a function g such that

$$\mathbb{E}[f(X(t)) | \mathcal{F}_s] = g(X(s))$$

Markov Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $\{\mathcal{F}_t\}_{t \geq 0}$ be a filtration on it. A stochastic process $X(t)$ is a Markov process if for any Borel function f and $s < t$, there is a function g such that

$$\mathbb{E}[f(X(t)) | \mathcal{F}_s] = g(X(s))$$

The estimate of $f(X(t))$ at time s depends only on the value of X at time s and not on the path of the process before time s .

Markov Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and $\{\mathcal{F}_t\}_{t \geq 0}$ be a filtration on it. A stochastic process $X(t)$ is a Markov process if for any Borel function f and $s < t$, there is a function g such that

$$\mathbb{E}[f(X(t)) | \mathcal{F}_s] = g(X(s))$$

The estimate of $f(X(t))$ at time s depends only on the value of X at time s and not on the path of the process before time s . The condition is equivalent to

$$\mathbb{E}[f(X(t)) | \mathcal{F}_s] = \mathbb{E}[f(X(t)) | X(s)]$$

Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

A stochastic process W is a Brownian motion if

1 $W(0) = 0$

Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

A stochastic process W is a Brownian motion if

- 1 $W(0) = 0$
- 2 For all $\omega \in \Omega$, $W(t, \omega)$ as a function of t is continuous.

Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

A stochastic process W is a Brownian motion if

- 1 $W(0) = 0$
- 2 For all $\omega \in \Omega$, $W(t, \omega)$ as a function of t is continuous.
- 3 For $0 = t_0 < t_1 < \dots < t_n$, the increments $W(t_1) - W(t_0), \dots, W(t_n) - W(t_{n-1})$ are independent.

Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

A stochastic process W is a Brownian motion if

- 1 $W(0) = 0$
- 2 For all $\omega \in \Omega$, $W(t, \omega)$ as a function of t is continuous.
- 3 For $0 = t_0 < t_1 < \dots < t_n$, the increments $W(t_1) - W(t_0), \dots, W(t_n) - W(t_{n-1})$ are independent.
- 4 For any $s < t$, $W(t) - W(s)$ are normally distributed with mean 0 and variance $t - s$.

Filtration for Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $\{\mathcal{F}_t\}_{t \geq 0}$ be a filtration. It is a filtration for the Brownian motion $W(t)$ if

- $W(t)$ is adapted to $\{\mathcal{F}_t\}$
- $\forall t > s$, $W(t) - W(s)$ is independent of \mathcal{F}_s

Properties of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Properties of Brownian Motion

- $W(t)$ is a martingale

Properties of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Properties of Brownian Motion

- $W(t)$ is a martingale
- $W(t)$ is a Markov process

Properties of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Proof of Markov Property.

For any Borel function f and $s < t$, we shall find a function g such that

$$\mathbb{E}[f(W(t))|\mathcal{F}_s] = g(W(s))$$

Properties of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Proof of Markov Property.

For any Borel function f and $s < t$, we shall find a function g such that

$$\mathbb{E}[f(W(t))|\mathcal{F}_s] = g(W(s))$$

Note that

$$\mathbb{E}[f(W(t))|\mathcal{F}_s] = \mathbb{E}[f(W(t) - W(s) + W(s))|\mathcal{F}_s]$$

Properties of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Proof of Markov Property.

For any Borel function f and $s < t$, we shall find a function g such that

$$\mathbb{E}[f(W(t))|\mathcal{F}_s] = g(W(s))$$

Note that

$$\mathbb{E}[f(W(t))|\mathcal{F}_s] = \mathbb{E}[f(W(t) - W(s) + W(s))|\mathcal{F}_s]$$

Since $W(t) - W(s)$ is independent of \mathcal{F}_s and $W(s)$ is \mathcal{F}_s -measurable, we can compute the conditional expectation on the RHS as

$$\mathbb{E}[f(W(t) - W(s) + W(s))|\mathcal{F}_s] = g(W(s))$$

where $g(x) = \mathbb{E}[f(W(t) - W(s) + x)]$. This proves that $W(t)$ is a Markov process.

Transition Density of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

In the above proof, we can compute the function g explicitly.

$$g(x) = \mathbb{E}[f(W(t) - W(s) + x)]$$

Transition Density of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

In the above proof, we can compute the function g explicitly.

$$\begin{aligned} g(x) &= \mathbb{E}[f(W(t) - W(s) + x)] \\ &= \int f(u + x) \frac{1}{\sqrt{2\pi(t-s)}} e^{-\frac{u^2}{2(t-s)}} du \end{aligned}$$

Transition Density of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

In the above proof, we can compute the function g explicitly.

$$\begin{aligned} g(x) &= \mathbb{E}[f(W(t) - W(s) + x)] \\ &= \int f(u + x) \frac{1}{\sqrt{2\pi(t-s)}} e^{-\frac{u^2}{2(t-s)}} du \\ &= \int f(y) \frac{1}{\sqrt{2\pi(t-s)}} e^{-\frac{(y-x)^2}{2(t-s)}} dy \end{aligned}$$

Transition Density of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Define $p(t-s, x, y) = \frac{1}{\sqrt{2\pi(t-s)}} e^{-\frac{(y-x)^2}{2(t-s)}}$, then

$$\mathbb{E}[f(W(t)) | \mathcal{F}_s] = \int f(y) p(t-s, W(s), y) dy$$

$p(t-s, x, y)$ is the transition density for $W(t)$ given $W(s) = x$.

Quadratic Variation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Fix $T > 0$, the quadratic variation of a function f is

$$\lim_{||\Pi|| \rightarrow 0} \sum_{i=1}^n (f(t_i) - f(t_{i-1}))^2$$

where $\Pi : 0 = t_0 < t_1 < \dots < t_n = T$ is a partition of the interval $[0, T]$ and $||\Pi|| = \max_i |t_i - t_{i-1}|$.

Quadratic Variation

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

If f is smooth enough (e.g., f has a continuous first-order derivative), $f(t_i) - f(t_{i-1})$ is small. And $(f(t_i) - f(t_{i-1}))^2$ is even smaller. Hence the summation in the definition will go to 0. However this is not the case for Brownian motion.

Quadratic Variation of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

For each $\omega \in \Omega$, $W(\cdot, \omega)$ can be viewed as a (continuous) function of t . Hence we can define quadratic variation for $W(\cdot, \omega)$ as

$$\lim_{\|\Pi\| \rightarrow 0} \sum_{i=1}^n (W(t_i, \omega) - W(t_{i-1}, \omega))^2$$

provided that the limit exists.

Quadratic Variation of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $X(t, \omega)$ be a stochastic process, if there exists a random variable Y such that for any $\delta > 0$

$$\lim_{\|\Pi\| \rightarrow 0} \mathbb{P} \left[\left| \sum_{i=1}^n (X(t_i) - X(t_{i-1}))^2 - Y \right| > \delta \right] = 0$$

Then we say Y is the quadratic variation of X on $[0, T]$, denoted by $[X, X](T)$.

Quadratic Variation of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Suppose t_i are equally spaced. Let $t_i - t_{i-1} = \frac{T}{n}$,

$$\sum_{i=1}^n (W(t_i, \omega) - W(t_{i-1}, \omega))^2$$

Quadratic Variation of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Suppose t_i are equally spaced. Let $t_i - t_{i-1} = \frac{T}{n}$,

$$\begin{aligned} & \sum_{i=1}^n (W(t_i, \omega) - W(t_{i-1}, \omega))^2 \\ &= \frac{T}{n} \sum_{i=1}^n \left(\frac{W(t_i, \omega) - W(t_{i-1}, \omega)}{\sqrt{T/n}} \right)^2 \end{aligned}$$

Quadratic Variation of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Suppose t_i are equally spaced. Let $t_i - t_{i-1} = \frac{T}{n}$,

$$\begin{aligned} & \sum_{i=1}^n (W(t_i, \omega) - W(t_{i-1}, \omega))^2 \\ &= \frac{T}{n} \sum_{i=1}^n \left(\frac{W(t_i, \omega) - W(t_{i-1}, \omega)}{\sqrt{T/n}} \right)^2 \\ &= T \left(\frac{1}{n} \sum_{i=1}^n \left(\frac{W(t_i, \omega) - W(t_{i-1}, \omega)}{\sqrt{T/n}} \right)^2 \right) \end{aligned}$$

Quadratic Variation of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Suppose t_i are equally spaced. Let $t_i - t_{i-1} = \frac{T}{n}$,

$$\begin{aligned} & \sum_{i=1}^n (W(t_i, \omega) - W(t_{i-1}, \omega))^2 \\ &= \frac{T}{n} \sum_{i=1}^n \left(\frac{W(t_i, \omega) - W(t_{i-1}, \omega)}{\sqrt{T/n}} \right)^2 \\ &= T \left(\frac{1}{n} \sum_{i=1}^n \left(\frac{W(t_i, \omega) - W(t_{i-1}, \omega)}{\sqrt{T/n}} \right)^2 \right) \end{aligned}$$

Note that $\left(\frac{W(t_i, \omega) - W(t_{i-1}, \omega)}{\sqrt{T/n}} \right)^2$ are i.i.d. with mean 1. From Strong Law of Law Number

$$\frac{1}{n} \left(\sum_{i=1}^n \left(\frac{W(t_i, \omega) - W(t_{i-1}, \omega)}{\sqrt{T/n}} \right)^2 \right) \rightarrow 1$$

Quadratic Variation of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Suppose t_i are equally spaced. Let $t_i - t_{i-1} = \frac{T}{n}$,

$$\begin{aligned} & \sum_{i=1}^n (W(t_i, \omega) - W(t_{i-1}, \omega))^2 \\ &= \frac{T}{n} \sum_{i=1}^n \left(\frac{W(t_i, \omega) - W(t_{i-1}, \omega)}{\sqrt{T/n}} \right)^2 \\ &= T \left(\frac{1}{n} \sum_{i=1}^n \left(\frac{W(t_i, \omega) - W(t_{i-1}, \omega)}{\sqrt{T/n}} \right)^2 \right) \end{aligned}$$

Note that $\left(\frac{W(t_i, \omega) - W(t_{i-1}, \omega)}{\sqrt{T/n}} \right)^2$ are i.i.d. with mean 1. From Strong Law of Law Number

$$\frac{1}{n} \left(\sum_{i=1}^n \left(\frac{W(t_i, \omega) - W(t_{i-1}, \omega)}{\sqrt{T/n}} \right)^2 \right) \rightarrow 1$$

Caveat: This is not a rigorous proof.

Quadratic Variation of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem

The quadratic variation of a Brownian motion W on $[0, T]$ is

$$[W, W](T) = T$$

Quadratic Variation of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem

The quadratic variation of a Brownian motion W on $[0, T]$ is

$$[W, W](T) = T$$

- The theorem also holds for almost sure convergence if the partition is carefully chosen.

Quadratic Variation of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem

The quadratic variation of a Brownian motion W on $[0, T]$ is

$$[W, W](T) = T$$

- The theorem also holds for almost sure convergence if the partition is carefully chosen.
- The quadratic variation of a process is usually a random variable. However for a Brownian motion it is a non-zero constant.

Quadratic Variation of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem

The quadratic variation of a Brownian motion W on $[0, T]$ is

$$[W, W](T) = T$$

- The theorem also holds for almost sure convergence if the partition is carefully chosen.
- The quadratic variation of a process is usually a random variable. However for a Brownian motion it is a non-zero constant.
- This will be the source of volatility in our models.

Quadratic Variation of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Brownian motion accumulates variation at the rate of 1 per unit time. Informally we write it as

$$dW(t)dW(t) = dt$$

Quadratic Variation of Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

It is easy to see that

$$\lim_{\|\Pi\| \rightarrow 0} \sum_{i=1}^n (W(t_i) - W(t_{i-1}))(t_i - t_{i-1}) = 0$$

and

$$\lim_{\|\Pi\| \rightarrow 0} \sum_{i=1}^n (t_i - t_{i-1})^2 = 0$$

This can be written as

$$dW(t)dt = 0$$

and

$$dtdt = 0$$

Realized Volatility

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Suppose $S(t)$ follows a geometric Brownian motion as in the BSM model, then

$$S(t) = S(0)e^{(r - \frac{\sigma^2}{2})t + \sigma W(t)}$$

We often use

$$\frac{1}{n} \sum_{i=1}^n \left(\ln \left(\frac{S(t_i)}{S(t_{i-1})} \right) \right)^2$$

to estimate σ^2 . Now we justify it.

Realized Volatility

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

$$\frac{1}{n} \sum_{i=1}^n \left[\ln \left(\frac{S(t_i)}{S(t_{i-1})} \right) \right]^2$$

Realized Volatility

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \left[\ln\left(\frac{S(t_i)}{S(t_{i-1})}\right) \right]^2 \\ = & \frac{1}{n} \sum_{i=1}^n \left[\left(r - \frac{1}{2}\sigma^2\right)\Delta t_i + \sigma\Delta W_{t_i} \right]^2 \end{aligned}$$

Realized Volatility

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \left[\ln\left(\frac{S(t_i)}{S(t_{i-1})}\right) \right]^2 \\ = & \frac{1}{n} \sum_{i=1}^n \left[\left(r - \frac{1}{2}\sigma^2\right)\Delta t_i + \sigma\Delta W_{t_i} \right]^2 \\ = & \frac{1}{n} \sum_{i=1}^n \left[\left(r - \frac{1}{2}\sigma^2\right)^2(\Delta t_i)^2 + \right. \\ & \left. \sigma^2(\Delta W_{t_i})^2 + \left(r - \frac{1}{2}\sigma^2\right)\sigma\Delta t_i\Delta W_{t_i} \right] \end{aligned}$$

Realized Volatility

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \left[\ln\left(\frac{S(t_i)}{S(t_{i-1})}\right) \right]^2 \\ = & \frac{1}{n} \sum_{i=1}^n \left[\left(r - \frac{1}{2}\sigma^2\right)\Delta t_i + \sigma\Delta W_{t_i} \right]^2 \\ = & \frac{1}{n} \sum_{i=1}^n \left[\left(r - \frac{1}{2}\sigma^2\right)^2(\Delta t_i)^2 + \right. \\ & \quad \left. \sigma^2(\Delta W_{t_i})^2 + \left(r - \frac{1}{2}\sigma^2\right)\sigma\Delta t_i\Delta W_{t_i} \right] \\ = & \frac{1}{n} \left[\left(r - \frac{1}{2}\sigma^2\right)^2 \sum_{i=1}^n (\Delta t_i)^2 + \right. \\ & \quad \left. \sigma^2 \sum_{i=1}^n (\Delta W_{t_i})^2 + \right. \\ & \quad \left. \left(r - \frac{1}{2}\sigma^2\right)\sigma \sum_{i=1}^n \Delta t_i\Delta W_{t_i} \right] \end{aligned}$$

Realized Volatility

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \left[\ln\left(\frac{S(t_i)}{S(t_{i-1})}\right) \right]^2 \\ = & \frac{1}{n} \sum_{i=1}^n \left[\left(r - \frac{1}{2}\sigma^2\right)\Delta t_i + \sigma\Delta W_{t_i} \right]^2 \\ = & \frac{1}{n} \sum_{i=1}^n \left[\left(r - \frac{1}{2}\sigma^2\right)^2(\Delta t_i)^2 + \right. \\ & \quad \left. \sigma^2(\Delta W_{t_i})^2 + \left(r - \frac{1}{2}\sigma^2\right)\sigma\Delta t_i\Delta W_{t_i} \right] \\ = & \frac{1}{n} \left[\left(r - \frac{1}{2}\sigma^2\right)^2 \sum_{i=1}^n (\Delta t_i)^2 + \right. \\ & \quad \left. \sigma^2 \sum_{i=1}^n (\Delta W_{t_i})^2 + \right. \\ & \quad \left. \left(r - \frac{1}{2}\sigma^2\right)\sigma \sum_{i=1}^n \Delta t_i\Delta W_{t_i} \right] \\ \rightarrow & \frac{T}{n}\sigma^2 \end{aligned}$$

which is daily variance if we take t_i to be daily.

Ito's Integral

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Suppose we hold $\Delta(t)$ shares of a stock and $W(t)$ is the stock price, the P&L of our portfolio between t_i to t_{i+1} is

$$\Delta(t)(W(t_{i+1}) - W(t_i)) = \Delta(t)dW(t_i)$$

where

$$dW(t_i) = W(t_{i+1}) - W(t_i)$$

Ito's Integral

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Suppose we hold $\Delta(t)$ shares of a stock and $W(t)$ is the stock price, the P&L of our portfolio between t_i to t_{i+1} is

$$\Delta(t)(W(t_{i+1}) - W(t_i)) = \Delta(t)dW(t_i)$$

where

$$dW(t_i) = W(t_{i+1}) - W(t_i)$$

The accumulated P&L over the time $[0, T]$ is

$$\sum_{i=1}^n \Delta(t_i)dW(t_i)$$

Ito's Integral

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

In the limit case as $dt_i (= t_{i+1} - t_i) \rightarrow 0$, it becomes

$$\int_0^T \Delta(t) dW(t)$$

provided the limit is well defined.

Ito's Integral

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

In calculus, we define the integral of a function f on $[a, b]$ as

$$\lim_{||\Pi|| \rightarrow 0} \sum_{i=1}^n f(x_i^*)(x_i - x_{i-1})$$

where x_i^* is an arbitrary point chosen from the interval $[x_{i-1}, x_i]$ and Π is a partition.

Ito's Integral

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

In calculus, we define the integral of a function f on $[a, b]$ as

$$\lim_{||\Pi|| \rightarrow 0} \sum_{i=1}^n f(x_i^*)(x_i - x_{i-1})$$

where x_i^* is an arbitrary point chosen from the interval $[x_{i-1}, x_i]$ and Π is a partition.

However, the difference between $W(t_i)$ and $W(t_{i+1})$ is much bigger and hence the choice of x_i^* does matter.

Ito's Integral

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

In calculus, we define the integral of a function f on $[a, b]$ as

$$\lim_{||\Pi|| \rightarrow 0} \sum_{i=1}^n f(x_i^*)(x_i - x_{i-1})$$

where x_i^* is an arbitrary point chosen from the interval $[x_{i-1}, x_i]$ and Π is a partition.

However, the difference between $W(t_i)$ and $W(t_{i+1})$ is much bigger and hence the choice of x_i^* does matter.

Ito's integral chooses $x_i^* = x_{i-1}$.

Ito's Integral

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Notations

- Probability space $(\Omega, \mathcal{F}, \mathbb{P})$
- $W(t, \omega)$ is a Brownian motion
- $\{\mathcal{F}_t\}_{t \geq 0}$ is a filtration for the Brownian motion W
- $\Delta(t, \omega)$ is a adapted stochastic process

Ito's Integral for Simple Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Fix a partition $\Pi : 0 = t_0 < \dots < t_n = T$.

Definition

The stochastic process $\Delta(t, \omega)$ is called a simple process if for any $\omega \in \Omega$, $\Delta(t, \omega)$ is constant in t on each interval $[t_{i-1}, t_i]$.

Ito's Integral for Simple Process


Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus



Ito's Integral for Simple Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $\Delta(t, \omega)$ be a simple process. The Ito's integral is defined as

$$\int_0^T \Delta(t, \omega) dW(t) = \sum_{i=1}^n \Delta(t_{i-1}, \omega) (W(t_i) - W(t_{i-1}))$$

Ito's Integral for General Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Let $\Delta(t, \omega)$ be an adapted process such that

$$\mathbb{E} \left[\int_0^T \Delta^2(t, \omega) dt < \infty \right]$$

Ito's Integral for General Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

We can define Ito's integral for Δ in the following steps:

Ito's Integral for General Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

We can define Ito's integral for Δ in the following steps:

- 1 Approximate Δ with a sequence of simple processes Δ_n such that

$$\mathbb{E} \left[\int_0^T (\Delta_n - \Delta)^2 dt \right] \rightarrow 0$$

Ito's Integral for General Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

We can define Ito's integral for Δ in the following steps:

- 1 Approximate Δ with a sequence of simple processes Δ_n such that

$$\mathbb{E} \left[\int_0^T (\Delta_n - \Delta)^2 dt \right] \rightarrow 0$$

- 2 Let $I_n(\omega) = \int_0^T \Delta_n dW(t)$, we can show that there exists a random variable X such that

$$\mathbb{E} [(I_n(\omega) - X(\omega))^2] \rightarrow 0$$

Ito's Integral for General Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

We can define Ito's integral for Δ in the following steps:

- 1 Approximate Δ with a sequence of simple processes Δ_n such that

$$\mathbb{E} \left[\int_0^T (\Delta_n - \Delta)^2 dt \right] \rightarrow 0$$

- 2 Let $I_n(\omega) = \int_0^T \Delta_n dW(t)$, we can show that there exists a random variable X such that

$$\mathbb{E} [(I_n(\omega) - X(\omega))^2] \rightarrow 0$$

- 3 Define

$$\int_0^T \Delta(t, \omega) dW(t) = X(\omega)$$

Properties of Ito's Integral

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Let $I(t, \omega) = \int_0^t \Delta(s, \omega) dW(s)$, we have

- *Continuous sample path*

$I(t, \omega)$ is continuous in t for each ω

Properties of Ito's Integral

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Let $I(t, \omega) = \int_0^t \Delta(s, \omega) dW(s)$, we have

- *Continuous sample path*
 $I(t, \omega)$ is continuous in t for each ω
- *Adaptivity*
 $I(t, \omega)$ is \mathcal{F}_t -measurable.

Properties of Ito's Integral

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Let $I(t, \omega) = \int_0^t \Delta(s, \omega) dW(s)$, we have

- *Continuous sample path*

$I(t, \omega)$ is continuous in t for each ω

- *Adaptivity*

$I(t, \omega)$ is \mathcal{F}_t -measurable.

- *Linearity*

Let Δ, Γ be two processes and α, β be constant, then

$$\begin{aligned} & \int_0^T (\alpha \Delta(t) + \beta \Gamma(t)) dW(t) \\ = & \alpha \int_0^T \Delta(t) dW(t) + \beta \int_0^T \Gamma(t) dW(t) \end{aligned}$$

Properties of Ito's Integral

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

■ *Martingale*

$I(t, \omega)$ is a martingale. In particular, $\mathbb{E}[I(t)] = I(0) = 0$

Properties of Ito's Integral

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

- *Martingale*

$I(t, \omega)$ is a martingale. In particular, $\mathbb{E}[I(t)] = I(0) = 0$

- *Ito Isometry*

$$\mathbb{E}[I^2(t, \omega)] = \mathbb{E} \left[\int_0^t \Delta^2(s, \omega) ds \right] = \int_0^t \mathbb{E}[\Delta^2(s, \omega)] ds$$

Properties of Ito's Integral

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

- *Martingale*

$I(t, \omega)$ is a martingale. In particular, $\mathbb{E}[I(t)] = I(0) = 0$

- *Ito Isometry*

$$\mathbb{E}[I^2(t, \omega)] = \mathbb{E} \left[\int_0^t \Delta^2(s, \omega) ds \right] = \int_0^t \mathbb{E}[\Delta^2(s, \omega)] ds$$

- *Quadratic Variation*

$$[I, I](t, \omega) = \int_0^t \Delta^2(s, \omega) ds$$

Differential Forms

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

- From $I(t, \omega) = \int_0^t \Delta(s, \omega) dW(s)$ we have

$$dI(t) = \Delta dW(t)$$

Differential Forms

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

- From $I(t, \omega) = \int_0^t \Delta(s, \omega) dW(s)$ we have

$$dI(t) = \Delta dW(t)$$

- From $[I, I](t, \omega) = \int_0^t \Delta^2(s, \omega) ds$ we have

$$dI(t)dI(t) = \Delta dW(t)\Delta dW(t) = \Delta^2(t)dt$$

Ito's Lemma

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Lemma

Let $f(t, x)$ be a function for which the partial derivatives f_t, f_x and f_{xx} are continuous, and let $W(t)$ be a Brownian motion. Then

$$f(T, W(T)) = f(0, W(0)) + \int_0^T f_t(s, W(s))ds + \int_0^T f_x(s, W(s))dW(s) + \frac{1}{2} \int_0^T f_{xx}(s, W(s))ds$$

Ito's Lemma

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

An intuitive proof.

Since

$$df(t, x) = f_t dt + f_x dx + \frac{1}{2} f_{xx} dx^2 + \frac{1}{2} f_{tt} dt^2 + f_{tx} dt dx + \dots$$

we have

$$\begin{aligned} df(t, W(t)) &= f_t(t, W(t))dt + f_x(t, W(t))dW(t) \\ &+ \frac{1}{2} f_{xx}(t, W(t))dW(t)dW(t) \\ &+ \frac{1}{2} f_{tt}(t, W(t))dt^2 + f_{tx}(t, W(t))dt dW(t) \\ &+ \dots \end{aligned}$$



Ito's Lemma

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

An intuitive proof.

Note that $dW(t)dt = 0$, $dt dt = 0$ and $dW(t)dW(t) = dt$, the above equation reduces to

$$\begin{aligned} df(t, W(t)) &= f_t(t, W(t))dt + f_x(t, W(t))dW(t) \\ &\quad + \frac{1}{2}f_{xx}(t, W(t))dt \end{aligned}$$

Integrating it we get Ito's lemma. □

Ito's Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let $W(t)$ be a Brownian motion, $\{\mathcal{F}_t\}_{t \geq 0}$ be a filtration for it, $\Delta(t)$ and $\Theta(t)$ be two adapted processes. Define

$$X(t) = X(0) + \int_0^t \Delta(s) dW(s) + \int_0^t \Theta(s) ds$$

where $X(0)$ is non-random. Then X is called an Ito process.

Quadratic Variation of Ito Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Lemma

$$[X, X](t) = \int_0^t \Delta^2(s) ds$$

Proof.

HW



Ito's Integral for Ito Process

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

Let

$$X(t) = X(0) + \int_0^t \Delta(s) dW(s) + \int_0^t \Theta(s) ds$$

be an Ito process, $\Gamma(t)$ be an adapted process. We define the integral with respect to $X(t)$ as

$$\int_0^t \Gamma(s) dX(s) = \int_0^t \Gamma(s) \Delta(s) dW(s) + \int_0^t \Gamma(s) \Theta(s) ds$$

Ito's Lemma

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Lemma

Let

$$X(t) = X(0) + \int_0^t \Delta(s) dW(s) + \int_0^t \Theta(s) ds$$

be an Ito process, $f(t, x)$ be a function whose partial derivatives f_t, f_x and f_{xx} are continuous. Then

$$\begin{aligned} & f(T, X(T)) \\ = & f(0, X(0)) + \int_0^T f_t(s, X(s)) ds + \int_0^T f_x(s, X(s)) dX(s) \\ & + \frac{1}{2} \int_0^T f_{xx}(s, X(s)) d[X, X](s) \end{aligned}$$

Ito's Lemma

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Lemma

Let

$$X(t) = X(0) + \int_0^t \Delta(s) dW(s) + \int_0^t \Theta(s) ds$$

be an Ito process, $f(t, x)$ be a function whose partial derivatives f_t, f_x and f_{xx} are continuous. Then

$$\begin{aligned} & f(T, X(T)) \\ = & f(0, X(0)) + \int_0^T f_t(s, X(s)) ds + \int_0^T f_x(s, X(s)) dX(s) \\ & + \frac{1}{2} \int_0^T f_{xx}(s, X(s)) d[X, X](s) \end{aligned}$$

Ito's Lemma

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Lemma

Let

$$X(t) = X(0) + \int_0^t \Delta(s) dW(s) + \int_0^t \Theta(s) ds$$

be an Ito process, $f(t, x)$ be a function whose partial derivatives f_t, f_x and f_{xx} are continuous. Then

$$\begin{aligned} & f(T, X(T)) \\ = & f(0, X(0)) + \int_0^T f_t(s, X(s)) ds + \int_0^T f_x(s, X(s)) dX(s) \\ & + \frac{1}{2} \int_0^T f_{xx}(s, X(s)) d[X, X](s) \\ = & f(0, X(0)) + \int_0^T f_t(s, X(s)) ds + \int_0^T f_x(s, X(s)) \Delta(s) dW(s) \\ & + \int_0^T f_x(s, X(s)) \Theta(s) ds + \frac{1}{2} \int_0^T f_{xx}(s, X(s)) \Delta^2(s) ds \end{aligned}$$

Ito's Lemma

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

We can write Ito's Lemma in differential form as

$$df(t, X) = f_t(t, X)dt + f_x(t, X)dX(t) + \frac{1}{2}f_{xx}(t, X)dX(t)dX(t)$$

Ito's Lemma

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

We can write Ito's Lemma in differential form as

$$df(t, X) = f_t(t, X)dt + f_x(t, X)dX(t) + \frac{1}{2}f_{xx}(t, X)dX(t)dX(t)$$

Expanding it we have

$$\begin{aligned} df(t, X) = & f_t(t, X)dt + f_x(t, X)\Delta(t)dW(t) \\ & + f_x(t, X)\Theta(t)dt + \frac{1}{2}f_{xx}(t, X)\Delta^2(t)dt \end{aligned}$$

Ito's Integral of a deterministic integrand

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem

*Let $\Delta(t)$ be a non-random function of time t ,
 $I(t) = \int_0^t \Delta(s) dW(s)$. Then $I(t)$ is normally distributed with
mean zero and variance $\int_0^t \Delta^2(s) ds$.*

Ito's Integral of a deterministic integrand

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem

*Let $\Delta(t)$ be a non-random function of time t ,
 $I(t) = \int_0^t \Delta(s) dW(s)$. Then $I(t)$ is normally distributed with
mean zero and variance $\int_0^t \Delta^2(s) ds$.*

Proof.

Since $I(t)$ is a martingale and $I(0) = 0$, $\mathbb{E}[I(t)] = I(0) = 0$.

Ito's Integral of a deterministic integrand

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem

Let $\Delta(t)$ be a non-random function of time t ,
 $I(t) = \int_0^t \Delta(s) dW(s)$. Then $I(t)$ is normally distributed with
mean zero and variance $\int_0^t \Delta^2(s) ds$.

Proof.

Since $I(t)$ is a martingale and $I(0) = 0$, $\mathbb{E}[I(t)] = I(0) = 0$.
By Ito Isometry, $\mathbb{E}[I^2(t)] = \int_0^t \Delta^2(s) ds$. Hence the variance of
 $I(t)$ is $\int_0^t \Delta^2(s) ds$.



Ito's Integral of a deterministic integrand

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Proof.

To show that $I(t)$ is normally distributed, we consider its moment generating function $m(s) = \mathbb{E}[e^{sI(t)}]$.

Ito's Integral of a deterministic integrand

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Proof.

To show that $I(t)$ is normally distributed, we consider its moment generating function $m(s) = \mathbb{E}[e^{sI(t)}]$.

We need to show that

$$m(s) = \mathbb{E}[e^{sI(t)}] = e^{\frac{1}{2}s^2 \int_0^t \Delta^2(u) du}$$

Ito's Integral of a deterministic integrand

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Proof.

To show that $I(t)$ is normally distributed, we consider its moment generating function $m(s) = \mathbb{E}[e^{sI(t)}]$.

We need to show that

$$m(s) = \mathbb{E}[e^{sI(t)}] = e^{\frac{1}{2}s^2 \int_0^t \Delta^2(u) du}$$

which is equivalent to showing

$$\mathbb{E}[e^{sI(t) - \frac{1}{2}s^2 \int_0^t \Delta^2(u) du}] = 1$$



Ito's Integral of a deterministic integrand

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Proof.

Fix s , define $Y(t) = sI(t) - \frac{1}{2}s^2 \int_0^t \Delta^2(u)du$. Since $Y(0) = 0$, if we can prove that $e^{Y(t)}$ is a martingale, then $\mathbb{E}[e^{Y(t)}] = 1$, we are done.

Ito's Integral of a deterministic integrand

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Proof.

Fix s , define $Y(t) = sl(t) - \frac{1}{2}s^2 \int_0^t \Delta^2(u)du$. Since $Y(0) = 0$, if we can prove that $e^{Y(t)}$ is a martingale, then $\mathbb{E}[e^{Y(t)}] = 1$, we are done.

Since

$$\begin{aligned} dY(t) &= s dl(t) - \frac{1}{2}s^2 \Delta^2(t)dt \\ &= s \Delta(t) dW(t) - \frac{1}{2}s^2 \Delta^2(t)dt \end{aligned}$$

Ito's Integral of a deterministic integrand

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Proof.

Fix s , define $Y(t) = sl(t) - \frac{1}{2}s^2 \int_0^t \Delta^2(u)du$. Since $Y(0) = 0$, if we can prove that $e^{Y(t)}$ is a martingale, then $\mathbb{E}[e^{Y(t)}] = 1$, we are done.

Since

$$\begin{aligned} dY(t) &= s dl(t) - \frac{1}{2}s^2 \Delta^2(t)dt \\ &= s \Delta(t) dW(t) - \frac{1}{2}s^2 \Delta^2(t)dt \end{aligned}$$

and

$$dY(t)dY(t) = s^2 \Delta^2(t)dt$$

Ito's Integral of a deterministic integrand

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Proof.

Fix s , define $Y(t) = sl(t) - \frac{1}{2}s^2 \int_0^t \Delta^2(u)du$. Since $Y(0) = 0$, if we can prove that $e^{Y(t)}$ is a martingale, then $\mathbb{E}[e^{Y(t)}] = 1$, we are done.

Since

$$\begin{aligned} dY(t) &= s dl(t) - \frac{1}{2}s^2 \Delta^2(t)dt \\ &= s \Delta(t) dW(t) - \frac{1}{2}s^2 \Delta^2(t)dt \end{aligned}$$

and

$$dY(t)dY(t) = s^2 \Delta^2(t)dt$$

we have

$$dY(t) + \frac{1}{2}dY(t)dY(t) = s\Delta(t)dW(t)$$



Ito's Integral of a deterministic integrand

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Proof.

Applying Ito's Lemma to $e^{Y(t)}$, we have

$$\begin{aligned} de^{Y(t)} &= e^{Y(t)}dY(t) + \frac{1}{2}e^{Y(t)}dY(t)dY(t) \\ &= e^{Y(t)}\left(dY(t) + \frac{1}{2}dY(t)dY(t)\right) \end{aligned}$$

Ito's Integral of a deterministic integrand

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Proof.

Applying Ito's Lemma to $e^{Y(t)}$, we have

$$\begin{aligned} de^{Y(t)} &= e^{Y(t)}dY(t) + \frac{1}{2}e^{Y(t)}dY(t)dY(t) \\ &= e^{Y(t)}\left(dY(t) + \frac{1}{2}dY(t)dY(t)\right) \end{aligned}$$

Substituting the equation from last slide, we have

$$de^{Y(t)} = s\Delta(t)e^{Y(t)}dW(t)$$

Ito's Integral of a deterministic integrand

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Proof.

Applying Ito's Lemma to $e^{Y(t)}$, we have

$$\begin{aligned} de^{Y(t)} &= e^{Y(t)}dY(t) + \frac{1}{2}e^{Y(t)}dY(t)dY(t) \\ &= e^{Y(t)}\left(dY(t) + \frac{1}{2}dY(t)dY(t)\right) \end{aligned}$$

Substituting the equation from last slide, we have

$$de^{Y(t)} = s\Delta(t)e^{Y(t)}dW(t)$$

Hence

$$e^{Y(t)} = 1 + s \int_0^t \Delta(s)e^{Y(s)}dW(s)$$

is a martingale. □

Multiple Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Definition

A d -dimensional Brownian motion is a process

$$W(t) = (W_1(t), \dots, W_d(t))$$

with the following properties

- Each $W_i(t)$ is a one-dimensional Brownian motion
- For $i \neq j$, the processes $W_i(t)$ and $W_j(t)$ are independent

Multiple Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space. A filtration $\{\mathcal{F}_t\}$ for the d -dimensional Brownian motion $W(t)$ is a collection of sub σ -algebra of \mathcal{F} such that

Multiple Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space. A filtration $\{\mathcal{F}_t\}$ for the d -dimensional Brownian motion $W(t)$ is a collection of sub σ -algebra of \mathcal{F} such that

$$\blacksquare \mathcal{F}_s \subset \mathcal{F}_t, \forall s \leq t$$

Multiple Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space. A filtration $\{\mathcal{F}_t\}$ for the d -dimensional Brownian motion $W(t)$ is a collection of sub σ -algebra of \mathcal{F} such that

- $\mathcal{F}_s \subset \mathcal{F}_t, \forall s \leq t$
- For each t , the random vector $W(t)$ is \mathcal{F}_t -measurable

Multiple Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space. A filtration $\{\mathcal{F}_t\}$ for the d -dimensional Brownian motion $W(t)$ is a collection of sub σ -algebra of \mathcal{F} such that

- $\mathcal{F}_s \subset \mathcal{F}_t, \forall s \leq t$
- For each t , the random vector $W(t)$ is \mathcal{F}_t -measurable
- For $s < t$, the increment $W(t) - W(s)$ is independent of \mathcal{F}_s .

Multiple Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Lemma

For each i ,

$$[W_i, W_i](T) = T$$

and $i \neq j$

$$[W_i, W_j](T) = 0$$



Multiple Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Lemma

For each i ,

$$[W_i, W_i](T) = T$$

and $i \neq j$

$$[W_i, W_j](T) = 0$$

This can be informally write as

$$dW_i(t)dW_i(t) = dt$$

and

$$dW_i(t)dW_j(t) = 0, \forall i \neq j$$

Multiple Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Lemma

For each i ,

$$[W_i, W_i](T) = T$$

and $i \neq j$

$$[W_i, W_j](T) = 0$$

This can be informally write as

$$dW_i(t)dW_i(t) = dt$$

and

$$dW_i(t)dW_j(t) = 0, \forall i \neq j$$

Proof.

HW



Two-dimensional Ito's Lemma

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem

Let $f(t, x, y)$ be a function whose partial derivatives $f_t, f_x, f_y, f_{xx}, f_{xy}, f_{yy}$ are continuous. Let $X(t)$ and $Y(t)$ be two Ito processes. Then we have

$$df(t, X, Y) = f_t dt + f_x dX(t) + f_y dY(t) + \frac{1}{2} f_{xx} dX(t) dX(t) + \frac{1}{2} f_{yy} dY(t) dY(t) + f_{xy} dX(t) dY(t)$$

Ito Product Rule

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Corollary

Let $X(t)$ and $Y(t)$ be two Ito processes, then

$$d(X(t)Y(t)) = Y(t)dX(t) + X(t)dY(t) + dX(t)dY(t)$$



Characterizing a Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

$W(t)$ Brownian motion \implies

- martingale
- continuous sample path
- $[W, W](t) = t$

Characterizing a Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

$W(t)$ Brownian motion \implies

- martingale
- continuous sample path
- $[W, W](t) = t$

It turns out the converse is also true.

Characterizing a Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem (Levy, one dimension)

Let $\{\mathcal{F}_t\}_{t \geq 0}$ be a filtration on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and $M(t)$ be a stochastic process with $M(0) = 0$. Suppose

- *$M(t)$ is a martingale*
- *$M(t)$ has a continuous sample path*
- *$[M, M](t) = t$*

Then $M(t)$ is a Brownian motion.

Characterizing a Brownian Motion

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Theorem (Levy, two dimensions)

Let $\{\mathcal{F}_t\}_{t \geq 0}$ be a filtration on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and $M_i(t), i = 1, 2$ be a stochastic process with $M_i(0) = 0$.

Suppose

- $M_i(t), i = 1, 2$ is a martingale
- $M_i(t), i = 1, 2$ has a continuous sample path
- $[M_i, M_i](t) = t$
- $[M_1, M_2](t) = 0$

Then $M_1(t)$ and $M_2(t)$ are independent Brownian motions.

Correlated Stock Prices

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Suppose two stock prices follow the following dynamics

$$\begin{aligned}\frac{dS_1(t)}{S_1(t)} &= \alpha_1 dt + \sigma_1 dW_1(t) \\ \frac{dS_2(t)}{S_2(t)} &= \alpha_2 dt + \sigma_2 \left(\rho dW_1(t) + \sqrt{1 - \rho^2} dW_2(t) \right)\end{aligned}$$

where $\alpha_1, \alpha_2, \sigma_1, \sigma_2, \rho$ are constant. We investigate how these two stock prices are correlated.

Correlated Stock Prices

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Let $W_3(t) = \rho W_1(t) + \sqrt{1 - \rho^2} W_2(t)$.

Correlated Stock Prices

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Let $W_3(t) = \rho W_1(t) + \sqrt{1 - \rho^2} W_2(t)$. Then

$$dW_3(t) = \rho dW_1(t) + \sqrt{1 - \rho^2} dW_2(t)$$

Correlated Stock Prices

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Let $W_3(t) = \rho W_1(t) + \sqrt{1 - \rho^2} W_2(t)$. Then

$$dW_3(t) = \rho dW_1(t) + \sqrt{1 - \rho^2} dW_2(t)$$

Hence

$$\begin{aligned} dW_3(t)dW_3(t) &= \rho^2 dW_1(t)dW_1(t) + (1 - \rho^2)dW_2(t)dW_2(t) \\ &\quad + 2\rho\sqrt{1 - \rho^2}dW_1(t)dW_2(t) \\ &= \rho^2 dt + (1 - \rho^2)dt \end{aligned}$$

Correlated Stock Prices

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Let $W_3(t) = \rho W_1(t) + \sqrt{1 - \rho^2} W_2(t)$. Then

$$dW_3(t) = \rho dW_1(t) + \sqrt{1 - \rho^2} dW_2(t)$$

Hence

$$\begin{aligned} dW_3(t)dW_3(t) &= \rho^2 dW_1(t)dW_1(t) + (1 - \rho^2)dW_2(t)dW_2(t) \\ &\quad + 2\rho\sqrt{1 - \rho^2}dW_1(t)dW_2(t) \\ &= \rho^2 dt + (1 - \rho^2)dt \\ &= dt \end{aligned}$$

Correlated Stock Prices

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

Let $W_3(t) = \rho W_1(t) + \sqrt{1 - \rho^2} W_2(t)$. Then

$$dW_3(t) = \rho dW_1(t) + \sqrt{1 - \rho^2} dW_2(t)$$

Hence

$$\begin{aligned} dW_3(t)dW_3(t) &= \rho^2 dW_1(t)dW_1(t) + (1 - \rho^2)dW_2(t)dW_2(t) \\ &\quad + 2\rho\sqrt{1 - \rho^2}dW_1(t)dW_2(t) \\ &= \rho^2 dt + (1 - \rho^2)dt \\ &= dt \end{aligned}$$

It is easy to verify that $W_3(t)$ is a martingale with continuous sample path. By Levy's Theorem, $W_3(t)$ is a Brownian motion.

Correlated Stock Prices

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

By Ito Product Rule,

$$\begin{aligned}d(W_1(t)W_3(t)) &= W_3dW_1(t) + W_1dW_3(t) + dW_1(t)dW_3(t) \\ &= W_3dW_1(t) + W_1dW_3(t) + \rho dt\end{aligned}$$

Correlated Stock Prices

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

By Ito Product Rule,

$$\begin{aligned}d(W_1(t)W_3(t)) &= W_3dW_1(t) + W_1dW_3(t) + dW_1(t)dW_3(t) \\ &= W_3dW_1(t) + W_1dW_3(t) + \rho dt\end{aligned}$$

Solving it we have

$$W_1(t)W_3(t) = \int_0^t W_3dW_1(t) + \int_0^t W_1dW_3(t) + \rho t$$

Correlated Stock Prices

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

By Ito Product Rule,

$$\begin{aligned}d(W_1(t)W_3(t)) &= W_3dW_1(t) + W_1dW_3(t) + dW_1(t)dW_3(t) \\ &= W_3dW_1(t) + W_1dW_3(t) + \rho dt\end{aligned}$$

Solving it we have

$$W_1(t)W_3(t) = \int_0^t W_3dW_1(t) + \int_0^t W_1dW_3(t) + \rho t$$

Hence

$$\mathbb{E}[W_1(t)W_3(t)] = \rho t$$

Correlated Stock Prices

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

By Ito Product Rule,

$$\begin{aligned}d(W_1(t)W_3(t)) &= W_3dW_1(t) + W_1dW_3(t) + dW_1(t)dW_3(t) \\ &= W_3dW_1(t) + W_1dW_3(t) + \rho dt\end{aligned}$$

Solving it we have

$$W_1(t)W_3(t) = \int_0^t W_3dW_1(t) + \int_0^t W_1dW_3(t) + \rho t$$

Hence

$$\mathbb{E}[W_1(t)W_3(t)] = \rho t$$

Since $W_1(t)$ and $W_3(t)$ have zero mean and variance t , the correlation between $W_1(t)$ and $W_3(t)$ is ρ .

Correlated Stock Prices

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

$S_2(t)$ can be written as

$$\frac{dS_2(t)}{S_2(t)} = \alpha_2 dt + \sigma_2 dW_3(t)$$

Correlated Stock Prices

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

$S_2(t)$ can be written as

$$\frac{dS_2(t)}{S_2(t)} = \alpha_2 dt + \sigma_2 dW_3(t)$$

Note that



$$dW_1(t)dW_3(t) = \rho dt$$

Correlated Stock Prices

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

$S_2(t)$ can be written as

$$\frac{dS_2(t)}{S_2(t)} = \alpha_2 dt + \sigma_2 dW_3(t)$$

Note that

■

$$dW_1(t)dW_3(t) = \rho dt$$

■

$$\frac{dS_1(t)}{S_1(t)} \frac{dS_2(t)}{S_2(t)} = \sigma_1 \sigma_2 \rho dt$$

.

Correlated Stock Prices

Mathematical
Foundation

Wu Lei

Probability

Stochastic
Process

Stochastic
Calculus

$S_2(t)$ can be written as

$$\frac{dS_2(t)}{S_2(t)} = \alpha_2 dt + \sigma_2 dW_3(t)$$

Note that

■

$$dW_1(t)dW_3(t) = \rho dt$$

■

$$\frac{dS_1(t)}{S_1(t)} \frac{dS_2(t)}{S_2(t)} = \sigma_1 \sigma_2 \rho dt$$

.

- ρ is also called instantaneous correlation (in particular when ρ is time-dependent).

Thank you!