

Storm Surges

March 31, 2014

Abstract

Abstract to be completed

1 Introduction

(Masson and Cummins, 2004)

2 Model Configuration

2.1 Model domain

The modelled domain extends from the Strait of Juan de Fuca to Puget Sound to Johnstone Strait as shown in Figure 1. Bathymetry from the Cascadia physiography dataset (Haugerud, 1999) was smoothed to limit the difference in depth across grid cells. For model stability, additional smoothing at the Strait of Juan de Fuca western boundary was imposed to achieve constant depth across the first ten grid cells.

2.2 Boundary conditions

2.2.1 Tidal forcing

The model was forced by tidal elevations and currents at the Juan de Fuca and Johnstone Strait boundaries. Tidal heights and currents for eight tidal constituents (K1, O1, P1, Q1, M2, K2, N2, S2) at grid points along the Juan de Fuca boundary were extracted from Webtide, an online web prediction model for the northeast Pacific Ocean, which is based on work by Foreman et al. (2000). The Johnstone Strait boundary was forced with current and elevation tidal harmonics measured and calculated by Thomson and Huggett (1980) for the major M2 and K1 constituents. Additionally, O1 and S2 elevation harmonics from their measurements were employed. The remaining constituents were extrapolated from Webtide.




Figure 1: Model domain including bathymetry, rivers (*), and storm surge locations (o) of interest.

2.2.2 Temperature and salinity

Temperature and salinity at the Juan de Fuca boundary were taken from a weekly climatology which was created from results from a model covering the Salish Sea and the west coast of Vancouver Island (Masson and Fine, 2012). Their results, originally on s-levels were interpolated onto z-levels and then onto the NEMO horizontal grid. To prepare the climatology all years (1995-2008) were averaged and results, approximately every 15 days, were interpolated to a weekly climatology.

2.2.3 Sea Surface Height

Sea surface height at the mouth of Juan de Fuca was set using values from the Tofino tide gauge. A monthly climatology was produced using daily averages from 2000-2010, binning them by month, averaging and setting the yearly mean to zero. For the storm surge simulations, hourly variations in sea surface height were used. These values are the Tofino tide gauge values, de-tided and with the zero reset as for the climatology.

2.2.4 Open Boundary Conditions

The model relaxed to the forced temperature and salinity over the 10 grid points (about 5km) closest to the open boundaries, using the NEMO FRS scheme (Madec, 2008). The tidal forcing and sea surface height was used in the barotropic velocity forcing which used

the NEMO Flather scheme Madec (2008). The baroclinic velocities at the boundary were set to be equal to the values inside the boundary (zero-gradient boundary conditions). This scheme is not part of core NEMO. Zero gradient conditions were chosen because the baroclinic velocity at the mouth of Juan de Fuca is primarily estuarine and thus set by density variations between inside and outside the domain.

2.3 River forcing

River input provides a significant volume of freshwater to the Salish Sea and can influence stratification, circulation and primary productivity. However, most rivers in the domain are not gauged so parameterisations were required to represent river flow. Morrison et al. (2011) provides a method for estimating freshwater runoff in the Salish Sea region based on precipitation. Monthly runoff volumes for each watershed for each year from 1970 to 2012 were acquired from Morrison et al. (2011), as well as monthly averages.

Freshwater runoff from each watershed was divided between the rivers in that watershed. The area drained by each river was estimated from Toporama maps by the Atlas of Canada and watershed maps available on the Washington State government website. The watersheds included in our model were Fraser (which represents approximately 44% of the freshwater input into our domain), Skagit (12%), East Vancouver Island (North and South) (12%), Howe (7%), Bute (7%), Puget (6%), Juan de Fuca (5%), Jervis (4%) and Toba (3%).

The monthly flow from each river was input as a point source in the three grid points closest to the surface at the model point closest to the mouth of each river. Incoming water was assumed to be fresh and at the temperature of the surface. A total of 150 rivers were parameterised by this method.

2.4 Atmospheric forcing

2.5 Initial conditions

Initial conditions for temperature and salinity were taken from a CTD cast in the middle Strait of Georgia taken in Sept 2002 (Pawlowicz et al., 2007). Conditions were initially uniform horizontally. Velocity was initialized at zero.

2.6 Spin-up

The model was spun up for a 15.5 months from the initial conditions above, starting Sep 16, 2002, using atmospheric forcing from 2002-2003, climatological temperature and salinity and sea surface height at the boundaries, with tides and climatological river output. All storm surge runs were started three days prior to the event of interest with zero initial velocities and sea surface height and a stratification profile from model spin up. The modelled sea surface height adjusted to forcing in less than one day.

3 Model Evaluation

3.1 Tidal evaluation

The model was initially evaluated qualitatively by comparing patterns of tidal amplitude and phase to results from (Foreman et al., 1995). For example, the amphidromic dome around Victoria was produced in the M_2 results, as well as the monotonic decrease in M_2 amplitude moving northwards along the Strait of Georgia. Initially, the Johnstone Strait boundary was closed, but modelled M_2 amplitudes were too small compared to measured amplitudes... TBC

Once our model was reproducing observed tidal patterns, model results were quantitatively evaluated by comparing modelled harmonic constituents to measured harmonic constituents at tidal measuring stations throughout the domain. Comparisons were made using the complex difference (D), defined by (Foreman et al., 1995) as:

$$D = [(A_0 \cos g_0 - A_m \cos g_m)^2 + (A_0 \sin g_0 - A_m \sin g_m)^2]^{1/2} \quad (1)$$

where A_0 , A_m , g_0 and g_m are observed and modelled amplitudes and phases.

Complex differences were less than ??cm at all stations in our domain, which was assumed to be acceptable for our purposes.

4 Storm Surge Hindcasts

5 Conclusions

References

- Foreman, M., W. Crawford, J. Cherniawsky, R. Henry, and M. Tarbotto (2000). A high-resolution assimilating tidal model for the northeast Pacific Ocean. *Journal of Geophysical Research* 105(C1), 28,629–28,652.
- Foreman, M. G. G., R. A. Walters, R. F. Henry, C. P. Keller, and A. G. Dolling (1995). A tidal model for eastern Juan de Fuca Strait and the southern Strait OF Georgia. *Journal of Geophysical Research-Oceans* 100(C1), 721–740.
- Haugerud, R. A. (1999). *Digital elevation model (DEM) of Cascadia, latitude 39N-53N, longitude 116W-133W*. U. S. Geological Survey Open-File Report 99-369. <http://pubs.usgs.gov/of/1999/0369/>.
- Madec, G. (2008). Nemo ocean engine.
- Masson, D. and P. F. Cummins (2004). Observations and modeling of seasonal variability in the Straits of Georgia and Juan de Fuca. *Journal of Marine Research* 62(4), 491–516.

- Masson, D. and I. Fine (2012). Modeling seasonal to interannual ocean variability of coastal british columbia. *Journal of Geophysical Research* 117(C10).
- Morrison, J., M. Foreman, and D. Masson (2011). A method for estimating monthly fresh-water discharge affecting British Columbia coastal waters. *Atmosphere-Ocean* 50(1), 1–8.
- Pawlowicz, R., O. Riche, and M. Halverson (2007). The circulation and residence time of the Strait of Georgia using a simple mixing-box approach. *Atmos. Ocean* 45, 173–193.
- Thomson, R. E. and W. S. Huggett (1980). M2 baroclinic tides in Johnstone Strait, British Columbia. *Journal of Physical Oceanography* 10, 1509–1539.