


Spatial dimension compactification in general relativity and duality symmetry

Adam Franchot-Fowler

Internship carried out from 08/06/2026 to 24/07/2026



Internship tutors : M. Camille Eloy

Referent teacher : M. Stéphane Douin

Éducational institution : Université Paris-Saclay / Magistère de physique fondamentale d'Orsay

Internship host company : Laboratoire de Physique (UMR CNRS 5672) ENS de Lyon 46, allée d'Italie F-69364 LYON CEDEX 07, FRANCE

Acknowledgment

Notations and abbreviations

For the abbreviations we will use

- GR for general relativity

We will during this internship use the vielbein formalism. We will use the same notation as in [1] which is

- Greek hatted letter for curved indices ($\hat{\mu}, \hat{\nu}, \hat{\rho}, \dots$)
- Latin hatted letter for flat indices ($\hat{p}, \hat{m}, \hat{n}, \dots$)
- For space-time indices taking D values, late greek letters are used in the curved case ($\mu, \nu, \rho, \dots$) and Latin letters in the flat case (p, m, n)
- For internal indices taking E values, early Greek letters in the curved case ($\alpha, \beta, \gamma, \dots$) and Latin letters in the flat case (a,b,c)

The general coordinate $x^{\hat{\mu}}$ will therefore be made of space time coordinate x^{μ} and internal coordinate y^{α} . The flat lorentz

Contents

Notation and abbreviation	1
1 Introduction	3
2 General relativity theoretical reminder	3
2.1 mathematical basis	3
2.2 spin connexion, vielbein and flatspace	3
3 Lagrangian field theory	3
3.1 Lagrangian density	3
3.2 General exemple for scalar and vectorial fields, and result for two forms	4
4 Reduction of hilbert-einstein Action in D+E dimension	4
4.1 Hypothesis	4
4.2 Rewriting the action	4
4.3 vielbein ansatz	4
4.3.1 Gauge transformations of the fields	5
4.3.2 Metrics	5
4.4 Latin connexion computation	5
4.5 finalisation	8
5 reduction of the new fields	8
5.1 scalar field reduction	9
5.2 Two form fiel reduction	9
6 Observation of new symmetry	9
7 Conclusion	9
A lagrangian	9

1 Introduction

Since the 19th-century unification of electricity and magnetism, and the later development of classical and quantum field theory, physicists have been looking for a single mathematical framework that could describe all fundamental interactions at once. A lot of progress has been made for the electromagnetic, weak and strong forces, which are now successfully unified in the Standard Model. Gravity, however, remains the main stumbling block. Unlike the other forces, it resists quantization in a straightforward way, and its description by General Relativity [2] sits completely apart from the quantum field-theoretic language we use for everything else.

One of the earliest attempts to change this was proposed by Kaluza in 1921 [3] and later refined by Klein: by adding one extra spatial dimension compactified on a small circle, five-dimensional General Relativity naturally reduces to four-dimensional Einstein gravity coupled to electromagnetism. Even if the original theory had problems, this Kaluza-Klein mechanism opened a whole new way of thinking about unification, and it is now at the heart of modern approaches like supergravity [4] and string theory.

2 General relativity theoretical reminder

Here you will find some GR reminder, which hopefully will make it easier to understand the mathematical foundation on which this internship was build

2.1 mathematical basis

Definition 1 Let V be an R -vector space of dimension N , we define the tensor of order $(r, s) \in \llbracket 0, 1 \rrbracket^2$ on $V^{(r,s)} \stackrel{\text{def}}{=} V^{\otimes r} \otimes V^{\otimes s}$ as a being a multilinear application on $V^{\star r} \times V^s$ to $\mathbb{R}$

Definition 2 We will call n -dimension ($n \in \mathbb{N}^*$) manifold every Hausdorff topological space $(\mathcal{M}, \tau)$ locally homeomorphic to $\mathbb{R}^n$

Definition 3 Let $(\mathcal{M}, g)$ be a smooth manifold. A metric tensor is a symmetric, non-degenerate bilinear form $g_{\mu\nu}$ on the tangent space $T_p\mathcal{M}$ at each point $p \in \mathcal{M}$, such that

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu$$

In General Relativity we use the signature $(-, +, +, +)$.

Definition 4 In general in RG we use the christoffels connexion as it is the sole one wich is mtreci and torsion free. We define it as being

$$\text{nabla}_\mu V^\nu = \partial_\mu V^\nu + \Gamma_{\mu\sigma}^\nu V^\sigma \quad (1)$$

where the christoffels symbols are

$$\Gamma_{\mu\nu}^\lambda = \frac{1}{2} g^{\lambda\sigma} (\partial_\mu g_{\nu\sigma} + \partial_\nu g_{\sigma\mu} - \partial_\sigma g_{\mu\nu}) \quad (2)$$

2.2 spin connexion, vielbein and flatspace

3 Lagrangian field theory

3.1 Lagrangian density

In field theory, just as Newton's second law is "volumized" into the Navier-Stokes equations by distributing quantities over space, we replace discrete degrees of freedom with continuous field densities defined at every point in spacetime. Therefore we will rewrite the Lagrangian as the integral over space of a new quantities called Lagrangian density, which will often refered to as just the Lagrangian. Thus we obtain this equation:

$$L = \iiint \mathcal{L}(\eta_\rho, \frac{\partial \eta_\rho}{\partial x^\nu}) dx dy dz \quad (3)$$

Now using esintein convention we can rewritte the action in a new light :

$$S = \int \mathcal{L} dx^\mu \quad (4)$$

3.2 General exemple for scalar and vectorial fields, and result for two forms

In field theory, the Lagrangian has a classical prescription for scalar and vector fields, but no general rule exists for other cases each field type requires its own treatment. For this work, only the scalar, vector, and two-form Lagrangians are needed, along with the metric, which is handled by Einstein's equations.

So for a scalar field we have :

$$\mathcal{L} = -\frac{1}{2}\partial_\mu\Phi\partial^\mu\Phi \quad (5)$$

and for a vectorial field we have

$$\mathcal{L} = -\frac{1}{2}F_{\mu\nu}F^{\mu\nu} \quad (6)$$

where for the field A

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$$

In a vacuum the einstein equation give us

$$\mathcal{L} = R$$

where R is ricci scalar finally for the lagrangian of a two form fields we will look in [5] where it is given by the equation :

$$\mathcal{L} = g^{\hat{\mu}\hat{\mu}'}g^{\nu\nu'}g^{\rho\rho'}H_{\mu\nu\rho}H_{\mu'\nu'\rho'} \quad (7)$$

where for the two form $B_{\nu\rho}$ we have

$$H_{\mu\nu\rho} = \partial_\mu B_{\nu\rho} + cyc.perm$$

4 Reduction of hilbert-einstein Action in D+E dimension

In this section we perform a Kaluza-Klein reduction. We consider a universe living in $D + E$ dimensions, with $D = 4$, and we try to determine how an observer confined to the D -dimensional world perceives the extra dimensions.

4.1 Hypothesis

We assume that none of the fields depend on the internal coordinates, i.e. for any field $\partial_\alpha A^{\hat{\mu}} = 0$. It is also worth noting that Kaluza originally performed his reduction on a circle S^1 ; here, instead, we consider a general internal space of dimension E .

4.2 Rewriting the action

We start from the usual Einstein-Hilbert action, the only difference being that we now work in $D + E$ dimensions. Following [1], this gives

$$S = \frac{-1}{\kappa^2} \int dx^D \int \frac{dy^E}{\rho(E)} \hat{V} \hat{R} \quad (8)$$

where $\rho(E)$ is a normalisation factor introduced in (8), corresponding in practice to the invariant volume of the internal space; κ is fixed such that $\kappa^2/4\pi$ is the usual Newton constant.

Working instead with the spin-connexion formalism, the action can be rewritten in a more transparent form:

$$S = \frac{-1}{\kappa^2} \int dx^D \int \frac{dy^E}{\rho(E)} \hat{V} (\hat{\omega}_{\hat{s}\hat{t}}^{\hat{r}} \hat{\omega}_{\hat{r}}^{\hat{s}\hat{t}} + \hat{\omega}_{\hat{r}\hat{t}}^{\hat{s}} \hat{\omega}_{\hat{s}}^{\hat{r}\hat{t}}) \quad (9)$$

where the spin connexion is given by

$$\hat{\omega}_{\hat{t}\hat{r}\hat{s}} = -\hat{\Omega}_{\hat{t}\hat{r}\hat{s}} + \hat{\Omega}_{\hat{r}\hat{s}\hat{t}} - \hat{\Omega}_{\hat{s}\hat{t}\hat{r}} \quad (10)$$

and the $\hat{\Omega}$ are expressed in terms of the inverse vielbein as

$$\hat{\Omega}_{\hat{r}\hat{s}\hat{t}} = \frac{1}{2}(\hat{V}_{\hat{r}}^{\hat{\mu}}\hat{V}_{\hat{s}}^{\hat{\nu}} - \hat{V}_{\hat{s}}^{\hat{\mu}}\hat{V}_{\hat{r}}^{\hat{\nu}})\partial_{\hat{\nu}}\hat{V}_{\hat{\mu}}^{\hat{t}} \quad (11)$$

4.3 vielbein ansatz

Let us now begin the actual computation. Since all the information about the curved-space metric is contained in the vielbein, we look for an ansatz directly on it. Moreover, following [1], the local Lorentz invariance in $D + E$ dimensions allows us to choose a triangular parametrisation. Our ansatz therefore reads

$$\hat{V}_{\hat{\mu}}^{\hat{r}} = \begin{pmatrix} \delta^\gamma V_\mu^r & 2\kappa A_\mu^\alpha \Phi_\alpha^r \\ 0 & \Phi_\alpha^r \end{pmatrix} \quad (12)$$

In this expression, $\delta = \det \Phi$, and γ is a free parameter corresponding to an arbitrary Weyl rescaling of the D -dimensional vielbein. The vielbein in (24) thus contains the D -dimensional vielbein (i.e. the graviton), E vector fields A_μ^α , and E^2 scalar fields Φ_α^a . However, because of the local Lorentz invariance in the internal directions, only $\frac{1}{2}E(E+1)$ of these scalars actually correspond to propagating degrees of freedom.

4.3.1 Gauge transformations of the fields

Using the fact that the theory is invariant under the classical transformation $x'^{\mu} = x^{\mu} + \xi^{\mu}(x)$ [1], which can be rewritten as

$$\delta \hat{V}_{\hat{\mu}}^{\hat{r}} = \xi^{\hat{\rho}} \partial_{\hat{\rho}} \hat{V}_{\hat{\mu}}^{\hat{r}} + \partial_{\hat{\mu}} \xi^{\hat{\rho}} \hat{V}_{\hat{\rho}}^{\hat{r}} \quad (13)$$

we now apply this transformation to each block of the vielbein, in order to see whether it corresponds to some familiar type of field. Let us start with the block $\hat{V}_{\mu}^r$:

$$\begin{aligned} \delta \hat{V}_{\mu}^r &= \xi^{\hat{\rho}} \partial_{\hat{\rho}} \hat{V}_{\mu}^r + \partial_{\mu} \xi^{\hat{\rho}} \hat{V}_{\hat{\rho}}^r \\ &= \xi^{\hat{\rho}} \partial_{\hat{\rho}} \hat{V}_{\mu}^r + \partial_{\mu} \xi^{\nu} \hat{V}_{\nu}^r + \underbrace{\partial_{\mu} \xi^{\alpha} \hat{V}_{\alpha}^r}_{\text{Ce terme est nul}} \\ &= \xi^{\hat{\rho}} \partial_{\hat{\rho}} (\delta^{\gamma} V_{\mu}^r + \delta^{\gamma} \partial_{\mu} \xi^{\nu} V_{\nu}^r) \\ &= \delta^{\gamma} (\xi^{\hat{\rho}} \partial_{\hat{\rho}} V_{\mu}^r + \partial_{\mu} \xi^{\nu} V_{\nu}^r) + \xi^{\hat{\rho}} \partial_{\hat{\rho}} \delta^{\gamma} V_{\mu}^r \end{aligned}$$

We can also redo this computation using a different method:

$$\begin{aligned} \delta \hat{V}_{\mu}^r &= \delta (\delta^{\gamma} V_{\mu}^r) \\ &= \delta (\delta^{\gamma}) V_{\mu}^r + \delta^{\gamma} \delta (V_{\mu}^r) \end{aligned}$$

and by identification we find

$$\delta V_{\mu}^r = \xi^{\hat{\rho}} \partial_{\hat{\rho}} V_{\mu}^r + \partial_{\mu} \xi^{\nu} V_{\nu}^r$$

which is exactly the transformation law of a vector field under a Lie derivative [4]. Applying the same method to the other blocks, we obtain, for Φ_{α}^a :

$$\delta \Phi_{\alpha}^a = \xi^{\rho} \partial_{\rho} \Phi_{\alpha}^a$$

which is exactly the transformation law of a scalar field. Finally, for A_{μ}^{α} we obtain:

$$A_{\mu}^{\alpha} = \xi^{\rho} \partial_{\rho} A_{\mu}^{\alpha} + \partial_{\mu} \xi^{\rho} A_{\rho}^{\alpha} + \frac{\partial_{\mu} \xi^{\alpha}}{2\kappa}$$

which corresponds to the transformation law of a vector field, up to an extra correction term. One may further note that, restricting to ξ^{α} only, we obtain

$$\delta V_{\mu}^r = \delta \Phi_{\alpha}^a = 0, \quad (14a)$$

$$\delta A_{\mu}^{\alpha} = \frac{\partial_{\mu} \xi^{\alpha}}{2\kappa}. \quad (14b)$$

so that what remains of the D -dimensional diffeomorphisms, once restricted to ξ^{α} independent of the internal coordinates y , corresponds to a set of E abelian gauge symmetries.

4.3.2 Metrics

Looking back at (24), we notice that Φ_{α}^a is in fact nothing but the internal vielbein. We can therefore define the internal metric $h_{\alpha\beta} = \Phi_{\alpha}^a \delta_{ab} \Phi_{\beta}^b$, and finally write down the full $(D + E)$ -dimensional metric $\hat{g}_{\hat{\mu}\hat{\nu}}$ as

$$\hat{g}_{\hat{\mu}\hat{\nu}} = \begin{pmatrix} \delta^{2\gamma} g_{\mu\nu} - 4\kappa^2 A_{\mu}^{\gamma} A_{\nu}^{\lambda} & -2\kappa A_{\mu\beta}^{\gamma} \\ -2\kappa A_{\nu\alpha}^{\gamma} & -h_{\alpha\beta} \end{pmatrix} \quad (15)$$

and, just as forthrightly, the inverse metric

$$\hat{g}^{\hat{\mu}\hat{\nu}} = \begin{pmatrix} \delta^{2\gamma} g^{\mu\nu} & -2\kappa A^{\mu\beta} \\ -2\kappa A^{\nu\alpha} & -h^{\alpha\beta} + 4\kappa^2 A_{\rho}^{\beta} A^{\rho\alpha} \end{pmatrix} \quad (16)$$

4.4 Latin connexion computation

We now need to compute some of the spin-connexion components. To see which ones are actually required, let us rewrite the action in terms of blocks of non-hatted (flat) indices:

$$\begin{aligned} S = -\frac{1}{4\kappa^2} \int d^D x \int \frac{d^E y}{\rho(E)} \hat{V} \Big[& \hat{\omega}_{bc}^a \hat{\omega}^{bc}_a + \hat{\omega}_{bt}^a \hat{\omega}^{bt}_a + \hat{\omega}_{sc}^a \hat{\omega}^{sc}_a + \hat{\omega}_{st}^a \hat{\omega}^{st}_a + \hat{\omega}_{bc}^r \hat{\omega}^{bc}_r + \hat{\omega}_{bt}^r \hat{\omega}^{bt}_r + \hat{\omega}_{sc}^r \hat{\omega}^{sc}_r \\ & \hat{\omega}_{st}^r \hat{\omega}^{st}_r + \hat{\omega}_{ac}^a \hat{\omega}^{bc}_b + \hat{\omega}_{at}^a \hat{\omega}^{bt}_b + \hat{\omega}_{ac}^a \hat{\omega}^{sc}_s + \hat{\omega}_{at}^a \hat{\omega}^{st}_s + \hat{\omega}_{rc}^r \hat{\omega}^{bc}_b + \hat{\omega}_{rt}^r \hat{\omega}^{bt}_b + \hat{\omega}_{rc}^r \hat{\omega}^{sc}_s + \hat{\omega}_{rt}^r \hat{\omega}^{st}_s \Big] \end{aligned} \quad (17)$$

In order to minimise the number of connexion components we need to compute, we use the identity

$$\omega_{rst} = -\omega_{rts} \quad (18)$$

Furthermore, for any pair of up/down indices, one can perform the following manipulation:

$$\omega_{rs}{}^t = \eta^{tn} \omega_{rsn} = -\eta^{tn} \omega_{rns} = -\omega_r{}^t{}_s$$

This gives a convenient trick to reduce, as much as possible, the number of connexion components that must be computed, allowing us to rewrite the action above as

$$S = -\frac{1}{4\kappa^2} \int d^D x \int \frac{d^E y}{\rho(E)} \hat{V} \left[\hat{\omega}^a{}_{bc} \hat{\omega}^{bc}{}_a + \hat{\omega}^a{}_{bt} \hat{\omega}^{bt}{}_a + \hat{\omega}^a{}_{sc} \hat{\omega}^{sc}{}_a + \hat{\omega}^a{}_{st} \hat{\omega}^{st}{}_a - \hat{\omega}^r{}_{bc} \hat{\omega}^b{}_r{}^c - \hat{\omega}^r{}_{tb} \hat{\omega}^{bt}{}_r - \hat{\omega}^r{}_{sc} \hat{\omega}^{cs}{}_r \right. \\ \left. + \hat{\omega}^r{}_{st} \hat{\omega}^{st}{}_r + \hat{\omega}^a{}_{ac} \hat{\omega}^b{}_b{}^c + \hat{\omega}^a{}_{ta} \hat{\omega}^{bt}{}_b + \hat{\omega}^a{}_{ac} \hat{\omega}^s{}_s{}^c - \hat{\omega}^a{}_{ta} \hat{\omega}^s{}_s{}^t + \hat{\omega}^r{}_{rc} \hat{\omega}^b{}_b{}^c - \hat{\omega}^r{}_{rt} \hat{\omega}^{bt}{}_b + \hat{\omega}^r{}_{rc} \hat{\omega}^s{}_s{}^c + \hat{\omega}^r{}_{rt} \hat{\omega}^s{}_s{}^t \right] \quad (19)$$

It then becomes clear that we only need to compute six components of ω , namely

$$\hat{\omega}_{pmn}, \hat{\omega}_{pma}, \hat{\omega}_{pab}, \\ \hat{\omega}_{cmn}, \hat{\omega}_{cma}, \hat{\omega}_{cab},$$

One can also check that computing these six components only requires the following eight components of Ω :

$$\hat{\Omega}_{pmn}, \\ \hat{\Omega}_{pma}, \hat{\Omega}_{pam}, \hat{\Omega}_{apm}, \\ \hat{\Omega}_{mab}, \hat{\Omega}_{amb}, \hat{\Omega}_{abm}, \\ \hat{\Omega}_{abc},$$

Using (10), (11) and (24), we can now set out a clear strategy: first compute $\hat{V}_{\hat{r}}^{\hat{\mu}}$ and $\hat{V}_{\hat{\mu}\hat{r}}$, then the Ω 's, and finally the ω 's. The inverse vielbein is easy to obtain, since

$$\hat{V}_{\hat{\mu}}^{\hat{s}} \hat{V}_{\hat{r}}^{\hat{\mu}} = \delta_{\hat{r}}^{\hat{s}}$$

so that

$$\hat{V}_{\hat{\mu}}^{\hat{r}} = \begin{pmatrix} \delta^{-\gamma} V_r^\mu & -2\kappa \delta^{-\gamma} A_r^\alpha \\ 0 & \Phi_a^\alpha \end{pmatrix} \quad (20)$$

Similarly, for $\hat{V}_{\hat{\mu}\hat{r}}$, using

$$\hat{V}_{\hat{r}}^{\hat{\mu}} \hat{V}_{\hat{\mu}\hat{s}} = \eta_{\hat{r}\hat{s}}$$

we obtain

$$\hat{V}_{\hat{\mu}\hat{s}} = \begin{pmatrix} \delta^\gamma V_{\mu s} & 2\kappa A_\mu^\alpha \Phi_{\alpha a} \\ 0 & \Phi_{\alpha a} \end{pmatrix} \quad (21)$$

Let us now turn to the computation of the Ω 's themselves. Starting with $\hat{\Omega}_{pmn}$:

$$\begin{aligned} \hat{\Omega}_{pmn} &= \frac{1}{2} (\hat{V}_p^{\hat{\mu}} \hat{V}_m^\nu - \hat{V}_m^{\hat{\mu}} \hat{V}_p^\nu) \partial_\nu \hat{V}_{\hat{\mu}n} \\ &= \frac{1}{2} (\hat{V}_p^\mu \hat{V}_m^\nu - \hat{V}_m^\mu \hat{V}_p^\nu) \partial_\nu \hat{V}_{\hat{\mu}n} \\ &= \frac{1}{2} (\hat{V}_p^\mu \hat{V}_m^\nu - \hat{V}_m^\mu \hat{V}_p^\nu) \partial_\nu \hat{V}_{\mu n} \\ &= \frac{\delta^{-2\gamma}}{2} (V_p^\mu V_m^\nu - V_m^\mu V_p^\nu) \partial_\nu (\delta^\gamma V_{\mu n}) \\ &= \frac{\delta^{-2\gamma}}{2} (V_p^\mu V_m^\nu - V_m^\mu V_p^\nu) (\partial_\nu (\delta^\gamma) V_{\mu n} + \delta^\gamma \partial_\nu (V_{\mu n})) \\ &= \frac{\delta^{-2\gamma}}{2} (V_p^\mu V_m^\nu - V_m^\mu V_p^\nu) (\gamma \partial_\nu (\delta) \delta^{\gamma-1} V_{\mu n} + \delta^\gamma \partial_\nu (V_{\mu n})) \\ &= \frac{\delta^{-\gamma}}{2} (V_p^\mu V_m^\nu - V_m^\mu V_p^\nu) (\gamma \partial_\nu (\delta) \delta^{-1} V_{\mu n} + \partial_\nu (V_{\mu n})) \\ &= \frac{\delta^{-\gamma}}{2} (V_p^\mu V_m^\nu - V_m^\mu V_p^\nu) (\gamma \partial_\nu (\log \delta) V_{\mu n} + \partial_\nu (V_{\mu n})) \end{aligned}$$

For $\hat{\Omega}_{pma}$, we find:

$$\begin{aligned}
\hat{\Omega}_{pma} &= \frac{1}{2}(\hat{V}_p^{\hat{\mu}}\hat{V}_m^{\nu} - \hat{V}_m^{\hat{\mu}}\hat{V}_p^{\nu})\partial_{\nu}\hat{V}_{\hat{\mu}a} \\
&= -\kappa\delta^{-2\gamma}(V_p^{\mu}V_m^{\nu} - V_m^{\mu}V_p^{\nu})\partial_{\nu}(A_{\mu}^{\alpha}\Phi_{a\alpha}) + \kappa\delta^{-\gamma}(A_p^{\alpha}V_m^{\nu} - \hat{A}_m^{\alpha}V_p^{\nu})\partial_{\nu}(\Phi_{a\alpha}) \\
&= -\kappa\delta^{-2\gamma}(V_p^{\mu}V_m^{\nu} - V_m^{\mu}V_p^{\nu})(\partial_{\nu}(A_{\mu}^{\alpha})\Phi_{a\alpha} + \partial_{\nu}(\Phi_{a\alpha})A_{\mu}^{\alpha}) + \kappa\delta^{-\gamma}(A_p^{\alpha}V_m^{\nu} - \hat{A}_m^{\alpha}V_p^{\nu})\partial_{\nu}(\Phi_{a\alpha}) \\
&= -\kappa\delta^{-2\gamma}(V_p^{\mu}V_m^{\nu} - V_m^{\mu}V_p^{\nu})(\partial_{\nu}(A_{\mu}^{\alpha})\Phi_{a\alpha} + \partial_{\nu}(\Phi_{a\alpha})A_{\mu}^{\alpha}) + (A_p^{\alpha}V_m^{\nu} - \hat{A}_m^{\alpha}V_p^{\nu})\partial_{\nu}(\Phi_{a\alpha}) \\
&= -\kappa\delta^{-2\gamma}(V_p^{\mu}V_m^{\nu} - V_m^{\mu}V_p^{\nu})\partial_{\nu}(A_{\mu}^{\alpha})\Phi_{a\alpha} \\
&= \kappa\delta^{-2\gamma}(F^{\alpha}_{pm}\Phi_{a\alpha} - \partial_{\nu}(V_p^{\mu}V_m^{\nu})A_{\mu}^{\alpha}\Phi_{a\alpha} + \partial_{\nu}(V_p^{\nu}V_m^{\mu})A_{\mu}^{\alpha}\Phi_{a\alpha}) \\
&= \kappa\delta^{-2\gamma}F^{\alpha}_{pm}\Phi_{a\alpha}
\end{aligned}$$

For $\hat{\Omega}_{pam}$:

$$\begin{aligned}
\hat{\Omega}_{pam} &= \frac{1}{2}(\hat{V}_p^{\hat{\mu}}\hat{V}_a^{\nu} - \hat{V}_a^{\hat{\mu}}\hat{V}_p^{\nu})\partial_{\nu}\hat{V}_{\hat{\mu}m} \\
&= \frac{1}{2}(\hat{V}_p^{\hat{\mu}}\hat{V}_a^{\nu} - \hat{V}_a^{\hat{\mu}}\hat{V}_p^{\nu})\partial_{\nu}\hat{V}_{\hat{\mu}m} \\
&= 0
\end{aligned}$$

The same result holds for $\hat{\Omega}_{apm}$.

For $\hat{\Omega}_{pab}$:

$$\begin{aligned}
\hat{\Omega}_{pab} &= \frac{1}{2}(\hat{V}_p^{\hat{\mu}}\hat{V}_a^{\nu} - \hat{V}_a^{\hat{\mu}}\hat{V}_p^{\nu})\partial_{\nu}\hat{V}_{\hat{\mu}b} \\
&= \frac{-\delta^{-\gamma}}{2}\hat{V}_a^{\hat{\mu}}V_p^{\nu}\partial_{\nu}\hat{V}_{\hat{\mu}b} \\
&= \frac{\delta^{-\gamma}}{2}\Phi_a^{\alpha}V_p^{\nu}\partial_{\nu}\Phi_{\alpha b} \\
&= \frac{\delta^{-\gamma}}{2}\Phi_a^{\alpha}V_p^{\nu}\partial_{\nu}(h_{\alpha\beta}\Phi_b^{\beta}) \\
&= \frac{\delta^{-\gamma}}{2}\Phi_a^{\alpha}\Phi_b^{\beta}V_p^{\nu}\partial_{\nu}h_{\alpha\beta} + \frac{\delta^{-\gamma}}{2}\Phi_a^{\alpha}V_p^{\nu}h_{\alpha\beta}\partial_{\nu}\Phi_b^{\beta} \\
&= \frac{\delta^{-\gamma}}{2}\Phi_a^{\alpha}\Phi_b^{\beta}V_p^{\nu}\partial_{\nu}h_{\alpha\beta} + \frac{\delta^{-\gamma}}{2}\Phi_{\beta a}V_p^{\nu}\partial_{\nu}\Phi_b^{\beta} \\
&= \frac{\delta^{-\gamma}}{2}\Phi_a^{\alpha}\Phi_b^{\beta}V_p^{\nu}\partial_{\nu}h_{\alpha\beta} + \frac{\delta^{-\gamma}}{2}V_p^{\nu}\partial_{\nu}(\underbrace{\Phi_{\beta a}\Phi_b^{\beta}}_{=\delta_{ab} \text{ i.e is constant}}) - \Phi_b^{\beta}V_p^{\nu}\partial_{\nu}\Phi_{\beta a} \\
&= \frac{\delta^{-\gamma}}{2}\Phi_a^{\alpha}\Phi_b^{\beta}V_p^{\nu}\partial_{\nu}h_{\alpha\beta} - \Phi_b^{\beta}V_p^{\nu}\partial_{\nu}\Phi_{\beta a} \\
&=
\end{aligned}$$

For $\hat{\Omega}_{apb}$:

$$\begin{aligned}
\hat{\Omega}_{apb} &= \frac{1}{2}(\hat{V}_a^{\hat{\mu}}\hat{V}_p^{\nu} - \hat{V}_p^{\hat{\mu}}\hat{V}_a^{\nu})\partial_{\nu}\hat{V}_{\hat{\mu}b} \\
&= \frac{1}{2}\hat{V}_a^{\hat{\mu}}\hat{V}_p^{\nu}\partial_{\nu}\hat{V}_{\hat{\mu}b} \\
&= \frac{1}{2}\hat{V}_a^{\alpha}\hat{V}_p^{\nu}\partial_{\nu}\hat{V}_{\alpha b} \\
&= \frac{-\delta^{-\gamma}}{2}\Phi_a^{\alpha}V_p^{\nu}\partial_{\nu}\Phi_{\alpha b}
\end{aligned}$$

For $\hat{\Omega}_{abp}$:

$$\begin{aligned}
\hat{\Omega}_{abp} &= \frac{1}{2}(\hat{V}_a^{\hat{\mu}}\hat{V}_b^{\nu} - \hat{V}_b^{\hat{\mu}}\hat{V}_a^{\nu})\partial_{\nu}\hat{V}_{\hat{\mu}p} \\
&= 0
\end{aligned}$$

The same holds for Ω_{abc} . Using (10), we can then obtain the ω 's straightforwardly:

$$\hat{\omega}_{pmn} = \delta^{-\gamma} [\omega_{pmn} + \gamma(\eta_{pm} V_n^\lambda - \eta_{pn} V_m^\lambda) \partial_\lambda], \quad (22a)$$

$$\hat{\omega}_{pma} = \kappa \delta^{-2\gamma} F_{pm}^\alpha \Phi_{\alpha a}, \quad (22b)$$

$$\hat{\omega}_{pab} = \frac{1}{2} \delta^{-\gamma} \Phi_a^\alpha V_p^\mu \partial_\mu \Phi_{\alpha\beta} - (a \leftrightarrow b), \quad (22c)$$

$$\hat{\omega}_{cmn} = -\kappa \delta^{-2\gamma} F_{mn}^\alpha \Phi_{\alpha c}, \quad (22d)$$

$$\hat{\omega}_{cab} = 0 \quad (22e)$$

where $F_{\mu\nu}^\alpha = \partial_\mu A_\nu^\alpha - \partial_\nu A_\mu^\alpha$.

The gauge invariance of these expressions is then manifest.

4.5 finalisation

Now that all this work has been done we can finally begin the computation of the action

$$S = \frac{-1}{4\kappa^2} \int dx^D \int \frac{dy^E}{\rho(E)} V \delta^{\gamma D+1} \left(\frac{1}{4} \delta^{-2\gamma} g^{\rho\lambda} \partial_\lambda h_{\alpha\beta} \partial_\rho h^{\alpha\beta} + \kappa^2 \delta^{-2\gamma} F^{\mu\nu\alpha} F_{\mu\nu}^\beta h_{\alpha\beta} + \right. \\ \left. \delta^{-2\gamma} (\omega_{st}^r \omega_r^{st} + \gamma^2 (2-D) g^{\lambda\rho} \partial_\rho \log \delta \partial_\lambda \log \delta) + \delta^{-2\gamma} (\omega_{rt}^r \omega_s^s{}^t) \right)$$

or rewritten in a nicer way

$$S = \int dx^D \int \frac{dy^E}{\rho(E)} V \delta^{\gamma D+1} \left(\frac{-1}{16} \delta^{-2\gamma} g^{\rho\lambda} \partial_\lambda h_{\alpha\beta} \partial_\rho h^{\alpha\beta} + \frac{-1}{4} \delta^{-2\gamma} F^{\mu\nu\alpha} F_{\mu\nu}^\beta h_{\alpha\beta} + \right. \\ \left. \frac{-1}{4\kappa^2} \delta^{-2\gamma} \gamma^2 (2-D) g^{\lambda\rho} \partial_\rho \log \delta \partial_\lambda \log \delta + \frac{-1}{4\kappa^2} \delta^{-2\gamma} (\omega_{st}^r \omega_r^{st} + \omega_{rt}^r \omega_s^s{}^t) \right)$$

so now we can easily identify R the D-dimension einstein action and see the factor $\delta^{\gamma(D-2)+1}$ the good thing is that γ is a free variable so I can just say that

$$S = \int dx^D V \delta^{\gamma D+1} \left(-\frac{1}{4\kappa^2} R + \frac{-1}{4} \delta^{\frac{D-2}{2}} F^{\mu\nu\alpha} F_{\mu\nu}^\beta h_{\alpha\beta} + \right. \\ \left. -\frac{1}{16} g^{\rho\lambda} \partial_\lambda h_{\alpha\beta} \partial_\rho h^{\alpha\beta} + \frac{1}{4\kappa^2(D-2)} \gamma^2 g^{\lambda\rho} \partial_\rho \log \delta \partial_\lambda \log \delta \right)$$

5 reduction of the new fields

Now that we have seen the classical reduction of the einstein action an interesting exercise we can do is to add two new fields namely the dilaton field and a second rank antisymmetric tensor as done in [5]. We will once again follow the same path for the reduction knowing that for this part [5] did follow the instruction of [1]. The main change in both papers is that [5] choose $\gamma = 0$ and not $\gamma = -\frac{1}{D-2}$, which as said earlier is just a choice who does not carry a lot of importance. [5] also chose to ignore the κ which gave us a modified action in comparison to our last result. Finally contrary to [1], [5] didn't choose the $(+, -, -, -, \dots)$ signature for the tangent space metric but $(-, +, +, +, \dots)$, so there will be some change of sign in the action. Once everything is taken into account we get the new reduced action

$$S = \int dx^D \sqrt{-g} \left(R - \frac{1}{4} F^{\mu\nu\alpha} F_{\mu\nu}^\beta h_{\alpha\beta} + \right. \\ \left. - g^{\rho\lambda} \partial_\lambda h_{\alpha\beta} \partial_\rho h^{\alpha\beta} - \frac{1}{2} g^{\lambda\rho} \partial_\rho \log \delta \partial_\lambda \log \delta \right) \quad (23)$$

and of course we can't forget to rewrite the vielbein and the high dimension metric

$$\hat{V}_{\hat{\mu}}^{\hat{r}} = \begin{pmatrix} V_\mu^r & A_\mu^\alpha \Phi_\alpha^a \\ 0 & \Phi_\alpha^a \end{pmatrix} \quad \text{and} \quad \hat{V}_{\hat{\mu}}^{\hat{r}} = \begin{pmatrix} V_r^\mu & -A_r^\alpha \\ 0 & \Phi_a^\alpha \end{pmatrix} \quad (24)$$

and the metric

$$\hat{g}_{\hat{\mu}\hat{\nu}} = \begin{pmatrix} g_{\mu\nu} + A_\mu^\gamma A_\nu^\lambda & A_{\mu\beta} \\ A_{\nu\alpha} & h_{\alpha\beta} \end{pmatrix} \quad \text{and} \quad \hat{g}^{\hat{\mu}\hat{\nu}} = \begin{pmatrix} g^{\mu\nu} & -A^{\mu\beta} \\ -A^{\nu\alpha} & -h^{\alpha\beta} + A_\rho^\beta A^{\rho\alpha} \end{pmatrix} \quad (25)$$

As a final step before beginning we can see that

$$\sqrt{-g} = \det \hat{V}_{\hat{\mu}}^{\hat{r}} = \det V_\mu^r \det \Phi_\alpha^a = \sqrt{-g} \sqrt{h} \quad (26)$$

5.1 scalar field reduction

Now let's add the dilaton field by modifying like this the original action

$$S = \int dx^D \int \frac{dy^E}{\rho(E)} \sqrt{-\hat{g}} e^{-\hat{\phi}} (\hat{R} + \hat{g}^{\hat{\mu}\hat{\nu}} \partial_{\hat{\mu}} \hat{\phi} \partial_{\hat{\nu}} \hat{\phi}) \quad (27)$$

We can then do what we did with the [1] and get and other manipulation

$$S = \int dx^D \sqrt{-g} \sqrt{h} e^{-\hat{\phi}} \left(R - \frac{1}{4} F^{\mu\nu\alpha} F_{\mu\nu}^{\beta} h_{\alpha\beta} + \right. \\ \left. - g^{\rho\lambda} \partial_{\lambda} h_{\alpha\beta} \partial_{\rho} h^{\alpha\beta} - \frac{1}{2} g^{\lambda\rho} \partial_{\rho} \log \delta \partial_{\lambda} \log \delta + \hat{g}^{\hat{\mu}\hat{\nu}} \partial_{\hat{\mu}} \hat{\phi} \partial_{\hat{\nu}} \hat{\phi} \right) \quad (28)$$

5.2 Two form fiel reduction

6 Observation of new symmetry

7 Conclusion

References

- [1] Joel Scherk and John H Schwarz. “How to get masses from extra dimensions”. In: *Nuclear Physics B* 153 (1979), pp. 61–88.
- [2] Albert Einstein. “Zur allgemeinen relativitätstheorie”. In: *Sitzungsber. preuss. Akad. Wiss* 44.2 (1915), pp. 778–786.
- [3] Th Kaluza. “Zum unitätsproblem der physik”. In: *Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.)* 1921.arXiv: 1803.08616 (1921), pp. 966–972.
- [4] Daniel Z Freedman and Antoine Van Proeyen. *Supergravity*. Cambridge university press, 2012.
- [5] Jnanadeva Maharana and John H Schwarz. “Noncompact symmetries in string theory”. In: *Nuclear Physics B* 390.1 (1993), pp. 3–32.

A lagrangian

By using the principle of least action and the variational principle we can see that

$$\begin{aligned} \delta S &= \delta \int \mathcal{L} dx^{\mu} \\ &= \delta \int \frac{\partial \mathcal{L}}{\partial \eta_{\rho}} \delta \eta_{\rho} + \frac{\partial \mathcal{L}}{\partial \partial_{\nu} \eta_{\rho}} \delta \partial_{\nu} \eta_{\rho} dx^{\mu} \\ &= \delta \int \frac{\partial \mathcal{L}}{\partial \eta_{\rho}} \delta \eta_{\rho} - \partial_{\nu} \frac{\partial \mathcal{L}}{\partial \partial_{\nu} \eta_{\rho}} \delta \eta_{\rho} + \underbrace{\partial_{\nu} \left(\frac{\partial \mathcal{L}}{\partial \partial_{\nu} \eta_{\rho}} \delta \eta_{\rho} \right)}_{\text{border terme equal to zero}} dx^{\mu} \\ &= 0 \end{aligned}$$

which gave the equation

$$\boxed{\frac{\partial \mathcal{L}}{\partial \eta_{\rho}} - \partial_{\nu} \frac{\partial \mathcal{L}}{\partial \partial_{\nu} \eta_{\rho}} = 0} \quad (29)$$

which is in fact the equation of mouvement for the field