


Spatial dimension compactification in general relativity and duality symmetry

Adam Franchot-Fowler

Internship carried out from 08/06/2026 to 24/07/2026



Internship tutors : M. Camille Eloy

Referent teacher : M. Stéphane Douin

Éducational institution : Université Paris-Saclay / Magistère de physique fondamentale d'Orsay

Internship host company : Laboratoire de Physique (UMR CNRS 5672) ENS de Lyon 46, allée d'Italie F-69364 LYON CEDEX 07, FRANCE

Acknowledgment

Notations and abbreviations

For the abbreviations we will use

- GR for general relativity

We will during this internship use the vielbein formalism. We will use the same notation as in [1] which is

- Greek hatted letter for curved indices ($\hat{\mu}, \hat{\nu}, \hat{\rho}, \dots$)
- Latin hatted letter for flat indices ($\hat{p}, \hat{m}, \hat{n}, \dots$)
- For space-time indices taking D values, late greek letters are used in the curved case ($\mu, \nu, \rho, \dots$) and Latin letters in the flat case (p, m, n)
- For internal indices taking E values, early Greek letters in the curved case ($\alpha, \beta, \gamma, \dots$) and Latin letters in the flat case (a,b,c)

The general coordinate $x^{\hat{\mu}}$ will therefore be made of space time coordinate x^{μ} and internal coordinate y^{α} . The flat lorentz

Contents

Notation and abbreviation	1
1 Introduction	3
2 General relativity theoretical reminder	3
2.1 mathematical basis	3
2.2 spin connexion, vielbein and flatspace	3
3 Lagrangian field theory	3
3.1 Lagrangian density	3
3.2 General exemple for scalar and vectorial fields, and result for two forms	4
4 Reduction of hilbert-einstein Action in D+E dimension	4
4.1 Hypothesis	4
4.2 Rewriting the action	4
4.3 vielbein anstazs	5
4.3.1 Fields Gauge	5
4.3.2 Metrics	5
4.4 Latin connexion computation	6
4.5 finalisation	8
5 reduction of the new fields	8
5.1 scalar field reduction	8
5.2 Two form fiel reduction	8
6 Observation of new symmetry	8
7 Conclusion	8
A lagrangian	8

1 Introduction

Since the 19th-century unification of electricity and magnetism, and the later development of classical and quantum field theory, physicists have been looking for a single mathematical framework that could describe all fundamental interactions at once. A lot of progress has been made for the electromagnetic, weak and strong forces, which are now successfully unified in the Standard Model. Gravity, however, remains the main stumbling block. Unlike the other forces, it resists quantization in a straightforward way, and its description by General Relativity [2] sits completely apart from the quantum field-theoretic language we use for everything else.

One of the earliest attempts to change this was proposed by Kaluza in 1921 [3] and later refined by Klein: by adding one extra spatial dimension compactified on a small circle, five-dimensional General Relativity naturally reduces to four-dimensional Einstein gravity coupled to electromagnetism. Even if the original theory had problems, this Kaluza-Klein mechanism opened a whole new way of thinking about unification, and it is now at the heart of modern approaches like supergravity [4] and string theory.

2 General relativity theoretical reminder

Here you will find some GR reminder, which hopefully will make it easier to understand the mathematical foundation on which this internship was build

2.1 mathematical basis

Definition 1 Let V be an R -vector space of dimension N , we define the tensor of order $(r, s) \in \llbracket 0, 1 \rrbracket^2$ on $V^{(r,s)} \stackrel{\text{def}}{=} V^{\otimes r} \otimes V^{\otimes s}$ as a being a multilinear application on $V^{\star r} \times V^s$ to $\mathbb{R}$

Definition 2 We will call n -dimension ($n \in \mathbb{N}^*$) manifold every Hausdorff topological space $(\mathcal{M}, \tau)$ locally homeomorphic to $\mathbb{R}^n$

Definition 3 Let $(\mathcal{M}, g)$ be a smooth manifold. A metric tensor is a symmetric, non-degenerate bilinear form $g_{\mu\nu}$ on the tangent space $T_p\mathcal{M}$ at each point $p \in \mathcal{M}$, such that

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu$$

In General Relativity we use the signature $(-, +, +, +)$.

Definition 4 In general in RG we use the christofells connexion as it is the sole one wich is mtreci and torsion free. We define it as being

$$\text{nabla}_\mu V^\nu = \partial_\mu V^\nu + \Gamma_{\mu\sigma}^\nu V^\sigma \quad (1)$$

where the christofells symbols are

$$\Gamma_{\mu\nu}^\lambda = \frac{1}{2} g^{\lambda\sigma} (\partial_\mu g_{\nu\sigma} + \partial_\nu g_{\sigma\mu} - \partial_\sigma g_{\mu\nu}) \quad (2)$$

2.2 spin connexion, vielbein and flatspace

3 Lagrangian field theory

3.1 Lagrangian density

In field theory, just as Newton's second law is "volumized" into the Navier-Stokes equations by distributing quantities over space, we replace discrete degrees of freedom with continuous field densities defined at every point in spacetime. Therefore we will rewrite the Lagrangian as the integral over space of a new quantities called Lagrangian density, which will often refered to as just the Lagrangian. Thus we obtain this equation:

$$L = \iiint \mathcal{L}(\eta_\rho, \frac{\partial \eta_\rho}{\partial x^\nu}) dx dy dz \quad (3)$$

Now using esintein convention we can rewritte the action in a new light :

$$S = \int \mathcal{L} dx^\mu \quad (4)$$

3.2 General exemple for scalar and vectorial fields, and result for two forms

In field theory, the Lagrangian has a classical prescription for scalar and vector fields, but no general rule exists for other cases each field type requires its own treatment. For this work, only the scalar, vector, and two-form Lagrangians are needed, along with the metric, which is handled by Einstein's equations.

So for a scalar field we have :

$$\mathcal{L} = -\frac{1}{2}\partial_\mu\Phi\partial^\mu\Phi \quad (5)$$

and for a vectorial field we have

$$\mathcal{L} = -\frac{1}{2}F_{\mu\nu}F^{\mu\nu} \quad (6)$$

where for the field A

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$$

In a vacuum the Einstein equation give us

$$\mathcal{L} = R$$

where R is Ricci scalar finally for the Lagrangian of a two form fields we will look in [5] where it is given by the equation :

$$\mathcal{L} = g^{\hat{\mu}\mu'}g^{\nu\nu'}g^{\rho\rho'}H_{\mu\nu\rho}H_{\mu'\nu'\rho'} \quad (7)$$

where for the two form $B_{\nu\rho}$ we have

$$H_{\mu\nu\rho} = \partial_\mu B_{\nu\rho} + cyc.perm$$

4 Reduction of Hilbert-Einstein Action in D+E dimension

Here I will do a Kaluza-Klein reduction. Basically we will suppose an universe in D+E dimension where D is 4 and we will try to see how an observer in D will perceive the new dimension.

4.1 Hypothesis

We will consider that the fields don't depend on the internal variable of the world so for any field $\partial_\alpha A^{\hat{\mu}} = 0$. Another thing to note is that originally Kaluza did a reduction on S^1 , but here we will not do that and we will do the reduction on a nondescript space, with E dimensions.

4.2 Rewriting the action

We began our journey with the classical Einstein action; with the sole difference being that we are in D+E dimension so we have as shown in [1]

$$S = \frac{-1}{\kappa^2} \int dx^D \int \frac{dy^E}{\rho(E)} \hat{V} \hat{R} \quad (8)$$

where $\rho(E)$ is the parameter set to normalise (8) and is in practice the invariant volume of the internal space. and we have set κ so that $\frac{\kappa^2}{4\pi}$ is the usual Newton constant.

Now working with spin connection formalism we can rewrite everything in a more understandable manner and we got the action. and get

$$S = \frac{-1}{\kappa^2} \int dx^D \int \frac{dy^E}{\rho(E)} \hat{V} (\hat{\omega}_{\hat{s}\hat{t}}^{\hat{r}} \hat{\omega}_{\hat{r}}^{\hat{s}\hat{t}} + \hat{\omega}_{\hat{r}\hat{t}}^{\hat{s}} \hat{\omega}_{\hat{s}}^{\hat{r}\hat{t}}) \quad (9)$$

Where the spin connection are given by

$$\hat{\omega}_{\hat{t}\hat{r}\hat{s}} = -\hat{\Omega}_{\hat{t}\hat{r}\hat{s}} + \hat{\Omega}_{\hat{r}\hat{s}\hat{t}} - \hat{\Omega}_{\hat{s}\hat{t}\hat{r}} \quad (10)$$

and the $\hat{\Omega}$ are given by the inverse vielbein

$$\hat{\Omega}_{\hat{r}\hat{s}\hat{t}} = \frac{1}{2} (\hat{V}_{\hat{r}}^{\hat{\mu}} \hat{V}_{\hat{s}}^{\hat{\nu}} - \hat{V}_{\hat{s}}^{\hat{\mu}} \hat{V}_{\hat{r}}^{\hat{\nu}}) \partial_{\hat{\nu}} \hat{V}_{\hat{\mu}\hat{t}} \quad (11)$$

4.3 vielbein anstazs

Now let's begin the true work. Firstly as the information of the curved space metric is fully contained within the vielbein we will do the anstaz of the solution on them. Furthermore according to [1] we can choose a triangular parametrization thanks to the local lorentz invariance in D+E dimension. So our anstazs will be

$$\hat{V}_{\hat{\mu}}^{\hat{r}} = \begin{pmatrix} \delta^{\gamma} V_{\mu}^r & 2\kappa A_{\mu}^{\alpha} \Phi_{\alpha}^a \\ 0 & \Phi_{\alpha}^a \end{pmatrix} \quad (12)$$

In this formula $\delta = \det \Phi$ and γ is a free parameter, corresponding to an arbitrary Weyl transformation of the D dimensional vielbein.

The vielbein in eq (12) has D dimensional vielbein (also called graviton), E vector fields A_{μ}^{α} and E^2 , however the local lorentz invariance in the internal direction, only $\frac{1}{2}E(E+1)$ of the scalar fields correspond to propagating particles.

4.3.1 Fields Gauge

Using the fact that the theory is invariant under the classical transformation $x'^{\mu} = x^{\mu} + \xi^{\mu} u(x)$ [1] which can be rewritten as being

$$\delta \hat{V}_{\hat{\mu}}^{\hat{r}} = \xi^{\hat{\rho}} \partial_{\hat{\rho}} \hat{V}_{\hat{\mu}}^{\hat{r}} + \partial_{\hat{\mu}} \xi^{\hat{\rho}} \hat{V}_{\hat{\rho}}^{\hat{r}} \quad (13)$$

Now we will apply this to each block of the vielbein to see if they respect some type of classical field so for the block $\hat{V}_{\mu}^r$

$$\begin{aligned} \delta \hat{V}_{\mu}^r &= \xi^{\hat{\rho}} \partial_{\hat{\rho}} \hat{V}_{\mu}^r + \partial_{\mu} \xi^{\hat{\rho}} \hat{V}_{\hat{\rho}}^r \\ &= \xi^{\hat{\rho}} \partial_{\hat{\rho}} \hat{V}_{\mu}^r + \partial_{\mu} \xi^{\nu} \hat{V}_{\nu}^r + \underbrace{\partial_{\mu} \xi^{\alpha} \hat{V}_{\alpha}^r}_{\text{Ce terme est nul}} \\ &= \xi^{\hat{\rho}} \partial_{\hat{\rho}} (\delta^{\gamma} V_{\mu}^r + \delta^{\gamma} \partial_{\mu} \xi^{\nu} V_{\nu}^r) \\ &= \delta^{\gamma} (\xi^{\hat{\rho}} \partial_{\hat{\rho}} V_{\mu}^r + \partial_{\mu} \xi^{\nu} V_{\nu}^r) + \xi^{\hat{\rho}} \partial_{\hat{\rho}} \delta^{\gamma} V_{\mu}^r \end{aligned}$$

Now we can redo this by this time using another method

$$\begin{aligned} \delta \hat{V}_{\mu}^r &= \delta(\delta^{\gamma} V_{\mu}^r) \\ &= \delta(\delta^{\gamma}) V_{\mu}^r + \delta^{\gamma} \delta(V_{\mu}^r) \end{aligned}$$

and by identification we can see that

$$\delta V_{\mu}^r = \xi^{\hat{\rho}} \partial_{\hat{\rho}} V_{\mu}^r + \partial_{\mu} \xi^{\nu} V_{\nu}^r$$

which is exactly the same transformation as a vector with Lie derivative [4] Now using the same method for the other block we get for the $\hat{V}_{\alpha}^a$

$$\delta \Phi_{\alpha}^a = \xi^{\hat{\rho}} \partial_{\hat{\rho}} \Phi_{\alpha}^a$$

which is exactly the transformation of a scalar field. Finally for the V_{μ}^a we got

$$A_{\mu}^{\alpha} = \xi^{\hat{\rho}} \partial_{\hat{\rho}} A_{\mu}^{\alpha} + \partial_{\mu} \xi^{\rho} A_{\rho}^{\alpha} + \frac{\partial_{\mu} \xi^{\alpha}}{2\kappa}$$

which is also the transformation of a vector field plus a correction term at the end. moreover another one might see is that if we vary ϵ^{α} we got

$$\delta V_{\mu}^r = \delta \Phi_{\alpha}^a = 0, \quad (14a)$$

$$\delta A_{\mu}^{\alpha} = \frac{\partial_{\mu} \xi^{\alpha}}{2\kappa}. \quad (14b)$$

so what remains of the D-dimensional ξ^{α} restricted to y-independent transformation is a set of E abelian Gauge invariance

4.3.2 Metrics

In (12) we may remark that Φ_{α}^a is in fact the internal vielbein therefore we can create the internal metric h with $h_{\alpha\beta} = \Phi_{\alpha}^a \delta_{ab} \Phi_{\beta}^b$ and finally we can write the D+E metric $\hat{g}_{\hat{\mu}\hat{\nu}}$ with

$$\hat{g}_{\hat{\mu}\hat{\nu}} = \begin{pmatrix} \delta^{2\gamma} g_{\mu\nu} - 4\kappa^2 A_{\mu}^{\gamma} A_{\nu}^{\lambda} & -2\kappa A_{\mu\beta}^{\gamma} \\ -2\kappa A_{\nu\alpha}^{\gamma} & -h_{\alpha\beta} \end{pmatrix} \quad (15)$$

and pretty easily the inverse metrics

$$\hat{g}^{\hat{\mu}\hat{\nu}} = \begin{pmatrix} \delta^{2\gamma} g^{\mu\nu} & -2\kappa A^{\mu\beta} \\ -2\kappa A^{\nu\alpha} & -h^{\alpha\beta} + 4\kappa^2 A_{\rho}^{\beta} A^{\rho\alpha} \end{pmatrix} \quad (16)$$

4.4 Latin connexion computation

Now we need to calculate somme spin connexion to discover which one let's rewrite the action by block of non-hatted index that gave us

$$S = -\frac{1}{4\kappa^2} \int d^D x \int \frac{d^E y}{\rho(E)} \hat{V} \left[\hat{\omega}^a_{bc} \hat{\omega}^{bc}_a + \hat{\omega}^a_{bt} \hat{\omega}^{bt}_a + \hat{\omega}^a_{sc} \hat{\omega}^{sc}_a + \hat{\omega}^a_{st} \hat{\omega}^{st}_a + \hat{\omega}^r_{bc} \hat{\omega}^{bc}_r + \hat{\omega}^r_{bt} \hat{\omega}^{bt}_r + \hat{\omega}^r_{sc} \hat{\omega}^{sc}_r \right. \\ \left. \hat{\omega}^r_{st} \hat{\omega}^{st}_r + \hat{\omega}^a_{ac} \hat{\omega}^b_{b^c} + \hat{\omega}^a_{at} \hat{\omega}^b_{b^t} + \hat{\omega}^a_{ac} \hat{\omega}^s_{s^c} + \hat{\omega}^a_{at} \hat{\omega}^s_{s^t} + \hat{\omega}^r_{rc} \hat{\omega}^b_{b^c} + \hat{\omega}^r_{rt} \hat{\omega}^b_{b^t} + \hat{\omega}^r_{rc} \hat{\omega}^s_{s^c} + \hat{\omega}^r_{rt} \hat{\omega}^s_{s^t} \right] \quad (17)$$

now to minimize the number of connexion we have to calculate we will use the fact that

$$\omega_{rst} = -\omega_{rts} \quad (18)$$

furthermore on can with any kind of updown index do the next manipulation

$$\omega_{rs}{}^t = \eta^{tn} \omega_{rsn} = -\eta^{tn} \omega_{rns} = -\omega_r{}^t{}_s$$

A neat trick wich can be done with that is to reduce to a maxim the number of connection we have to compute so we rewrote the thing above as being

$$S = -\frac{1}{4\kappa^2} \int d^D x \int \frac{d^E y}{\rho(E)} \hat{V} \left[\hat{\omega}^a_{bc} \hat{\omega}^{bc}_a + \hat{\omega}^a_{bt} \hat{\omega}^{bt}_a + \hat{\omega}^a_{sc} \hat{\omega}^{sc}_a + \hat{\omega}^a_{st} \hat{\omega}^{st}_a - \hat{\omega}^r_{bc} \hat{\omega}^b_{r^c} - \hat{\omega}^r_{tb} \hat{\omega}^{bt}_r - \hat{\omega}^r_{sc} \hat{\omega}^{cs}_r \right. \\ \left. + \hat{\omega}^r_{st} \hat{\omega}^{st}_r + \hat{\omega}^a_{ac} \hat{\omega}^b_{b^c} + \hat{\omega}^a_{ta} \hat{\omega}^{bt}_b + \hat{\omega}^a_{ac} \hat{\omega}^s_{s^c} - \hat{\omega}^a_{ta} \hat{\omega}^s_{s^t} + \hat{\omega}^r_{rc} \hat{\omega}^b_{b^c} - \hat{\omega}^r_{rt} \hat{\omega}^{bt}_b + \hat{\omega}^r_{rc} \hat{\omega}^s_{s^c} + \hat{\omega}^r_{rt} \hat{\omega}^s_{s^t} \right] \quad (19)$$

It become quite obvious that we must only calculate the result for 6 ω being

$$\hat{\omega}_{pmn}, \hat{\omega}_{pma}, \hat{\omega}_{pab}, \\ \hat{\omega}_{cmn}, \hat{\omega}_{cma}, \hat{\omega}_{cab},$$

One may also see that to calculate them we only need the results of 8 Ω being this time

$$\hat{\Omega}_{pmn}, \\ \hat{\Omega}_{pma}, \hat{\Omega}_{pam}, \hat{\Omega}_{apm}, \\ \hat{\Omega}_{mab}, \hat{\Omega}_{amb}, \hat{\Omega}_{abm}, \\ \hat{\Omega}_{abc},$$

and we can esaly create a course of action from (10), (11) and (12) which is compute $\hat{V}_{\hat{r}}^{\hat{\mu}}$ and $\hat{V}_{\hat{\mu}\hat{r}}$ then the Ω then the ω . for the inverse vielbein the computation are quite easy as we have

$$\hat{V}_{\hat{\mu}}^{\hat{s}} \hat{V}_{\hat{r}}^{\hat{\mu}} = \delta_{\hat{r}}^{\hat{s}}$$

and so we have

$$\hat{V}_{\hat{\mu}}^{\hat{r}} = \begin{pmatrix} \delta^{-\gamma} V_r^\mu & -2\kappa \delta^{-\gamma} A_r^\alpha \\ 0 & \Phi_a^\alpha \end{pmatrix} \quad (20)$$

the same go for $\hat{V}_{\hat{\mu}\hat{r}}$ as we have the equation

$$\hat{V}_{\hat{r}}^{\hat{\mu}} \hat{V}_{\hat{\mu}\hat{s}} = \eta_{\hat{r}\hat{s}}$$

whcih gave us

$$\hat{V}_{\hat{\mu}\hat{s}} = \begin{pmatrix} \delta^\gamma V_{\mu s} & 2\kappa A_\mu^\alpha \Phi_{\alpha a} \\ 0 & Phi_{\alpha a} \end{pmatrix} \quad (21)$$

Now for the first part the Ω for $\hat{\Omega}_{pmn}$

$$\begin{aligned}
\hat{\Omega}_{pmn} &= \frac{1}{2}(\hat{V}_p^{\hat{\mu}}\hat{V}_m^{\nu} - \hat{V}_m^{\hat{\mu}}\hat{V}_p^{\nu})\partial_{\nu}\hat{V}_{\hat{m}n} \\
&= \frac{1}{2}(\hat{V}_p^{\mu}\hat{V}_m^{\nu} - \hat{V}_m^{\mu}\hat{V}_p^{\nu})\partial_{\nu}\hat{V}_{\mu n} \\
&= \frac{1}{2}(\hat{V}_p^{\mu}\hat{V}_m^{\nu} - \hat{V}_m^{\mu}\hat{V}_p^{\nu})\partial_{\nu}\hat{V}_{\mu n} \\
&= \frac{\delta^{-2\gamma}}{2}(V_p^{\mu}V_m^{\nu} - V_m^{\mu}V_p^{\nu})\partial_{\nu}(\delta^{\gamma}V_{\mu n}) \\
&= \frac{\delta^{-2\gamma}}{2}(V_p^{\mu}V_m^{\nu} - V_m^{\mu}V_p^{\nu})(\partial_{\nu}(\delta^{\gamma})V_{\mu n} + \delta^{\gamma}\partial_{\nu}(V_{\mu n})) \\
&= \frac{\delta^{-2\gamma}}{2}(V_p^{\mu}V_m^{\nu} - V_m^{\mu}V_p^{\nu})(\gamma\partial_{\nu}(\delta)\delta^{\gamma-1}V_{\mu n} + \delta^{\gamma}\partial_{\nu}(V_{\mu n})) \\
&= \frac{\delta^{-\gamma}}{2}(V_p^{\mu}V_m^{\nu} - V_m^{\mu}V_p^{\nu})(\gamma\partial_{\nu}(\delta)\delta^{-1}V_{\mu n} + \partial_{\nu}(V_{\mu n})) \\
&= \frac{\delta^{-\gamma}}{2}(V_p^{\mu}V_m^{\nu} - V_m^{\mu}V_p^{\nu})(\gamma\partial_{\nu}(\log \delta)V_{\mu n} + \partial_{\nu}(V_{\mu n}))
\end{aligned}$$

for $\hat{\Omega}_{pma}$

$$\begin{aligned}
\hat{\Omega}_{pma} &= \frac{1}{2}(\hat{V}_p^{\hat{\mu}}\hat{V}_m^{\nu} - \hat{V}_m^{\hat{\mu}}\hat{V}_p^{\nu})\partial_{\nu}\hat{V}_{\hat{m}a} \\
&= -\kappa\delta^{-2\gamma}(V_p^{\mu}V_m^{\nu} - V_m^{\mu}V_p^{\nu})\partial_{\nu}(A_{\mu}^{\alpha}\Phi_{a\alpha}) + \kappa\delta^{-\gamma}(A_p^{\alpha}V_m^{\nu} - \hat{A}_m^{\alpha}V_p^{\nu})\partial_{\nu}(\Phi_{a\alpha}) \\
&= -\kappa\delta^{-2\gamma}(V_p^{\mu}V_m^{\nu} - V_m^{\mu}V_p^{\nu})(\partial_{\nu}(A_{\mu}^{\alpha})\Phi_{a\alpha} + \partial_{\nu}(\Phi_{a\alpha})A_{\mu}^{\alpha}) + \kappa\delta^{-\gamma}(A_p^{\alpha}V_m^{\nu} - \hat{A}_m^{\alpha}V_p^{\nu})\partial_{\nu}(\Phi_{a\alpha}) \\
&= -\kappa\delta^{-2\gamma}(V_p^{\mu}V_m^{\nu} - V_m^{\mu}V_p^{\nu})(\partial_{\nu}(A_{\mu}^{\alpha})\Phi_{a\alpha} + \partial_{\nu}(\Phi_{a\alpha})A_{\mu}^{\alpha}) + (A_p^{\alpha}V_m^{\nu} - \hat{A}_m^{\alpha}V_p^{\nu})\partial_{\nu}(\Phi_{a\alpha}) \\
&= -\kappa\delta^{-2\gamma}(V_p^{\mu}V_m^{\nu} - V_m^{\mu}V_p^{\nu})\partial_{\nu}(A_{\mu}^{\alpha})\Phi_{a\alpha} \\
&= \kappa\delta^{-2\gamma}(F^{\alpha}_{pm}\Phi_{a\alpha} - \partial_{\nu}(V_p^{\mu}V_m^{\nu})A_{\mu}^{\alpha}\Phi_{a\alpha} + \partial_{\nu}(V_p^{\nu}V_m^{\mu})A_{\mu}^{\alpha}\Phi_{a\alpha}) \\
&= \kappa\delta^{-2\gamma}F^{\alpha}_{pm}\Phi_{a\alpha}
\end{aligned}$$

for $\hat{\Omega}_{pam}$

$$\begin{aligned}
\hat{\Omega}_{pam} &= \frac{1}{2}(\hat{V}_p^{\hat{\mu}}\hat{V}_a^{\nu} - \hat{V}_a^{\hat{\mu}}\hat{V}_p^{\nu})\partial_{\nu}\hat{V}_{\hat{m}m} \\
&= \frac{1}{2}(\hat{V}_p^{\mu}\hat{V}_a^{\nu} - \hat{V}_a^{\mu}\hat{V}_p^{\nu})\partial_{\nu}\hat{V}_{\mu m} \\
&= 0
\end{aligned}$$

same result for $\hat{\Omega}_{pam}$

for $\hat{\Omega}_{pab}$

$$\begin{aligned}
\hat{\Omega}_{pab} &= \frac{1}{2}(\hat{V}_p^{\hat{\mu}}\hat{V}_a^{\nu} - \hat{V}_a^{\hat{\mu}}\hat{V}_p^{\nu})\partial_{\nu}\hat{V}_{\hat{m}b} \\
&= \frac{-\delta^{-\gamma}}{2}\hat{V}_a^{\hat{\mu}}V_p^{\nu}\partial_{\nu}\hat{V}_{\hat{m}b} \\
&= \frac{\delta^{-\gamma}}{2}\Phi_a^{\alpha}V_p^{\nu}\partial_{\nu}\Phi_{\alpha b} \\
&= \frac{\delta^{-\gamma}}{2}\Phi_a^{\alpha}V_p^{\nu}\partial_{\nu}(h_{\alpha\beta}\Phi_b^{\beta}) \\
&= \frac{\delta^{-\gamma}}{2}\Phi_a^{\alpha}\Phi_b^{\beta}V_p^{\nu}\partial_{\nu}h_{\alpha\beta} + \frac{\delta^{-\gamma}}{2}\Phi_a^{\alpha}V_p^{\nu}h_{\alpha\beta}\partial_{\nu}\Phi_b^{\beta} \\
&= \frac{\delta^{-\gamma}}{2}\Phi_a^{\alpha}\Phi_b^{\beta}V_p^{\nu}\partial_{\nu}h_{\alpha\beta} + \frac{\delta^{-\gamma}}{2}\Phi_a^{\alpha}V_p^{\nu}\Phi_{\alpha}^c\delta_{cd}\Phi_{\beta}^d\partial_{\nu}\Phi_b^{\beta} \\
&= \frac{\delta^{-\gamma}}{2}\Phi_a^{\alpha}\Phi_b^{\beta}V_p^{\nu}\partial_{\nu}h_{\alpha\beta} + \frac{\delta^{-\gamma}}{2}\Phi_a^{\beta}V_p^{\nu}\partial_{\nu}\Phi_b^{\beta}
\end{aligned}$$

For $\hat{\Omega}_{apb}$

$$\begin{aligned}\hat{\Omega}_{apb} &= \frac{1}{2}(\hat{V}_a^{\hat{\mu}}\hat{V}_p^{\nu} - \hat{V}_p^{\hat{\mu}}\hat{V}_a^{\nu})\partial_{\nu}\hat{V}_{\hat{\mu}b} \\ &= \frac{1}{2}\hat{V}_a^{\hat{\mu}}\hat{V}_p^{\nu}\partial_{\nu}\hat{V}_{\hat{\mu}b} \\ &= \frac{1}{2}\hat{V}_a^{\alpha}\hat{V}_p^{\nu}\partial_{\nu}\hat{V}_{\alpha}^b\end{aligned}$$

For $\hat{\Omega}_{abp}$

$$\begin{aligned}\hat{\Omega}_{abp} &= \frac{1}{2}(\hat{V}_a^{\hat{\mu}}\hat{V}_b^{\nu} - \hat{V}_b^{\hat{\mu}}\hat{V}_a^{\nu})\partial_{\nu}\hat{V}_{\hat{\mu}p} \\ &= 0\end{aligned}$$

pareil pour Ω_{abc} then using (10) we can get the ω fairly easily

$$\hat{\omega}_{pmn} = \delta^{-\gamma}[\omega_{pmn} + \gamma(\eta_{pm}V_n^{\lambda} - \eta_{pn}V_m^{\lambda})\partial_{\lambda}], \quad (22a)$$

$$\hat{\omega}_{pma} = \kappa\delta^{-2\gamma}F_{pm}^{\alpha}\Phi_{\alpha a}, \quad (22b)$$

$$\hat{\omega}_{pab} = \frac{1}{2}\delta^{-\gamma}\Phi_a^{\alpha}V_p^{\mu}\partial_{\mu}\Phi_{\alpha b} - (a \leftrightarrow b), \quad (22c)$$

$$\hat{\omega}_{cmn} = -\kappa\delta^{-2\gamma}F_{mn}^{\alpha}\Phi_{\alpha c}, \quad (22d)$$

$$\hat{\omega}_{cab} = 0 \quad (22e)$$

where $F_{\mu\nu}^{\alpha} = \partial_{\mu}A_{\nu}^{\alpha} - \partial_{\nu}A_{\mu}^{\alpha}$ We can easily see the gauge invariance of theses expression.

4.5 finalisation

5 reduction of the new fields

5.1 scalar field reduction

5.2 Two form field reduction

6 Observation of new symmetry

7 Conclusion

References

- [1] Joel Scherk and John H Schwarz. “How to get masses from extra dimensions”. In: *Nuclear Physics B* 153 (1979), pp. 61–88.
- [2] Albert Einstein. “Zur allgemeinen relativitätstheorie”. In: *Sitzungsber. preuss. Akad. Wiss* 44.2 (1915), pp. 778–786.
- [3] Th Kaluza. “Zum unitätsproblem der physik”. In: *Sitzungsber. Preuss. Akad. Wiss. Berlin (Math. Phys.)* 1921.arXiv: 1803.08616 (1921), pp. 966–972.
- [4] Daniel Z Freedman and Antoine Van Proeyen. *Supergravity*. Cambridge university press, 2012.
- [5] Jnanadeva Maharana and John H Schwarz. “Noncompact symmetries in string theory”. In: *Nuclear Physics B* 390.1 (1993), pp. 3–32.

A lagrangian

By using the principle of least action and the variational principle we can see that

$$\begin{aligned}
\delta S &= \delta \int \mathcal{L} dx^\mu \\
&= \delta \int \frac{\partial \mathcal{L}}{\partial \eta_\rho} \delta \eta_\rho + \frac{\partial \mathcal{L}}{\partial \partial_\nu \eta_\rho} \delta \partial_\nu \eta_\rho dx^\mu \\
&= \delta \int \frac{\partial \mathcal{L}}{\partial \eta_\rho} \delta \eta_\rho - \partial_\nu \frac{\partial \mathcal{L}}{\partial \partial_\nu \eta_\rho} \delta \eta_\rho + \underbrace{\partial_\nu \left(\frac{\partial \mathcal{L}}{\partial \partial_\nu \eta_\rho} \delta \eta_\rho \right)}_{\text{border terme equal to zero}} dx^\mu \\
&= 0
\end{aligned}$$

which gave the equation

$$\boxed{\frac{\partial \mathcal{L}}{\partial \eta_\rho} - \partial_\nu \frac{\partial \mathcal{L}}{\partial \partial_\nu \eta_\rho} = 0} \quad (23)$$

which is in fact the equation of mouvement for the field