


Dimensional Reduction in Supergravity and Duality Symmetries

Adam Franchot–Fowler

Internship carried out from 08/06/2026 to 24/07/2026



Internship tutor: M. Camille Eloy

Referent teacher: M. Stéphane Douin

Educational institution: Université Paris-Saclay / Magistère de physique fondamentale d'Orsay

Internship host institution: Laboratoire de Physique (UMR CNRS 5672), ENS de Lyon

Acknowledgment

Notations and Abbreviations

For quick reference, we summarize here the index conventions used throughout this report, following [1]; their motivation is explained in Section 2.

- Hatted Greek letters denote curved indices $(\hat{\mu}, \hat{\nu}, \hat{\rho}, \dots)$.
- Hatted Latin letters denote flat indices $(\hat{p}, \hat{m}, \hat{n}, \dots)$.
- For space-time indices taking D values, late Greek letters are used in the curved case $(\mu, \nu, \rho, \dots)$ and Latin letters in the flat case (p, m, n) .
- For internal indices taking E values, early Greek letters are used in the curved case $(\alpha, \beta, \gamma, \dots)$ and Latin letters in the flat case (a, b, c) .

Contents

Notation and abbreviations	1
1 Introduction	3
2 Vielbein Formalism	3
3 Lagrangian Field Theory	4
3.1 Lagrangian Density	4
3.2 General Example for Scalar and Vector Fields, and Result for Two-Forms	5
3.3 NS-NS Action of Supergravity	6
3.4 Different Parts of the Action	6
4 Gravitational Action Reduction	6
4.1 Rewriting the Action	6
4.2 Vielbein Ansatz	7
4.3 Gauge Observation	7
4.4 Ricci Scalar Reduction	8
4.5 Adding the Dilaton Field Reduction	9
5 Two-Form Field Reduction	10
5.1 The Reduction	10
5.2 Observation of Gauge	11
6 Complete Reduced Action	11
7 Observation of a New Symmetry	11
7.1 The $O(E,E)$ Group	11
7.2 Rewriting the Reduced Action	12
7.3 Action Invariance under $O(E,E)$	12
8 Conclusion	13
A General Relativity Background	14
B Euler–Lagrange Equation	14
C Summary of the Reduced Field Content	15

1 Introduction

Since the 19th-century unification of electricity and magnetism, and the later development of classical and quantum field theory, physicists have been looking for a single mathematical framework that could describe all fundamental interactions at once. A great deal of progress has been made for the electromagnetic, weak, and strong forces, which are now successfully unified in the Standard Model. Gravity, however, remains the main stumbling block: indeed, its description by General Relativity [2] sits completely apart from the quantum field-theoretic language we use for every other force. One way to address this issue is to add a new symmetry to nature, supersymmetry. This symmetry is mainly used in two theories: the first is supergravity, which, as the name suggests, attempts to combine the principle of supersymmetry with general relativity. The other major theory is superstring theory, which attempts to do the same thing within the framework of string theory. An advantage of the former over the latter is that supergravity does not require extra dimensions, whereas superstring theory does. This is precisely why we work within the framework of supergravity here.

The tool we use to make the role of these extra dimensions manifest is dimensional reduction, in the spirit of the original Kaluza–Klein idea: a theory living in $D + E$ dimensions can be reinterpreted, by a D -dimensional observer, as a theory in D dimensions containing extra scalar and vector fields that encode the geometry of the E hidden dimensions. Carrying this out consistently and identifying what those extra fields are is the main technical goal of this internship, and it is what ultimately allows us to uncover a hidden symmetry relating theories that look different in $D + E$ dimensions but turn out to be equivalent once reduced.

Concretely, we begin by describing what a field theory is, in order to work out how to write down the actions of the fields we will use. We then follow the steps of [1] by performing a dimensional reduction from $D + E$ dimensions down to 4 dimensions. However, following [3], we do not simply use the classical Einstein action, but the NS-NS action of supergravity, which adds two new fields. We do this using the vielbein formalism, as it is useful for the description of fermions, even though here we only describe bosonic fields. We then show that, although in high dimensions only the $SL(E, E)$ symmetry is detectable without much work, an additional $O(E, E)$ symmetry appears once we go to low dimensions; this means that a number of theories that look distinct in $D + E$ dimensions are in fact equivalent.

Throughout, we work in the simplest setting: we assume that none of the fields depend on the internal coordinates, so that for any field we have $\partial_\alpha A^{\hat{\mu}} = 0$.

2 Vielbein Formalism

Throughout this internship we will use the vielbein formalism, following the same notation as in [1] (summarized on page 1): hatted Greek letters ($\hat{\mu}, \hat{\nu}, \hat{\rho}, \dots$) denote curved indices and hatted Latin letters ($\hat{p}, \hat{m}, \hat{n}, \dots$) denote flat indices; among these, the D space-time directions are labelled by late Greek letters ($\mu, \nu, \rho, \dots$) in the curved case and late Latin letters (p, m, n) in the flat case, while the E internal directions are labelled by early Greek letters ($\alpha, \beta, \gamma, \dots$) in the curved case and early Latin letters (a, b, c) in the flat case. The general coordinate $x^{\hat{\mu}}$ is therefore made up of a space-time coordinate x^μ and an internal coordinate y^α . The flat Lorentz indices are precisely the ones on which the vielbein formalism, introduced below, lets us use the flat Minkowski metric η_{ab} to raise and lower indices, in place of the curved metric $g_{\mu\nu}$.

In general relativity we often encounter metrics that are difficult to handle, such as the Kerr–Newman metric. A useful idea is then to work in an orthonormal basis $\{e_\mu^a\}$, which satisfies

$$g_{\mu\nu} V_a^\mu V_b^\nu = \eta_{ab}, \quad (1)$$

where g is the original metric and η is the Minkowski metric.

We call this basis the vielbein. In this basis, indices are raised and lowered using the Minkowski metric. The dual basis, also called the inverse vielbein, is defined by

$$V_\nu^a V_b^\nu = \delta_b^a. \quad (2)$$

Consequently, the curved metric can be reconstructed from the vielbein:

$$g_{\mu\nu} = \eta_{ab} V_\mu^a V_\nu^b. \quad (3)$$

The vielbein also allows us to transform vectors, covectors, and tensors between curved and flat indices:

$$T^a = V_\mu^a T^\mu, \quad (4)$$

$$T_a = V_a^\mu T_\mu, \quad (5)$$

$$T_b^a = V_\mu^a V_b^\nu T_\nu^\mu. \quad (6)$$

The introduction of flat indices requires a corresponding modification of the usual transformation law. In particular, the Christoffel connection must be replaced by the spin connection, defined by

$$\nabla_\mu X_b^a = \partial_\mu X_b^a + \omega_\mu^a{}_c X_b^c - \omega_\mu^c{}_b X_c^a. \quad (7)$$

The spin connection remains related to the Levi–Civita connection through

$$\Gamma_{\mu\lambda}^\nu = V_a^\nu \partial_\mu V_\lambda^a + V_a^\nu V_\lambda^b \omega_\mu^a{}_b, \quad (8)$$

$$\omega_\mu^a{}_b = V_\nu^a V_b^\lambda \Gamma_{\mu\lambda}^\nu - V_b^\lambda \partial_\mu V_\lambda^a. \quad (9)$$

With these relations we can prove the antisymmetry of the last two indices of the spin connection:

$$\begin{aligned}\nabla_\mu \eta_{ab} &= \partial_\mu X_{ab} - \omega_\mu{}^a{}_c \eta_{cb} - \omega_\mu{}^c{}_b \eta_{ac} \\ &= -\omega_{\mu ab} - \omega_{\mu ba} \\ &= 0.\end{aligned}\tag{10}$$

So, in effect, we have

$$-\omega_{\mu ab} = \omega_{\mu ba}.\tag{11}$$

3 Lagrangian Field Theory

3.1 Lagrangian Density

In this internship we are going to use Lagrangian field theory to describe the objects we will be working with. The main interest of this framework is that it lets us work with an infinite number of degrees of freedom. To better understand what we are going to do, let us start from a simple case in 1D, which we will then generalize to ND . The "simple" example is going to be an infinitely long elastic rod undergoing small longitudinal vibrations. We can model this with a system of discrete points of equal mass, connected by springs of natural length a , as shown in Figure 1. Point i is displaced from its equilibrium position by a quantity η_i .




Figure 1: A discrete chain of equal masses connected by springs of natural length a . Top: the chain at equilibrium. Bottom: the chain displaced from equilibrium, with η_i the displacement of mass i .

We can then obtain the potential and kinetic energy as

$$V = \frac{1}{2} \sum_i m \dot{\eta}_i^2,\tag{12a}$$

$$T = \frac{1}{2} \sum_i k(\eta_{i+1} - \eta_i)^2,\tag{12b}$$

and now we can write the Lagrangian of the system, which is

$$L = T - V = \frac{1}{2} \sum_i m \dot{\eta}_i^2 - k(\eta_{i+1} - \eta_i)^2,$$

which can be rewritten as

$$L = T - V = \frac{1}{2} \sum_i a \left[\frac{m}{a} \dot{\eta}_i^2 - ak \left(\frac{\eta_{i+1} - \eta_i}{a} \right)^2 \right] = \sum_i a L_i.\tag{13}$$

From this we obtain

$$\frac{m}{a} \ddot{\eta}_i^2 - ka \left(\frac{\eta_{i+1} - \eta_i}{a} \right)^2 + ka \left(\frac{\eta_i - \eta_{i-1}}{a} \right)^2 = 0$$

as the equation of motion for each η_i . If we pass to the limit $a \rightarrow 0$, we can see that $\frac{m}{a} \rightarrow \mu$, which is the mass per unit length. The limit of ka is far less evident, and to determine it we have to use Hooke's law, since we have an elastic rod, for which the extension of the rod per unit length is proportional to the applied force. Thus the relation is

$$F = Y\xi,\tag{14}$$

where ξ is the elongation per unit length and Y is Young's modulus. Now we can write $\xi = \frac{\eta_{i+1} - \eta_i}{a}$. The force necessary to stretch the spring by this amount is

$$F = k(\eta_{i+1} - \eta_i) = ka \left(\frac{\eta_{i+1} - \eta_i}{a} \right) = ka\xi,$$

which means that ka is the Young's modulus of the continuous rod. As we go from the discrete to the continuous case, the integer i identifying the particular mass point becomes the continuous position coordinate x . We now have $\eta(x)$ instead of η_i . Furthermore, as a goes to zero, we have

$$\frac{\eta_{i+1} - \eta_i}{a} = \frac{\eta(x+a) - \eta(x)}{a} \rightarrow \frac{d\eta}{dx},$$

since a was playing the role of dx . As it approaches zero, the sum we had earlier turns into an integral over the length of the rod, and the Lagrangian becomes

$$L = \frac{1}{2} \int \left[\mu \dot{\eta}^2 - Y \left(\frac{d\eta}{dx} \right)^2 \right] dx. \quad (15)$$

In the limit $a \rightarrow 0$ the last two terms of the equation of motion become

$$\lim_{a \rightarrow 0} -\frac{Y}{a} \left[\frac{d\eta}{dx} \Big|_x - \frac{d\eta}{dx} \Big|_{x-a} \right],$$

which is, in effect, a second derivative of η . Hence, the equation of motion for the continuous case is

$$\mu \frac{d^2 \eta}{dt^2} - Y \frac{d^2 \eta}{dx^2} = 0, \quad (16)$$

which is the wave equation in one dimension, with propagation velocity

$$v = \sqrt{\frac{Y}{\mu}}.$$

And so we recover d'Alembert's equation.

Now we can see that, even though we have an infinite number of points, there is not an infinite number of Lagrange equations, but merely one. This also means that we can write the Lagrangian as

$$L = \int \mathcal{L} dx,$$

where here $\mathcal{L}$ is the Lagrangian per unit length. This generalizes to ND , where the Lagrangian can be written as

$$L = \iiint \mathcal{L}(\eta_\rho, \frac{\partial \eta_\rho}{\partial x^\nu}) dx^N. \quad (17)$$

Now, using Einstein's convention, we can rewrite the action as

$$S = \int \mathcal{L} dx^\mu. \quad (18)$$

As a final note, one can see that both the variational method and the Lagrange equation work in the same way as before the transformation.

3.2 General Example for Scalar and Vector Fields, and Result for Two-Forms

In field theory, the Lagrangian has a classical form for any field (often the simplest case). For this work, only the scalar, vector, and two-form Lagrangians are needed, along with the metric, which is handled by Einstein's equations.

So, for a scalar field we have

$$\boxed{\mathcal{L} = -\frac{1}{2} \partial_\mu \Phi \partial^\mu \Phi}, \quad (19)$$

which gives us $\square \Phi = 0$ as the equation of motion.

And for a vector field we have

$$\boxed{\mathcal{L} = -\frac{1}{4} F_{\mu\nu} F^{\mu\nu}}, \quad (20)$$

where, for the field A ,

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu.$$

The equation of motion is $\partial_\mu F^{\mu\nu} = 0$, which is exactly Maxwell's equations. In vacuum, the Einstein equation gives us

$$\boxed{\mathcal{L} = R},$$

where R is the Ricci scalar, and the equation of motion is exactly Einstein's equation.

Finally, for the Lagrangian of a two-form field we look to [3], where it is given by

$$\boxed{\mathcal{L} = \frac{1}{12} g^{\hat{\mu}\mu'} g^{\nu\nu'} g^{\rho\rho'} H_{\mu\nu\rho} H_{\mu'\nu'\rho'}}, \quad (21)$$

where, for the two-form $B_{\nu\rho}$, we have

$$H_{\mu\nu\rho} = \partial_\mu B_{\nu\rho} + \text{cyclic permutations},$$

and the equation of motion is $\partial_\mu H^{\mu\nu\rho} = 0$.

3.3 NS-NS Action of Supergravity

With these objects we can construct the NS-NS action, which has the form

$$S = \int dx^D \int \frac{dy^E}{\rho(E)} \sqrt{-\hat{g}} e^{\hat{\phi}} \left(\hat{R} + \hat{g}^{\hat{\mu}\hat{\nu}} \partial_{\hat{\mu}} \hat{\phi} \partial^{\hat{\nu}} \hat{\phi} - \frac{1}{12} g^{\hat{\mu}\hat{\mu}'} g^{\nu\nu'} g^{\rho\rho'} H_{\mu\nu\rho} H_{\mu'\nu'\rho'} \right), \quad (22)$$

where $\rho(E)$ is a normalization factor, corresponding in practice to the invariant volume of the internal space, and where $\hat{R}$ is similar to the classical Hilbert–Einstein action but in $D+E$ dimensions, $\hat{\phi}$ is a dilaton field, and $-\frac{1}{12} H_{\mu\nu\rho} H^{\mu\nu\rho}$ is the action of a two-form. Finally, since no field depends on the internal coordinates, and given $\rho(E)$, we can simplify this integral. We will now separate this action into three parts, one for each field. Now that we have this action in $D+E$ dimensions, we need, as in [1], to reduce it, which basically means looking at how a D -dimensional observer would perceive those extra dimensions.

3.4 Different Parts of the Action

We can separate the action into two main components: one being the gravitational part with the dilaton, and the other being the two-form part. So we have

$$S_{\hat{g}} = \int dx^D \sqrt{-\hat{g}} e^{-\hat{\phi}} [\hat{R} + \hat{g}^{\hat{\mu}\hat{\nu}} \partial_{\hat{\mu}} \hat{\phi} \partial^{\hat{\nu}} \hat{\phi}], \quad (23a)$$

$$S_{\hat{B}} = \int dx^D \sqrt{-\hat{g}} e^{-\hat{\phi}} \left(-\frac{1}{12} H_{\mu\nu\rho} H^{\mu\nu\rho} \right). \quad (23b)$$

4 Gravitational Action Reduction

We will begin with the gravitational part of the NS-NS action.

4.1 Rewriting the Action

For this part we do not need the $\hat{g}^{\hat{\mu}\hat{\nu}} \partial_{\hat{\mu}} \hat{\phi} \partial^{\hat{\nu}} \hat{\phi}$ part of the action, so we will ignore it entirely. Now, using the fact that $\hat{R}$ can be written as

$$\hat{R} = \hat{V}^{\hat{\mu}\hat{\nu}} \hat{V}^{\hat{\rho}\hat{s}} \hat{R}_{\hat{\mu}\hat{\nu}\hat{\rho}\hat{s}},$$

where

$$\hat{R}_{\hat{\mu}\hat{\nu}\hat{\rho}\hat{s}} = \partial_{\hat{\mu}} \hat{\omega}_{\hat{\nu}\hat{\rho}\hat{s}} + \hat{\omega}_{\hat{\mu}\hat{\rho}}^{\hat{t}} \hat{\omega}_{\hat{\nu}\hat{t}\hat{s}} - (\mu \leftrightarrow \nu),$$

and the ω 's are defined by

$$\hat{\omega}_{\hat{t}\hat{r}\hat{s}} = -\hat{\Omega}_{\hat{t}\hat{r}\hat{s}} + \hat{\Omega}_{\hat{r}\hat{s}\hat{t}} - \hat{\Omega}_{\hat{s}\hat{t}\hat{r}},$$

and finally the Ω 's are defined via the vielbein by

$$\hat{\Omega}_{\hat{r}\hat{s}\hat{t}} = \frac{1}{2} (\hat{V}_{\hat{r}}^{\hat{\mu}} \hat{V}_{\hat{s}}^{\hat{\nu}} - \hat{V}_{\hat{s}}^{\hat{\mu}} \hat{V}_{\hat{r}}^{\hat{\nu}}) \partial_{\hat{\nu}} \hat{V}_{\hat{\mu}\hat{t}},$$

with all of that we can now rewrite the action $S_{\hat{g}}$ in terms of $\hat{\omega}$:

$$\begin{aligned} S_{\hat{g}} &= \int dx^D \hat{V} e^{-\hat{\phi}} \hat{R} \\ &= \int dx^D \hat{V} e^{-\hat{\phi}} \hat{V}^{\hat{\mu}\hat{\nu}} \hat{V}^{\hat{\rho}\hat{s}} \hat{R}_{\hat{\mu}\hat{\nu}\hat{\rho}\hat{s}} \\ &= \int dx^D \hat{V} e^{-\hat{\phi}} \hat{V}^{\hat{\mu}\hat{\nu}} \hat{V}^{\hat{\rho}\hat{s}} \left([\partial_{\hat{\mu}} \hat{\omega}_{\hat{\nu}\hat{\rho}\hat{s}} - \partial_{\hat{\nu}} \hat{\omega}_{\hat{\mu}\hat{\rho}\hat{s}}] + [\hat{\omega}_{\hat{\mu}\hat{\rho}}^{\hat{t}} \hat{\omega}_{\hat{\nu}\hat{t}\hat{s}} - \hat{\omega}_{\hat{\nu}\hat{\rho}}^{\hat{t}} \hat{\omega}_{\hat{\mu}\hat{t}\hat{s}}] \right) \\ &\text{Integrating the first two terms by parts and dropping the resulting boundary term,} \\ &= \int dx^D \{ \hat{V} e^{-\hat{\phi}} [\hat{\omega}_{\hat{r}}^{\hat{t}} \hat{\omega}_{\hat{s}}^{\hat{t}} \hat{\omega}_{\hat{t}\hat{s}}^{\hat{r}} - \hat{\omega}_{\hat{s}}^{\hat{t}} \hat{\omega}_{\hat{r}}^{\hat{t}} \hat{\omega}_{\hat{t}\hat{s}}^{\hat{r}}] - 2 \hat{\omega}_{\hat{\nu}\hat{\rho}\hat{s}} [\hat{V} e^{-\hat{\phi}} \hat{V}^{\hat{\mu}\hat{\nu}} \partial_{\hat{\mu}} \hat{V}^{\hat{\rho}\hat{s}} + \hat{V} e^{-\hat{\phi}} \hat{V}^{\hat{\nu}\hat{s}} \partial_{\hat{\mu}} \hat{V}^{\hat{\mu}\hat{\rho}} + e^{-\hat{\phi}} \hat{V}^{\hat{\mu}\hat{\nu}} \hat{V}^{\hat{\rho}\hat{s}} \partial_{\hat{\mu}} \hat{V} + \hat{V} \hat{V}^{\hat{\mu}\hat{\nu}} \hat{V}^{\hat{\rho}\hat{s}} \partial_{\hat{\mu}} e^{-\hat{\phi}}] \} \\ &\text{Using the raising/lowering rules and the equation above to simplify,} \\ &= \int dx^D \hat{V} e^{-\hat{\phi}} \{ \hat{\omega}_{\hat{s}}^{\hat{t}} \hat{\omega}_{\hat{r}}^{\hat{t}} \hat{\omega}_{\hat{t}\hat{s}}^{\hat{r}} + \hat{\omega}_{\hat{r}}^{\hat{t}} \hat{\omega}_{\hat{s}}^{\hat{t}} \hat{\omega}_{\hat{t}\hat{s}}^{\hat{r}} + 2 \hat{\omega}_{\hat{s}}^{\hat{r}} \hat{V}_{\hat{r}}^{\hat{\mu}} \partial_{\hat{\mu}} \hat{\phi} \}. \end{aligned} \quad (24)$$

4.2 Vielbein Ansatz

Now that we have explicitly written the action in terms of the Latin connection and the two other fields, what remains is to assume an ansatz for the vielbein. Why the vielbein and not the metric, one may ask? Because, as stated in Section 2, all the information contained in the metric is also contained in the vielbein, and since we primarily work with the vielbein in the action, it is easier to formulate the ansatz on it. We now use the same ansatz as in [1], except that we take the parameter γ used there to be zero from the start, and we use the same parameter conventions as in [3]. Finally, thanks to local Lorentz invariance we can choose a triangular parametrization, giving us

$$\hat{V}_{\hat{\mu}}^{\hat{r}} = \begin{pmatrix} V_{\mu}^r & A_{\mu}^{(1)\beta} \Phi_{\beta}^a \\ 0 & \Phi_{\alpha}^a \end{pmatrix}. \quad (25)$$

Now, still using (2), we can compute the inverse vielbein and then the metric and inverse metric, which gives us

$$\hat{V}_{\hat{\mu}}^{\hat{r}} = \begin{pmatrix} V_r^{\mu} & -A_r^{(1)\alpha} \\ 0 & \Phi_a^{\alpha} \end{pmatrix} \quad (26)$$

for the inverse vielbein, and

$$\hat{g}_{\hat{\mu}\hat{\nu}} = \begin{pmatrix} g_{\mu\nu} - A_{\mu}^{\gamma} A_{\nu\lambda}^{(1)} & -A_{\mu\beta}^{(1)} \\ A_{\nu\alpha}^{(1)} & -h_{\alpha\beta} \end{pmatrix}, \quad \hat{g}^{\hat{\mu}\hat{\nu}} = \begin{pmatrix} g^{\mu\nu} & -A^{(1)\mu\beta} \\ -A^{(1)\nu\alpha} & -h^{\alpha\beta} + A_{\rho}^{(1)\beta} A^{\rho\alpha} \end{pmatrix} \quad (27)$$

for the metric.

4.3 Gauge Observation

Now that we have set up the ansatz, we would like to identify what each object is. To that end, we can use the classical gauge transformation $x'^{\mu} = x^{\mu} + \xi^{\mu}(x)$ and use the fact that it can easily be rewritten as a Lie derivative. Indeed, since we already know how different objects transform under a Lie derivative, we can easily identify what each object is.

So, firstly, let us list every known transformation:

$$\delta\phi(x) = \xi^{\mu} \partial_{\mu} \phi(x), \quad (28)$$

$$\delta U^{\mu} = \xi^{\rho} \partial_{\rho} U^{\mu} - (\partial_{\rho} \xi^{\mu}) U^{\rho}, \quad (29)$$

$$\delta\omega_{\mu} = \xi^{\rho} \partial_{\rho} \omega_{\mu} + (\partial_{\rho} \xi^{\mu}) \omega_{\rho}, \quad (30)$$

$$\delta T_{\nu}^{\mu} = \xi^{\rho} \partial_{\rho} T_{\nu}^{\mu} - (\partial_{\rho} \xi^{\mu}) T_{\nu}^{\rho} + (\partial_{\nu} \xi^{\rho}) T_{\rho}^{\mu}. \quad (31)$$

We now apply the transformation to each block of the vielbein, in order to see whether it corresponds to some familiar type of field, still using (28). Let us start with the block $\hat{V}_{\mu}^r$:

$$\begin{aligned} \delta \hat{V}_{\mu}^r &= \delta V_{\mu}^r = \xi^{\rho} \partial_{\rho} \hat{V}_{\mu}^r + \partial_{\mu} \xi^{\rho} \hat{V}_{\rho}^r \\ &= \xi^{\rho} \partial_{\rho} \hat{V}_{\mu}^r + \partial_{\mu} \xi^{\nu} \hat{V}_{\nu}^r \\ &= \xi^{\rho} \partial_{\rho} V_{\mu}^r + \partial_{\mu} \xi^{\nu} V_{\nu}^r, \end{aligned}$$

which is exactly the transformation law of a vector field under a Lie derivative [4]. Applying the same method to the other blocks, we obtain, for Φ_{α}^a ,

$$\delta \Phi_{\alpha}^a = \xi^{\rho} \partial_{\rho} \Phi_{\alpha}^a,$$

which is exactly the transformation law of a scalar field. Finally, for $A_{\mu}^{(1)\alpha}$ we obtain

$$\delta A_{\mu}^{(1)\alpha} = \xi^{\rho} \partial_{\rho} A_{\mu}^{(1)\alpha} + \partial_{\mu} \xi^{\rho} A_{\rho}^{(1)\alpha} + \partial_{\mu} \xi^{\alpha},$$

which corresponds to the transformation law of a vector field, up to an extra correction term. One may further note that, restricting to ξ^{α} only, we obtain

$$\delta V_{\mu}^r = \delta \Phi_{\alpha}^a = 0, \quad (32a)$$

$$\delta A_{\mu}^{(1)\alpha} = \partial_{\mu} \xi^{\alpha}. \quad (32b)$$

So, what remains of the D -dimensional diffeomorphisms, once restricted to ξ^{α} independent of the internal coordinates y , corresponds to a set of E abelian gauge symmetries. The vielbein in (25) thus contains the D -dimensional vielbein (i.e., the graviton), E vector fields $A_{\mu}^{(1)\alpha}$, and E^2 scalar fields Φ_{α}^a . However, because of local Lorentz invariance in the internal directions, only $\frac{1}{2}E(E+1)$ of these scalars actually correspond to propagating degrees of freedom. We also introduce $\delta = \det \Phi$.

4.4 Ricci Scalar Reduction

Now that we are better acquainted with the tools we will use, we can begin the calculation. To start, we consider the Einstein action in $D + E$ dimensions, modified by the addition of the dilaton.

First, let us fully expand $S_{\hat{g}}$: this shows that we technically need to compute 8 ω 's; however, thanks to the antisymmetry of the last two indices, this reduces to 6. So the action is

$$S = -\frac{1}{4\kappa^2} \int d^D x \int \frac{d^E y}{\rho(E)} \hat{V} \left[\hat{\omega}^a_{bc} \hat{\omega}^{bc}_a - \hat{\omega}^a_{tb} \hat{\omega}^{bt}_a + \hat{\omega}^a_{sc} \hat{\omega}^{sc}_a + \hat{\omega}^a_{st} \hat{\omega}^{st}_a - \hat{\omega}^r_{bc} \hat{\omega}^b_{r^c} - \hat{\omega}^r_{tb} \hat{\omega}^{bt}_r - \hat{\omega}^r_{sc} \hat{\omega}^s_{r^c} \right. \\ \left. - \hat{\omega}^r_{st} \hat{\omega}^{st}_r + \hat{\omega}^a_{ac} \hat{\omega}^b_{b^c} + \hat{\omega}^a_{ta} \hat{\omega}^{bt}_b + \hat{\omega}^a_{ac} \hat{\omega}^s_{s^c} - \hat{\omega}^a_{ta} \hat{\omega}^s_{s^t} + \hat{\omega}^r_{rc} \hat{\omega}^b_{b^c} - \hat{\omega}^r_{rt} \hat{\omega}^{bt}_b + \hat{\omega}^r_{rc} \hat{\omega}^s_{s^c} + \hat{\omega}^r_{rt} \hat{\omega}^s_{s^t} + \right. \\ \left. 2\hat{\omega}_s{}^{rs} \hat{V}_r{}^\mu \partial_\mu \hat{\phi} + 2\hat{\omega}_b{}^{rb} \hat{V}_r{}^\mu \partial_\mu \hat{\phi} \right] \quad (33)$$

It then becomes clear that we only need to compute six components of ω , namely

$$\hat{\omega}_{pmn}, \hat{\omega}_{pma}, \hat{\omega}_{pab}, \\ \hat{\omega}_{cmn}, \hat{\omega}_{cma}, \hat{\omega}_{cab}.$$

One can also check that computing these six components only requires the following eight components of Ω :

$$\hat{\Omega}_{pmn}, \\ \hat{\Omega}_{pma}, \hat{\Omega}_{pam}, \hat{\Omega}_{apm}, \\ \hat{\Omega}_{mab}, \hat{\Omega}_{amb}, \hat{\Omega}_{abm}, \\ \hat{\Omega}_{abc}.$$

Using (26), (4.1) and (4.1), we can now set out a clear strategy: first the Ω 's, then the ω 's. Let us begin the computation of the Ω 's, starting with $\hat{\Omega}_{pmn}$:

$$\hat{\Omega}_{pmn} = \frac{1}{2} (\hat{V}_p{}^\mu \hat{V}_m{}^\nu - \hat{V}_m{}^\mu \hat{V}_p{}^\nu) \partial_\nu \hat{V}_{\mu n} \\ = \frac{1}{2} (V_p{}^\mu V_m{}^\nu - V_m{}^\mu V_p{}^\nu) \partial_\nu (V_{\mu n}).$$

For $\hat{\Omega}_{pma}$, we find:

$$\hat{\Omega}_{pma} = \frac{1}{2} (\hat{V}_p{}^\mu \hat{V}_m{}^\nu - \hat{V}_m{}^\mu \hat{V}_p{}^\nu) \partial_\nu \hat{V}_{\mu a} \\ = -(V_p{}^\mu V_m{}^\nu - V_m{}^\mu V_p{}^\nu) \partial_\nu (A_\mu^{(1)\alpha} \Phi_{a\alpha}) + (A_p^{(1)\alpha} V_m{}^\nu - A_m^{(1)\alpha} V_p{}^\nu) \partial_\nu (\Phi_{a\alpha}) \\ = -(V_p{}^\mu V_m{}^\nu - V_m{}^\mu V_p{}^\nu) (\partial_\nu (A_\mu^{(1)\alpha}) \Phi_{a\alpha} + \partial_\nu (\Phi_{a\alpha}) A_\mu^{(1)\alpha}) + (A_p^{(1)\alpha} V_m{}^\nu - A_m^{(1)\alpha} V_p{}^\nu) \partial_\nu (\Phi_{a\alpha}) \\ = -(V_p{}^\mu V_m{}^\nu - V_m{}^\mu V_p{}^\nu) \partial_\nu (A_\mu^{(1)\alpha}) \Phi_{a\alpha} \\ = F^\alpha{}_{pm} \Phi_{a\alpha} - \partial_\nu (V_p{}^\mu V_m{}^\nu) A_\mu^{(1)\alpha} \Phi_{a\alpha} + \partial_\nu (V_p{}^\nu V_m{}^\mu) A_\mu^{(1)\alpha} \Phi_{a\alpha} \\ = F^\alpha{}_{pm} \Phi_{a\alpha}.$$

For $\hat{\Omega}_{pam}$:

$$\hat{\Omega}_{pam} = \frac{1}{2} (\hat{V}_p{}^\mu \hat{V}_a{}^\nu - \hat{V}_a{}^\mu \hat{V}_p{}^\nu) \partial_\nu \hat{V}_{\mu m} \\ = 0.$$

The same result holds for $\hat{\Omega}_{apm}$.

For $\hat{\Omega}_{pab}$:

$$\begin{aligned}
\hat{\Omega}_{pab} &= \frac{1}{2}(\hat{V}_p^{\hat{\mu}}\hat{V}_a^{\nu} - \hat{V}_a^{\hat{\mu}}\hat{V}_p^{\nu})\partial_{\nu}\hat{V}_{\hat{\mu}b} \\
&= \frac{-1}{2}\hat{V}_a^{\hat{\mu}}V_p^{\nu}\partial_{\nu}\hat{V}_{\hat{\mu}b} \\
&= \frac{1}{2}\Phi_a^{\alpha}V_p^{\nu}\partial_{\nu}\Phi_{\alpha b} \\
&= \frac{1}{2}\Phi_a^{\alpha}V_p^{\nu}\partial_{\nu}(h_{\alpha\beta}\Phi_b^{\beta}) \\
&= \frac{1}{2}\Phi_a^{\alpha}\Phi_b^{\beta}V_p^{\nu}\partial_{\nu}h_{\alpha\beta} + \frac{1}{2}\Phi_a^{\alpha}V_p^{\nu}h_{\alpha\beta}\partial_{\nu}\Phi_b^{\beta} \\
&= \frac{1}{2}\Phi_a^{\alpha}\Phi_b^{\beta}V_p^{\nu}\partial_{\nu}h_{\alpha\beta} + \frac{1}{2}\Phi_{\beta a}V_p^{\nu}\partial_{\nu}\Phi_b^{\beta} \\
&= \frac{1}{2}\Phi_a^{\alpha}\Phi_b^{\beta}V_p^{\nu}\partial_{\nu}h_{\alpha\beta} + \frac{1}{2}V_p^{\nu}\partial_{\nu}(\underbrace{\Phi_{\beta a}\Phi_b^{\beta}}_{=\delta_{ab}, \text{ i.e. constant}}) - \Phi_b^{\beta}V_p^{\nu}\partial_{\nu}\Phi_{\beta a} \\
&= \frac{1}{2}\Phi_a^{\alpha}\Phi_b^{\beta}V_p^{\nu}\partial_{\nu}h_{\alpha\beta} - \Phi_b^{\beta}V_p^{\nu}\partial_{\nu}\Phi_{\beta a}.
\end{aligned}$$

For $\hat{\Omega}_{apb}$:

$$\begin{aligned}
\hat{\Omega}_{apb} &= \frac{1}{2}(\hat{V}_a^{\hat{\mu}}\hat{V}_p^{\nu} - \hat{V}_p^{\hat{\mu}}\hat{V}_a^{\nu})\partial_{\nu}\hat{V}_{\hat{\mu}b} \\
&= \frac{1}{2}\hat{V}_a^{\hat{\mu}}\hat{V}_p^{\nu}\partial_{\nu}\hat{V}_{\hat{\mu}b} \\
&= \frac{1}{2}\hat{V}_a^{\alpha}\hat{V}_p^{\nu}\partial_{\nu}\hat{V}_{\alpha b} \\
&= \frac{-1}{2}\Phi_a^{\alpha}V_p^{\nu}\partial_{\nu}\Phi_{\alpha b}.
\end{aligned}$$

For $\hat{\Omega}_{abp}$:

$$\begin{aligned}
\hat{\Omega}_{abp} &= \frac{1}{2}(\hat{V}_a^{\hat{\mu}}\hat{V}_b^{\nu} - \hat{V}_b^{\hat{\mu}}\hat{V}_a^{\nu})\partial_{\nu}\hat{V}_{\hat{\mu}p} \\
&= 0.
\end{aligned}$$

The same holds for Ω_{abc} . Using (25), we can then obtain the ω 's:

$$\hat{\omega}_{pmn} = \omega_{pmn}, \quad (34a)$$

$$\hat{\omega}_{pma} = F_{pm}^{\alpha}\Phi_{\alpha a}, \quad (34b)$$

$$\hat{\omega}_{pab} = \frac{1}{2}\Phi_a^{\alpha}V_p^{\mu}\partial_{\mu}\Phi_{\alpha\beta} - (a \leftrightarrow b), \quad (34c)$$

$$\hat{\omega}_{cmn} = -F_{mn}^{\alpha}\Phi_{\alpha c}, \quad (34d)$$

$$\hat{\omega}_{cab} = 0. \quad (34e)$$

We can now conclude this section by carrying out the multiplication and examining the result, which gives

$$\boxed{S = \int dx^D \int V \delta \left(\frac{-1}{4} g^{\rho\lambda} \partial_{\lambda} h_{\alpha\beta} \partial_{\rho} h^{\alpha\beta} + \frac{1}{4} F^{\mu\nu\alpha} F_{\mu\nu}^{\beta} h_{\alpha\beta} + \right.} \quad (35)$$

$$\left. + R + \frac{1}{4} \partial_{\hat{\mu}} \log(h) \partial^{\hat{\mu}} \log(h) + \partial_{\hat{\mu}} \log(h) \partial^{\hat{\mu}} \hat{\phi} \right).$$

We can observe some interesting things. First, we recover the classical Einstein action, coupled to E electromagnetic fields. So we can conclude that gravity in $D + E$ dimensions is equivalent to gravity in D dimensions plus E electromagnetic fields, and we find a few additional fields which will prove useful for the dilaton field reduction.

4.5 Adding the Dilaton Field Reduction

For this part we sum $S_{\hat{g}}$ and $S_{\hat{\phi}}$, which gives

$$\begin{aligned}
S_{\hat{g}+\hat{\phi}} &= \int dx^D \int V \delta \left(\frac{-1}{4} g^{\rho\lambda} \partial_{\lambda} h_{\alpha\beta} \partial_{\rho} h^{\alpha\beta} + \frac{1}{4} F^{\mu\nu\alpha} F_{\mu\nu}^{\beta} h_{\alpha\beta} + \right. \\
&\quad \left. + R + \frac{1}{4} \partial_{\hat{\mu}} \log(h) \partial^{\hat{\mu}} \log(h) + \partial_{\hat{\mu}} \log(h) \partial^{\hat{\mu}} \hat{\phi} + \partial_{\hat{\mu}} \hat{\phi} \partial^{\hat{\mu}} \hat{\phi} \right).
\end{aligned}$$

We then introduce the shifted dilaton

$$\phi = \hat{\phi} - \frac{1}{2} \log h = \hat{\phi} - \log \delta, \quad (36)$$

which gives

$$S = \int dx^D V e^{-\phi} \left(R - \frac{1}{4} F^{\mu\nu\alpha} F_{\mu\nu}^{\beta} h_{\alpha\beta} + \right. \\ \left. - g^{\rho\lambda} \partial_\lambda h_{\alpha\beta} \partial_\rho h^{\alpha\beta} + \hat{g}^{\hat{\mu}\hat{\nu}} \partial_{\hat{\mu}} \phi \partial_{\hat{\nu}} \phi \right). \quad (37)$$

5 Two-Form Field Reduction

5.1 The Reduction

For the two-form we will carry out the reduction separately from the other two fields; this causes no issue, as the action is linear in it.

We can rewrite (21) as

$$\mathcal{L}_{\hat{B}} = \frac{-1}{12} \hat{H}_{\hat{\mu}\hat{\nu}\hat{\rho}} \hat{H}^{\hat{\mu}\hat{\nu}\hat{\rho}}$$

Here we use (4) to pass the lower-index H to the flat-space basis, and reuse the identity to change the upper index as well:

$$= \frac{-1}{12} \hat{H}_{\hat{r}\hat{s}\hat{t}} \hat{V}_{\hat{\mu}}^{\hat{r}} \hat{V}_{\hat{\nu}}^{\hat{s}} \hat{V}_{\hat{\rho}}^{\hat{t}} \hat{H}^{\hat{\mu}\hat{\nu}\hat{\rho}} \\ = \frac{-1}{12} (\hat{H}_{\hat{r}\hat{s}\hat{t}} \hat{H}^{\hat{r}\hat{s}\hat{t}})$$

We then expand the term, separating the internal space from space-time:

$$= \frac{-1}{12} (\hat{H}_{rst} \hat{H}^{rst} + \hat{H}_{rsc} \hat{H}^{rsc} + \hat{H}_{rbt} \hat{H}^{rbt} + \hat{H}_{rbc} \hat{H}^{rbc} + \hat{H}_{ast} \hat{H}^{ast} + \hat{H}_{asc} \hat{H}^{asc} + \hat{H}_{abs} \hat{H}^{abs} + \hat{H}_{abc} \hat{H}^{abc}).$$

However, thanks to the fact that $\hat{B}_{\hat{\mu}\hat{\nu}}$ is antisymmetric, one can easily show that $\hat{H}_{\hat{\mu}\hat{\nu}\hat{\rho}}$ is antisymmetric under exchange of any two neighboring indices. So we obtain

$$\mathcal{L}_{\hat{B}} = \frac{-1}{12} (\hat{H}_{rst} \hat{H}^{rst} + 3 \hat{H}_{rsc} \hat{H}^{rsc} + 3 \hat{H}_{asc} \hat{H}^{asc} + \hat{H}_{abc} \hat{H}^{abc}), \quad (38)$$

and, finally, the action for B is written as

$$S_{\hat{b}} = - \int dx^D \sqrt{-g} e^{-\phi} \left\{ \frac{1}{12} H_{\alpha\beta\gamma} H^{\alpha\beta\gamma} + \frac{1}{4} H_{\mu\alpha\beta} H^{\mu\alpha\beta} + \frac{1}{4} H_{\mu\nu\alpha} H^{\mu\nu\alpha} + \frac{1}{12} H_{\mu\nu\rho} H^{\mu\nu\rho} \right\}. \quad (39)$$

The absence of a hat is due to the fact that we define the reduced, lower-dimensional form as

$$T^\mu = V_r^\mu \hat{V}_{\hat{\mu}}^r \hat{T}^{\hat{\mu}}, \quad (40a)$$

$$T^\alpha = V_a^\alpha \hat{V}_{\hat{\mu}}^a \hat{T}^{\hat{\mu}}, \quad (40b)$$

$$T_\mu = V_\mu^r \hat{V}_{\hat{\mu}}^r \hat{T}_{\hat{\mu}}, \quad (40c)$$

$$T_\alpha = V_\alpha^a \hat{V}_{\hat{\mu}}^a \hat{T}_{\hat{\mu}}. \quad (40d)$$

With that we obtain (39).

We now compute each component of the un-hatted H as a function of the hatted H .

Starting slowly, as we have seen above $\hat{B}_{\alpha\beta} = B_{\alpha\beta}$, and using the y -independence we get $H_{\alpha\beta\gamma} = 0$.

We continue with $H_{\mu\alpha\beta}$:

$$H_{\mu\alpha\beta} = V_\mu^r \hat{V}_{\hat{\mu}}^r \hat{H}_{\hat{\mu}\alpha\beta} \\ = \partial_\mu B_{\alpha\beta}. \quad (41)$$

Those two were fairly straightforward; the next two are more involved, and we introduce new objects along the way. Now for $H_{\mu\nu\alpha}$:

$$H_{\mu\nu\alpha} = V_\mu^r V_\nu^s \hat{V}_{\hat{\mu}}^r \hat{V}_{\hat{\nu}}^s \hat{H}_{\hat{\mu}\hat{\nu}\alpha} \\ = \hat{H}_{\mu\nu\alpha} - (A_\mu^{(1)\alpha} \hat{H}_{\alpha\nu\rho} + A_\rho^{(1)\alpha} \hat{H}_{\mu\alpha\nu}) \\ = \partial_\mu \hat{B}_{\nu\alpha} + \partial_\nu \hat{B}_{\alpha\mu} - A_\mu^{(1)\beta} \partial_\nu \hat{B}_{\alpha\beta} - A_\nu^{(1)\beta} \partial_\mu \hat{B}_{\beta\alpha} \\ = \partial_\mu \hat{B}_{\nu\alpha} + \partial_\nu \hat{B}_{\alpha\mu} - \partial_\nu (A_\mu^{(1)\beta} \hat{B}_{\alpha\beta}) + \partial_\nu A_\mu^{(1)\beta} \hat{B}_{\alpha\beta} - \partial_\mu (A_\nu^{(1)\beta} \hat{B}_{\beta\alpha}) + \partial_\mu A_\nu^{(1)\beta} \hat{B}_{\beta\alpha} \\ = \partial_\mu \hat{B}_{\nu\alpha} - \partial_\nu \hat{B}_{\mu\alpha} - \partial_\nu (A_\mu^{(1)\beta} \hat{B}_{\alpha\beta}) + \partial_\nu A_\mu^{(1)\beta} \hat{B}_{\alpha\beta} + \partial_\mu (A_\nu^{(1)\beta} \hat{B}_{\alpha\beta}) - \partial_\mu A_\nu^{(1)\beta} \hat{B}_{\alpha\beta} \\ = \partial_\mu (\hat{B}_{\nu\alpha} + A_\nu^{(1)\beta} \hat{B}_{\alpha\beta}) - \partial_\nu (\hat{B}_{\mu\alpha} + A_\mu^{(1)\beta} \hat{B}_{\alpha\beta}) - \hat{B}_{\alpha\beta} (\partial_\mu A_\nu^{(1)\beta} - \partial_\nu A_\mu^{(1)\beta}) \\ = F_{\mu\nu\alpha}^{(2)} - \hat{B}_{\alpha\beta} F_{\mu\nu}^{(1)\beta}, \quad (42)$$

where

$$F_{\mu\nu\alpha}^{(2)} = \partial_\mu A_{\nu\alpha}^{(2)} - \partial_\nu A_{\mu\alpha}^{(2)},$$

and

$$A_{\mu\alpha}^{(2)} = \hat{B}_{\mu\alpha} + B_{\alpha\beta} A_\mu^{(1)\beta}.$$

Finally, for $H_{\mu\nu\rho}$ we use exactly the same technique as before, which gives

$$\begin{aligned} H_{\mu\nu\rho} &= V_\mu^r V_\nu^s V_\rho^t \hat{V}_r^{\hat{\mu}} \hat{V}_s^{\hat{\nu}} \hat{V}_t^{\hat{\rho}} \hat{H}_{\hat{\mu}\hat{\nu}\hat{\rho}} \\ &= \hat{H}_{\mu\nu\rho} - A_\mu^{(1)\alpha} \hat{H}_{\alpha\nu\rho} - A_\nu^{(1)\alpha} \hat{H}_{\alpha\rho\mu} - A_\rho^{(1)\alpha} \hat{H}_{\alpha\mu\nu} + A_\mu^{(1)\alpha} A_\nu^{(1)\beta} \hat{H}_{\alpha\beta\rho} + A_\nu^{(1)\alpha} A_\rho^{(1)\beta} \hat{H}_{\alpha\beta\mu} + A_\rho^{(1)\alpha} A_\mu^{(1)\beta} \hat{H}_{\alpha\beta\nu} \\ &= \partial_\mu B_{\nu\rho} - \frac{1}{2} (A_\mu^{(1)\alpha} F_{\nu\rho\alpha}^{(2)} + A_{\mu\alpha}^{(2)} F_{\nu\rho}^{(1)}) + \partial_\nu B_{\rho\mu} - \frac{1}{2} (A_\nu^{(1)\alpha} F_{\rho\mu\alpha}^{(2)} + A_{\nu\alpha}^{(2)} F_{\rho\mu}^{(1)}) + \partial_\rho B_{\mu\nu} - \frac{1}{2} (A_\rho^{(1)\alpha} F_{\mu\nu\alpha}^{(2)} + A_{\rho\alpha}^{(2)} F_{\mu\nu}^{(1)}), \end{aligned} \quad (43)$$

where

$$B_{\mu\nu} = \hat{B}_{\mu\nu} + \frac{1}{2} A_\mu^{(1)\alpha} A_{\nu\alpha}^{(2)} - \frac{1}{2} A_\nu^{(1)\alpha} A_{\mu\alpha}^{(2)} - A_\mu^{(1)\alpha} B_{\alpha\beta} A_\nu^{(2)\beta}. \quad (44)$$

5.2 Observation of Gauge

Now, as in Section 4.3, we check the gauge transformations to make sure every object transforms as it should. Let us begin with $A^{(2)}$, which should transform under the Lie derivative and under the new transformation $B_{\mu\nu} \rightarrow B_{\mu\nu} + \partial_\mu \Lambda_\nu$. So let us begin with $A_{\mu\alpha}^{(2)}$:

$$\begin{aligned} \delta A_{\mu\alpha}^{(2)} &= \delta(\hat{B}_{\mu\alpha} + B_{\alpha\beta} A_\mu^{(1)\beta}) \\ &= \delta(\hat{B}_{\mu\alpha}) + \delta(B_{\alpha\beta} A_\mu^{(1)\beta}) \\ &= \delta(\hat{B}_{\mu\alpha}) + \delta(B_{\alpha\beta}) A_\mu^{(1)\beta} + B_{\alpha\beta} \delta(A_\mu^{(1)\beta}) \\ &= L_\xi \hat{B}_{\mu\alpha} + \partial_\mu \Lambda_\alpha^{(2)} + A_\mu^{(1)\beta} L_\xi \hat{B}_{\alpha\beta} + \hat{B}_{\alpha\beta} \partial_\mu \Lambda^{(1)\alpha} \\ &= L_\xi \hat{B}_{\mu\alpha} + \partial_\mu \Lambda_\alpha^{(2)} + L_\xi (A_\mu^{(1)\beta} \hat{B}_{\alpha\beta}) \\ &= \partial_\mu \Lambda_\alpha^{(2)} + L_\xi A_{\mu\alpha}^{(2)}, \end{aligned} \quad (45)$$

so we obtain the correct transformation for this object.

Doing the same for $B_{\mu\nu}$, we find the transformation

$$\delta B_{\mu\nu} = \frac{1}{2} (\Lambda^{(1)\alpha} F_{\mu\nu\alpha}^{(2)} + \Lambda_\alpha^{(2)} F_{\mu\nu}^{(1)\alpha}), \quad (46)$$

which is exactly the right term to compensate for the additional term in the transformation of $H_{\mu\nu\rho}$.

6 Complete Reduced Action

To summarize, the complete reduced action takes the form

$$S = \int dx^D V e^{-\phi} \mathcal{L}, \quad (47)$$

where we decompose S as $S = S_1 + S_2 + S_3 + S_4$, with

$$\begin{aligned} S_1 &= \int dx^D V e^{-\phi} (R + g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi), \\ S_2 &= \int dx^D V e^{-\phi} \left(\frac{1}{4} g^{\mu\nu} (\partial_\mu h_{\alpha\beta} \partial_\nu h^{\alpha\beta} - h^{\alpha\beta} h^{\gamma\delta} \partial_\mu B_{\alpha\gamma} \partial_\nu B_{\beta\delta}) \right), \\ S_3 &= \int dx^D V e^{-\phi} \left(-\frac{1}{4} g^{\mu\rho} g^{\nu\lambda} (h_{\alpha\beta} F_{\mu\nu}^{(1)\alpha} F_{\rho\lambda}^{(1)\beta} + h^{\alpha\beta} H_{\mu\nu\alpha} H_{\rho\lambda\beta}) \right), \\ S_4 &= \int dx^D V e^{-\phi} \left(-\frac{1}{12} H_{\mu\nu\rho} H^{\mu\nu\rho} \right). \end{aligned} \quad (48)$$

7 Observation of a New Symmetry

7.1 The O(E,E) Group

For this part we explore a very important topic in physics research today: symmetry. In the present context it will allow us to establish equivalences between multiple theories. First, we note that we have an $SL(E, \mathbb{R})$ symmetry, given by every $E \times E$ matrix of determinant one. Another symmetry can be seen from the fact that, since B only appears within a derivative, we may add a constant to it as we please; and, as shown earlier in Sections 5.2 and 4.3, we also have some gauge freedom that we may use as we please. However, all of these symmetries are manifest in high dimensions. What interests us in the reduction is that it neither creates nor destroys any symmetry: if we observe a new symmetry in the reduced form, that means we have effectively discovered a new symmetry in the unreduced form. Following [3], we observe an $O(E, E)$ symmetry.

Definition 1 A matrix M is in $O(E, E)$ if it is a square $E \times E$ matrix and, for the matrix

$$\zeta = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix},$$

which can be used as a metric for the group $O(E, E)$, as it has $d - 1$ negative eigenvalues and d positive eigenvalues in a basis rotated from the one with diagonal metric, the matrix M satisfies

$$\boxed{M^T \zeta M = \zeta.}$$

7.2 Rewriting the Reduced Action

We now define a new matrix M ,

$$M = \begin{pmatrix} h^{-1} & -h^{-1}B \\ Bh^{-1} & h - Bh^{-1}B \end{pmatrix}. \quad (49)$$

We now try to rewrite the action in terms of M , to make things clearer. S_1 is a lost cause, as neither g nor ϕ is contained in M . So let us begin with S_2 , which can be rewritten in matrix form as

$$S_2 = \frac{1}{4} \text{tr}(\partial_\mu h^{-1} \partial^\mu h + h^{-1} \partial_\mu B h^{-1} \partial^\mu B).$$

We now perform a few manipulations:

$$\begin{aligned} S_2 &= \frac{1}{4} \text{tr}(\partial_\mu h^{-1} \partial^\mu h + h^{-1} \partial_\mu B h^{-1} \partial^\mu B) \\ &= \frac{1}{8} \text{tr}(2\partial_\mu h \partial^\mu h^{-1} + 2h^{-1} \partial_\mu B h^{-1} \partial^\mu B) \\ &= \frac{1}{8} \text{tr}(2\partial_\mu h \partial^\mu h^{-1} - 2\partial_\mu(Bh^{-1}B) \partial^\mu h^{-1} + \partial_\mu(Bh^{-1}) \partial^\mu(Bh^{-1}) + \partial_\mu(h^{-1}B) \partial^\mu(h^{-1}B)) \\ &= \frac{1}{8} \text{tr}(\partial_\mu(h - Bh^{-1}B) \partial^\mu h^{-1} + \partial_\mu(Bh^{-1}) \partial^\mu(Bh^{-1}) + \partial_\mu(h^{-1}B) \partial^\mu(h^{-1}B) + \partial^\mu h^{-1} \partial_\mu(h - Bh^{-1}B)) \\ &= \frac{1}{8} \text{tr}(\partial_\mu M^{-1} \partial^\mu M). \end{aligned} \quad (50)$$

We now rewrite S_3 by expanding the H terms, to get

$$\begin{aligned} S_3 &= -\frac{1}{4} \left[F_{\mu\nu}^{(1)\alpha} h_{\alpha\beta} F^{(1)\mu\nu\beta} + (F_{\mu\nu}^{(2)} - B_{\alpha\gamma} F_{\mu\nu}^{(1)\gamma}) h^{\alpha\beta} (F_{\beta}^{(2)\mu\nu} - B_{\beta\delta} F^{(1)\mu\nu\delta}) \right] \\ &= -\frac{1}{4} \mathcal{F}_{\mu\nu}^i (M^{-1})_{ij} \mathcal{F}^{\mu\nu j}. \end{aligned} \quad (51)$$

Here $\mathcal{F}_{\mu\nu}^i$ is the $2E$ -component vector of field strengths,

$$\mathcal{F}_{\mu\nu}^i = \begin{pmatrix} F_{\mu\nu}^{(1)\alpha} \\ F_{\mu\nu}^{(2)} \end{pmatrix} = \partial_\mu \mathcal{A}_\nu^i - \partial_\nu \mathcal{A}_\mu^i. \quad (52)$$

Finally, we turn to S_4 : first, let us rewrite $H_{\mu\nu\rho}$,

$$\begin{aligned} H_{\mu\nu\rho} &= \partial_\mu B_{\nu\rho} - \frac{1}{2} (A_\mu^{(1)\alpha} F_{\nu\rho\alpha}^{(2)} + A_{\mu\alpha}^{(2)} F_{\nu\rho}^{(1)}) + \partial_\nu B_{\rho\mu} - \frac{1}{2} (A_\nu^{(1)\alpha} F_{\rho\mu\alpha}^{(2)} + A_{\nu\alpha}^{(2)} F_{\rho\mu}^{(1)}) + \partial_\rho B_{\mu\nu} - \frac{1}{2} (A_\rho^{(1)\alpha} F_{\mu\nu\alpha}^{(2)} + A_{\rho\alpha}^{(2)} F_{\mu\nu}^{(1)}) \\ &= \partial_\mu B_{\nu\rho} - \frac{1}{2} \mathcal{A}_\mu^i \zeta_{ij} \mathcal{F}_{\nu\rho}^j + (\text{cyclic perms.}) \end{aligned} \quad (53)$$

We can now rewrite the complete action in terms of M , which gives

$$\begin{aligned} S &= \int dx^D V e^{-\phi} \left[R + g^{\mu\nu} \partial_\mu \phi \partial_\nu \phi + \frac{1}{8} \text{tr}(\partial_\mu M^{-1} \partial^\mu M) - \frac{1}{4} \mathcal{F}_{\mu\nu}^i (M^{-1})_{ij} \mathcal{F}^{\mu\nu j} \right. \\ &\quad \left. - \frac{1}{12} \left(\partial_\mu B_{\nu\rho} - \frac{1}{2} \mathcal{A}_\mu^i \zeta_{ij} \mathcal{F}_{\nu\rho}^j + (\text{cyclic perms.}) \right) \left(\partial^\mu B^{\nu\rho} - \frac{1}{2} \mathcal{A}^{i\mu} \zeta_{ij} \mathcal{F}^{j\nu\rho} + (\text{cyclic perms.}) \right) \right]. \end{aligned} \quad (54)$$

7.3 Action Invariance under $O(E, E)$

We now study the effect of the $O(E, E)$ group on this action. Since it acts only on the internal indices, its action on the different parts of the action is

$$g_{\mu\nu} \rightarrow g_{\mu\nu}, \quad (55)$$

$$\phi \rightarrow \phi. \quad (56)$$

It is worth noting that, while ϕ as a whole is invariant, each of its individual components ($\hat{\phi}$ and $\frac{1}{2} \log(h)$) is not invariant under $O(E, E)$ on its own; this is precisely why we needed to introduce the shifted dilaton.

We also need to define the action of the group on the two-form, and on $A^{(2)}$ when it carries internal indices. So we have

$$A^i \rightarrow \Omega_j^i A^j, \quad (57)$$

$$M_{ij} \rightarrow \Omega_i^l M_{lk} \Omega_j^k, \quad (58)$$

where $\Omega \in O(E, E)$. We now show that $M \in O(E, E)$:

$$\begin{aligned} M^T \zeta M &= \begin{pmatrix} h^{-1} & Bh^{-1} \\ -h^{-1}B & h - Bh^{-1}B \end{pmatrix} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} h^{-1} & -h^{-1}B \\ Bh^{-1} & h - Bh^{-1}B \end{pmatrix} \\ &= \begin{pmatrix} Bh^{-1} & h^{-1} \\ h - Bh^{-1}B & -h^{-1}B \end{pmatrix} \begin{pmatrix} h^{-1} & -h^{-1}B \\ Bh^{-1} & h - Bh^{-1}B \end{pmatrix} \\ &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} = \zeta. \end{aligned}$$

We can now verify that every part of the action is invariant under the action of $O(E, E)$. For S_1 the invariance is manifest, as nothing depends on the internal space. For S_2 we can use the cyclicity of the trace, together with the action of $O(E, E)$ on M ,

$$M \rightarrow \Omega M \Omega^T,$$

to make the invariance under $O(E, E)$ more visible. For S_3 , using $\Omega^T \zeta \Omega = \zeta$, the second term is $O(E, E)$ -invariant, and the first term is invariant provided we require that $B_{\nu\rho}$ does not transform. This shows that the action hides a non-trivial symmetry, providing a useful framework for comparing equivalent theories. Explicitly, for S_3 we have, using (57),

$$S_{3'} = -\frac{1}{4} \mathcal{F}_{\mu\nu}^i \Omega_i^l \Omega_n^i (M^{-1})_{ij} \Omega_m^j \Omega_j^k \mathcal{F}^{\mu\nu j} \quad (59)$$

$$= -\frac{1}{4} \mathcal{F}_{\mu\nu}^i \Omega_j^k \eta_{kn} \Omega_n^i (M^{-1})_{ij} \Omega_m^j \eta_{ml} \Omega_i^l \mathcal{F}^{\mu\nu j} \quad (60)$$

$$= -\frac{1}{4} \mathcal{F}_{\mu\nu}^i \eta_{ji} (M^{-1})_{ij} \eta_{ij} \mathcal{F}^{\mu\nu j} \quad (61)$$

$$= -\frac{1}{4} \mathcal{F}_{\mu\nu}^i (M^{-1})_{ij} \mathcal{F}^{\mu\nu j} \quad (62)$$

$$= S_3. \quad (63)$$

The same argument can be repeated for S_4 , with the same result. We have thus proven that the whole action is $O(E, E)$ -invariant. This demonstrates that the action hides a non-trivial symmetry, providing a useful framework for comparing equivalent theories.

8 Conclusion

In this internship we first developed a description of General Relativity through the vielbein formalism, then laid out the ground rules of classical field theory. Using these two frameworks we constructed the NS-NS action in $D + E$ dimensions and carried out its Kaluza-Klein reduction to D dimensions, under the assumption that no field depends on the internal coordinates. This reduction produced a D -dimensional action containing the graviton, a dilaton, E vector fields, a two-form, and the scalar matrix $h_{\alpha\beta}$ encoding the geometry of the internal space. Combining h and the reduced two-form B into the single matrix M , we showed that the reduced action is invariant under an $O(E, E)$ group acting on M and on the vector fields, a symmetry that is not visible in the original $D + E$ -dimensional theory.

This hidden $O(E, E)$ symmetry means that theories which appear distinct before reduction differing, for instance, in the background values of h and B describe the same D -dimensional physics once reduced. Practically, this lets us treat such theories as equivalent, using whichever formulation is most convenient, rather than working them out independently.

Two natural directions extend this work. The first is to relax the assumption of internal-coordinate independence and study the resulting field dependence explicitly. The second is to look for larger hidden symmetry groups, such as $O(E, E + n)$, which would extend the equivalences found here to a wider class of theories.

References

- ¹J. Scherk and J. H. Schwarz, “How to get masses from extra dimensions”, Nuclear Physics B **153**, 61–88 (1979).
- ²A. Einstein, “Zur allgemeinen relativitätstheorie”, Sitzungsber. preuss. Akad. Wiss **44**, 778–786 (1915).
- ³J. Maharana and J. H. Schwarz, “Noncompact symmetries in string theory”, Nuclear Physics B **390**, 3–32 (1993).
- ⁴D. Z. Freedman and A. Van Proeyen, *Supergravity* (Cambridge university press, 2012).

A General Relativity Background

Definition 2 Let V be an $\mathbb{R}$ -vector space of dimension N . We define the tensor of order $(r, s) \in \llbracket 0, 1 \rrbracket^2$ on $V^{(r,s)} \stackrel{\text{def}}{=} V^{\otimes r} \otimes V^{\otimes s}$ as a multilinear map from $V^{\star r} \times V^s$ to $\mathbb{R}$.

Definition 3 We call an n -dimensional ($n \in \mathbb{N}^*$) manifold every Hausdorff topological space $(\mathcal{M}, \tau)$ locally homeomorphic to $\mathbb{R}^n$.

Definition 4 Let $(\mathcal{M}, g)$ be a smooth manifold. A metric tensor is a symmetric, non-degenerate bilinear form $g_{\mu\nu}$ on the tangent space $T_p\mathcal{M}$ at each point $p \in \mathcal{M}$, such that

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu.$$

In General Relativity we use the signature $(-, +, +, +)$.

Definition 5 In General Relativity we generally use the Christoffel connection, as it is the only connection that is both metric-compatible and torsion-free.

We define it as

$$\nabla_\mu V^\nu = \partial_\mu V^\nu + \Gamma_{\mu\sigma}^\nu V^\sigma, \quad (64)$$

where the Christoffel symbols are

$$\Gamma_{\mu\nu}^\lambda = \frac{1}{2} g^{\lambda\sigma} (\partial_\mu g_{\nu\sigma} + \partial_\nu g_{\sigma\mu} - \partial_\sigma g_{\mu\nu}). \quad (65)$$

Definition 6 The Riemann curvature tensor measures the failure of covariant derivatives to commute:

$$[\nabla_\mu, \nabla_\nu] V^\rho = R^\rho_{\sigma\mu\nu} V^\sigma, \quad (66)$$

and is given in terms of the Christoffel symbols by

$$R^\rho_{\sigma\mu\nu} = \partial_\mu \Gamma_{\nu\sigma}^\rho - \partial_\nu \Gamma_{\mu\sigma}^\rho + \Gamma_{\mu\lambda}^\rho \Gamma_{\nu\sigma}^\lambda - \Gamma_{\nu\lambda}^\rho \Gamma_{\mu\sigma}^\lambda. \quad (67)$$

Contracting the first and third indices gives the Ricci tensor, $R_{\mu\nu} = R^\rho_{\mu\rho\nu}$, and contracting again with the inverse metric gives the Ricci scalar,

$$R = g^{\mu\nu} R_{\mu\nu}. \quad (68)$$

It is this object, written $\hat{R}$ in $D + E$ dimensions, that appears in the Einstein–Hilbert term of (3.2) and in the NS-NS action (22); in the vielbein formalism used throughout this report, it is expressed via $\hat{\omega}$ rather than Γ , as derived in Section 2 and Eq. (4.1).

B Euler–Lagrange Equation

This is the general derivation underlying the field equations of motion used throughout the report, in particular (17) and (18).

By using the principle of least action and the variational principle, we can see that

$$\begin{aligned} \delta S &= \delta \int \mathcal{L} dx^\mu \\ &= \delta \int \left(\frac{\partial \mathcal{L}}{\partial \eta_\rho} \delta \eta_\rho + \frac{\partial \mathcal{L}}{\partial \partial_\nu \eta_\rho} \delta \partial_\nu \eta_\rho \right) dx^\mu \\ &= \delta \int \left(\frac{\partial \mathcal{L}}{\partial \eta_\rho} \delta \eta_\rho - \partial_\nu \frac{\partial \mathcal{L}}{\partial \partial_\nu \eta_\rho} \delta \eta_\rho \right) + \underbrace{\partial_\nu \left(\frac{\partial \mathcal{L}}{\partial \partial_\nu \eta_\rho} \delta \eta_\rho \right)}_{\text{boundary term, equal to zero}} dx^\mu \\ &= 0, \end{aligned}$$

which gives the equation

$$\boxed{\frac{\partial \mathcal{L}}{\partial \eta_\rho} - \partial_\nu \frac{\partial \mathcal{L}}{\partial \partial_\nu \eta_\rho} = 0}, \quad (69)$$

which is the equation of motion for the field, applied e.g. to obtain $\square \Phi = 0$, $\partial_\mu F^{\mu\nu} = 0$ and $\partial_\mu H^{\mu\nu\rho} = 0$ from (19), (20) and (21) respectively.

C Summary of the Reduced Field Content

Because the reduction in Sections 4–6 introduces a large number of new fields, this table collects them for reference: where each one comes from, and where it is first defined.

Field	Origin	First defined
V_μ^r	D -dimensional vielbein (graviton)	(25)
Φ_α^a	internal-space vielbein (moduli), $h_{\alpha\beta} = \Phi_\alpha^a \Phi_{a\beta}$	(25), Sec. 4.2
$A_\mu^{(1)\alpha}$	off-diagonal vielbein block; Kaluza–Klein gauge field	(25)
ϕ	shifted (four-dimensional) dilaton, $\phi = \hat{\phi} - \log \delta$	Sec. 4.6
$B_{\mu\nu}$	reduced two-form	Sec. 5.1
$A_{\mu\alpha}^{(2)}$	mixed component of $\hat{B}$; second gauge field	Sec. 5.1
$\mathcal{A}_\mu^i = (A_\mu^{(1)\alpha}, A_{\mu\alpha}^{(2)})$	$2E$ -component vector combining both gauge fields	Sec. 6.2
M	moduli matrix built from h, B ; carries the $O(E, E)$ action	Sec. 6.2