

---

# PROBABILITY THEORY

---

## Study Note

### **Author**

Sam Ren  
Grinnell College  
February 5, 2024

# Contents

<b>1</b>	<b>Measure Theory</b>	<b>4</b>
<b>2</b>	<b>Probability Measure</b>	<b>5</b>
2.1	Discrete & Continuous Probability measure . . . . .	6
2.2	Binomial Distribution . . . . .	6
2.3	Product Probability Spaces . . . . .	7
2.4	Hypergeometric Distribution(Multivariate) . . . . .	7
<b>3</b>	<b>Conditional Probability</b>	<b>7</b>

## Acknowledgements

I finished this study notes by simply collected the concepts and terms from @brightsideof-maths youtube videos, here is the link: [Video link](#). At the same time I would like to sincerely thank to my teachers and friends. Without them this note would not be done.

# 1 Measure Theory

This section is not necessary for later studying but it helps with understanding those abstract concepts

## 2 Probability Measure

Probability Measure is the measure with total mass 1, where we call it sample space  $\Omega$ ,  $Area(\Omega) = 1$ . In math we write:

$$\mathbb{P} : \mathcal{A} \rightarrow \mathbb{R}$$

Where  $\mathcal{A}$  is the collection of subsets, then we have:

$$\mathbb{P}(\Omega) = 1, \mathbb{P}(A) \in [0, 1]$$

It is easy to notice that if two sets  $(A, B)$  are disjoint ( $A \cap B = \emptyset$ ) then we have:

$$\mathbb{P}(A \cup B) = \mathbb{P}(A) + \mathbb{P}(B)$$

**Theorem 2.1.** If we have a sequence of disjoint sets  $A_i, i \in \mathbb{N}$  then we have:

$$\mathbb{P}\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} \mathbb{P}(A_i)$$

**Definition 2.2.** Let  $\Omega$  be a set. A collection of subsets  $\mathcal{A} \subseteq \mathcal{P}(\Omega)$  ( $\mathcal{P}$  is the power set) is called a sigma algebra ( $\sigma$ -Algebra) if:

1.  $\Omega, \emptyset \in \mathcal{A}$
2. If  $A \in \mathcal{A}$  then  $A^c \in \mathcal{A}$
3. If  $A_1, A_2, A_3, \dots \in \mathcal{A}$  then  $\bigcup_{i=1}^{\infty} A_i \in \mathcal{A}$

In Probability theory we call the elements in  $\sigma$ -algebra events. Then we are going to define the probability measure properly.

**Definition 2.3.** Let  $\mathcal{A} \subseteq \mathcal{P}(\Omega)$  is a sigma-algebra. A map  $\mathbb{P} : \mathcal{A} \rightarrow [0, 1]$  is called probability measure if:

1.  $\mathbb{P}(\Omega) = 1$
2.  $\mathbb{P}(\emptyset) = 0$
3.  $\mathbb{P}(\bigcup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} \mathbb{P}(A_i)$  where  $i \neq j$

**example 2.4.** If we have  $\Omega = \{1, 2, 3, 4, 5, 6\}$ , then  $\mathcal{A} = \mathcal{P}(\Omega)$  we have:

$$\mathbb{P} : \mathcal{A} \rightarrow [0, 1], \mathbb{P}(A) := \frac{|A|}{|\Omega|}$$

*Proof.* Prove that  $\mathbb{P}(A^c) = 1 - \mathbb{P}(A)$

□

## 2.1 Discrete & Continuous Probability measure

We are going to discuss two situations where the sample space is either discrete or continuous. Where at here the term **Discrete** means the sample space is either countable or finite. For term **Continuous** the sample space is either infinite or uncountable.

Type	discrete	continuous
Sample space	$\Omega$ countable or finite	$\Omega \subseteq \mathbb{R}^n$ uncountable, $\Omega \in \mathcal{B}(\mathbb{R}^n)$
$\sigma$ -algebra	$\mathcal{A} = \mathcal{P}(\Omega)$	$\mathcal{A} = \mathcal{B}(\Omega)$
Prob Measure	$\mathbb{P} : \mathcal{A} \rightarrow [0, 1]$	$\mathbb{P} : \mathcal{A} \rightarrow [0, 1]$
Determined by	$\mathbb{P}(\{\omega\}), \forall \omega \in \Omega$	
mass function	$(p_\omega)_{\omega \in \Omega}$ with $p_\omega \geq 0, \sum_{\omega \in \Omega} p_\omega = 1$	$f : \Omega \rightarrow \mathbb{R}$ with $f(x) \geq 0, \int_{\Omega} f(x) dx = 1$
Definition	$\mathbb{P}(A) = \sum_{\omega \in A} p_\omega = 1$	$\mathbb{P}(A) = \int_A f(x) dx$

Table 1: Discrete and continuous sample space explanation table

For continous probability measure we need to ensure that  $\Omega$  is measuralbe which we are going to discuss later. Here is an example for continuous measure:

**example 2.5.** For  $\Omega = [0, 2]$  we have prob density function  $f(x) = \frac{1}{2}, f : \Omega \rightarrow \mathbb{R}$ . Hence:

$$\mathbb{P}(\Omega) = \int_0^2 f(x) dx = \frac{1}{2} \int_0^2 1 dx = 1$$

Therefore we would have conclusion:

$$\mathbb{P}(A) = \int_A f(x) dx = \frac{1}{2} \int_A 1 dx = \frac{1}{2} (\text{Lebesgue measure}) \text{Length of } A$$

We would discuss the Lebesgue measure later.

## 2.2 Binomial Distribution

A binomial(two outcomes) distribution is foundational to the probability theory, and it also require a definition(For now this definition is improper):

**Definition 2.6.** A binomial distribution can be determined by following features:

1. There is no order
2. With replacements (That is, does not change the sample space)
3. There are only two outcomes

If it is a binomial distribution we denote it as  $\mathbb{P} = B(n, p)$ . For a binomial distribution we have:

$$\Omega = \{0, 1, 2, 3, \dots, n\}, \mathbb{P}(\{k\}) = \binom{n}{k} p^k (1-p)^{n-k}$$

## 2.3 Product Probability Spaces

In previous sections we already introduced the probability measure( $\mathbb{P}$ ), the  $\sigma$ -algebra( $\mathcal{A}$ ) and sample space( $\Omega$ ). In this section we are going to introduce the probability space which is important in precisely proof and mathematics content.

**Definition 2.7.** Probability space is a triple combination( $\Omega_n, \mathcal{A}_n, \mathbb{P}_n$ ),  $n \in \{1, 2, 3, \dots\}$ . Then the product space( $\Omega, \mathcal{A}, \mathbb{P}$ ) is given by:

- $\Omega = \Omega_1 \times \Omega_2 \times \Omega_3 \times \dots = \prod_{i \in \mathbb{N}} \Omega_i$
- $\mathcal{A} = \sigma(\Omega_1 \times A_2 \times \Omega_3 \times \dots)$  The choice of  $A_n \in \mathcal{A}$  is not restricted by 2, it can be any number. Hence we just simply replace one  $\Omega_i$  with  $A_i$  to construct.
- $\mathbb{P}(A_1 \times A_2 \times A_3 \times \dots \times A_m \times \Omega_{m+1} \times \Omega_{m+2} \times \dots) = \mathbb{P}(A_1) \cdot \mathbb{P}(A_2) \cdot \dots \cdot \mathbb{P}(A_m)$

You can find examples in this [Link](#).

## 2.4 Hypergeometric Distribution(Multivariate)

For this distribution we consider the sample size  $n$  and without replacement and unordered. Let's begin with an example:

**example 2.8.** Let elements in set  $S$  be the color of balls:

$$S = \{0, 1, 2, 3\}$$

If we have two 0 and 1 for others then we have a  $(2, 1, 1, 1)$ . At the same time there are many functions where mapping set  $S$  to  $\mathbb{N}$ . We denote the set of these functions as  $\mathbb{N}^S$ . We now can define the sample space:

$$\Omega = \{(k_s)_{s \in S} \in \mathbb{N}^S \mid \sum_{s \in S} k_s = n\}$$

Let's denote  $N_s$  as the number of balls in color  $s$  and  $N$  as the total number: Then we would have:

$$\mathbb{P}((k_s)_{s \in S}) = \frac{\prod_{s \in S} \binom{N_s}{k_s}}{\binom{N}{k}}$$

**Definition 2.9.** For hypergeometric distribution, we have (variant)set  $S = \{0, 1, 2, \dots, n\}$  and sample space  $\Omega = \{0, 1, 2, 3, \dots, n\}$ :

$$\mathbb{P}((k_s)_{s \in S}) = \frac{\prod_{s \in S} \binom{N_s}{k_s}}{\binom{N}{k}}$$

## 3 Conditional Probability

## References