

CLS Example

This example shows how to compute Classical Least Squares (CLS) predictions.

Copyright 2022-2023 Battelle Memorial Institute

Supporting Information

A Practical Guide to Chemometric Analysis of Optical Spectroscopic Data

Hope E. Lackey,^{1,2} Rachel L. Sell,¹ Gilbert L. Nelson,^{3*} Thomas A. Bryan,^{4*} Amanda M. Lines,^{1,2*} Samuel A. Bryan,^{1,2*}

¹ Pacific Northwest National Laboratory, 902 Battelle Boulevard, Richland, WA 99352

² Washington State University, Department of Chemistry, Pullman WA 99164

³ College of Idaho, Department of Chemistry, 2112 Cleveland Blvd, Caldwell, ID 83605

⁴ The MathWorks, 3 Apple Hill Drive, Natick, MA 01760-2098

*Email: sam.bryan@pnnl.gov Phone 1 509 375 5648; orcid.org/0000-0002-8826-0880

*Email: tbryan@mathworks.com Phone 1 508 647 7669

*Email: amanda.lines@pnnl.gov Phone: 1 509 375 5689

*Email: gnelson@collegeofidaho.edu Phone: 1 208 459 5241

CLS Algorithm

The CLS Algorithm is encapsulated in the following MATLAB function.

```
function [C_cls, B_cls, K_cls] = pnnl_cls(A_train, C_train, A_unknown)
%pnnl_cls Classical least squares (CLS) regression
%
% [C_cls, B_cls, K_cls] = pnnl_cls(A_train, C_train, A_unknown)
% returns concentration matrix C_cls, the least-squares solution that
% minimizes norm(C_cls*K_cls - A_unknown), of the Beer's law
% relationship CK=A. Extinction coefficient matrix K_cls is the
% least-squares solution that minimizes norm(C_train*K - A_train)
% where A_train is a matrix of training spectra corresponding to
% known concentrations in the C_train matrix. Multiplier matrix
% B_cls is the pseudo-inverse of Beer's law extinction coefficient
% matrix K_cls such that C_cls = A_unknown * B_cls.
%
% Example:
%
% load pnnl_napalm_data
% [C_cls, B_cls, K_cls] = pnnl_cls(A_train, C_train, A_unknown);
%
% See also pnnl_pcr, pnnl_pls.
%
% Copyright 2022-2023 Battelle Memorial Institute
%
% Compute K that minimizes norm(CK - A) given C and A
```

```

K_cls = C_train \ A_train;

% Compute C that minimizes norm(CK - A) given A and K
C_cls = A_unknown / K_cls;

% Multiplier matrix B_cls is the pseudo-inverse of Beer's law
% extinction coefficient matrix K_cls such that
% C_cls = A_unknown * B_cls.
B_cls = pinv(K_cls);

```

```
end
```

Concentration Data

The concentrations of the training data are in matrix C_{train} and the concentrations of the validation data are in matrix $C_{\text{validation}}$. Column 1 corresponds to the concentrations in constituent 1 (benzene). Column 2 corresponds to the concentrations in constituent 2 (polystyrene). Column 3 corresponds to the concentrations in constituent 3 (gasoline).

$$C_{\text{train}} = \begin{array}{c} \begin{array}{ccc} \textit{benzene} & \textit{polystyrene} & \textit{gasoline} \end{array} \\ \begin{bmatrix} 0 & 0 & 100.0000 \\ 5.1309 & 0 & 94.8691 \\ 10.0660 & 0 & 89.9300 \\ 20.1799 & 0 & 79.8201 \\ 40.0120 & 0 & 59.9878 \\ 59.9972 & 0 & 40.0028 \\ 79.8412 & 0 & 20.1588 \\ 89.8273 & 0 & 10.1727 \\ 100.0000 & 0 & 0 \\ 90.0264 & 9.9736 & 0 \\ 80.1375 & 19.8625 & 0 \\ 64.9950 & 35.0005 & 0 \\ 21.0228 & 45.9197 & 33.0575 \\ 49.9507 & 5.0599 & 44.9895 \\ 40.0182 & 20.0385 & 39.9433 \\ 40.0154 & 10.0036 & 49.9810 \\ 30.0059 & 10.0282 & 59.9659 \\ 40.0340 & 39.9670 & 19.9990 \\ 49.9393 & 3.3748 & 46.6859 \\ 46.6501 & 13.4658 & 39.8840 \end{bmatrix} \end{array} \begin{array}{c} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 8 \\ 9 \\ 10 \\ 11 \\ 12 \\ 13 \\ 14 \\ 15 \\ 16 \\ 17 \\ 18 \\ 19 \\ 20 \end{array}$$

$$C_{\text{validation}} = \begin{array}{c} \begin{array}{ccc} \textit{benzene} & \textit{polystyrene} & \textit{gasoline} \end{array} \\ \begin{bmatrix} 22.1665 & 39.0384 & 38.7951 \\ 21.6874 & 37.0596 & 41.2530 \\ 22.1665 & 39.6980 & 38.1355 \\ 26.9575 & 40.3576 & 32.6849 \\ 25.9993 & 33.7616 & 40.2391 \\ 23.1247 & 37.0596 & 39.8157 \\ 22.6456 & 39.0384 & 38.3160 \\ 22.6456 & 39.0384 & 38.3160 \\ 27.4366 & 42.3364 & 30.2270 \\ 27.4366 & 37.7192 & 34.8442 \\ 26.4784 & 40.3576 & 33.1640 \\ 26.9575 & 37.7192 & 35.3233 \end{bmatrix} \end{array} \begin{array}{c} 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 8 \\ 9 \\ 10 \\ 11 \\ 12 \end{array}$$

Clear variables and load the PNNL napalm data.

```
clearvars
load pnnl_napalm_data
```

Compute CLS

Set up the plot title and color.

```
title_string = sprintf('CLS');
nConstituents = size(C_validation,2);
colorOrder = pnnl_colorOrder(nConstituents);
```

Use all the columns of C_train to compute CLS. Use all the columns of C_validation to compute RMSEP (root mean square error predicted). Use the columns of C_train to compute RMSEC (root mean square error calibration) and RMSECV (root mean square error cross validation).

Compute CLS

```
C_predicted = pnnl_cls(A_train,C_train,A_unknown);
C_calibration = pnnl_cls(A_train,C_train,A_train);
C_cross_validation = pnnl_cross_validation(@pnnl_cls,A_train,C_train);
```

Compute RMSE

```
RMSEP = pnnl_rmse(C_validation,C_predicted);
RMSEC = pnnl_rmse(C_train,C_calibration);
RMSECV = pnnl_rmse(C_train,C_cross_validation);
```

Display RMSE

```
pnnl_display_rmse(title_string,ConstituentNames,...
    RMSEC,RMSECV,RMSEP);
```

```
-----
      CLS      Benzene      Polystyrene      Gasoline
      RMSEC      4.2721      2.3784      5.6857
      RMSECV      5.1043      3.0807      6.7847
      RMSEP      7.895      2.6988      9.1105
      -----
```

Plot the results


```
figure
h = gobjects(nConstituents,1);
for k = 1:nConstituents
    % Plot Concentrations
    hold on
    % Validation vs. Predicted
    h(k) =
    plot(C_validation(:,k),C_predicted(:,k),'.','MarkerSize',35,'Color',colorOrder(k,:),
        'DisplayName',ConstituentNames{k});
    % Train vs. Calibration
    plot(C_train(:,k),C_calibration(:,k),'o','MarkerSize',10,'LineWidth',1,'Color',colorOrder(k,:))
```

% Train vs. Crosss Validation

```

plot(C_train(:,k),C_cross_validation(:,k),'+', 'MarkerSize',10, 'LineWidth',1,
'Color',colorOrder(k,:))
% 1-1 line
line(C_train(:,k),C_train(:,k), 'Color', 'k')
title(title_string)
xlabel(['Known constituent (' ,ConcentrationUnits,')'])
ylabel(['Predicted constituent (' ,ConcentrationUnits,')'])
set(gca, 'FontSize',14)
box on
axis square
hold off
end
legend(h, 'Location', 'northwest')

```



Legend: Dot is predicted. Circle is calibration. Cross is cross-validation.')

Disclaimer

This material was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the United States Department of Energy, nor Battelle, nor any of their employees, nor any jurisdiction or organization that has cooperated in the development of these

materials, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness or any information, apparatus, product, software, or process disclosed, or represents that its use would not infringe privately owned rights.

Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or Battelle Memorial Institute. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof.

PACIFIC NORTHWEST NATIONAL LABORATORY

operated by

BATTELLE

for the

UNITED STATES DEPARTMENT OF ENERGY

under Contract DE-AC05-76RL01830