

**EFFECTS OF DIEL-CYCLING HYPOXIA ON THE CARDIAC
ACTIVITY AND GROWTH OF *ARGOPECTEN IRRADIANS***

A Thesis Presented

by

Samuel Jonathan Gurr

to

The Graduate School

in Partial Fulfillment of the

Requirements

for the Degree of

Master of Science

in

Marine Science

Stony Brook University

August 2017

ProQuest Number: 10615476

All rights reserved

INFORMATION TO ALL USERS

The quality of this reproduction is dependent upon the quality of the copy submitted.

In the unlikely event that the author did not send a complete manuscript and there are missing pages, these will be noted. Also, if material had to be removed, a note will indicate the deletion.



ProQuest 10615476

Published by ProQuest LLC (2017). Copyright of the Dissertation is held by the Author.

All rights reserved.

This work is protected against unauthorized copying under Title 17, United States Code
Microform Edition © ProQuest LLC.

ProQuest LLC.
789 East Eisenhower Parkway
P.O. Box 1346
Ann Arbor, MI 48106 – 1346

Stony Brook University
The Graduate School

Samuel Jonathan Gurr

We, the thesis committee for the above candidate for the
Master of Science in Marine Science degree, hereby recommend
acceptance of this thesis.

Nils Volkenborn, Ph.D.
Advisor
School of Marine and Atmospheric Science
Stony Brook University

Christopher J. Gobler, Ph.D.
Co-Advisor, Second Reader
School of Marine and Atmospheric Science
Stony Brook University

Robert Cerrato, Ph.D.
Committee Advisor, Third Reader
School of Marine and Atmospheric Science
Stony Brook University

This thesis is accepted by the Graduate School

Charles Tabor
Dean of Graduate School

Thesis
Abstract of the Thesis

**EFFECTS OF DIEL-CYCLING HYPOXIA ON THE CARDIAC
ACTIVITY AND GROWTH OF *ARGOPECTEN IRRADIANS***

by

Samuel Jonathan Gurr
Master of Science

in

Marine Science
Stony Brook University

2017

ABSTRACT

Bottom water oxygen concentrations in coastal environments can oscillate between fully oxygenated and hypoxic conditions on a daily basis. How benthic organisms deal with such drastic changes in oxygen availability is not well understood. Specifically, we do not know the magnitude, duration, and frequency at which diel-cycling hypoxia conditions become harmful. Here we have used non-invasive infrared sensors to measure the cardiac activity of the Atlantic bay scallops, *Argopecten irradians*, in response to diel-cycling hypoxia *in-situ* over one-month periods as well as in the laboratory in controlled incubations using animals conditioned to contrasting field conditions. In the field, heartbeat rates at a well-oxygenated site were relatively stable while scallops deployed at two sites with pronounced diel-cycling hypoxia were elevated and more variable. Maximal heartbeat rates during diel-cycling hypoxia were commonly recorded around dawn when oxygen concentrations fell to 5 mg L⁻¹ indicating a sub-lethal response to dissolved oxygen (DO) concentrations higher than what is typically defined as hypoxia. Laboratory incubations confirmed the tight link between DO and cardiac activity. An increase of heartbeat rate in response to an initial decline from fully oxygenated conditions was indicative of a regulatory response in which cardiac activity was enhanced presumably to maintain oxygen supply. At DO below 3 mg L⁻¹ heartbeat rates declined reaching a state of acardia during anoxia, suggesting a conformer response to severe hypoxia. *In-situ* and laboratory data was integrated into a

novel conceptual model to characterize four phases that interpret cardiac and respiratory activity in diel-cycling hypoxia. Heartbeat frequency was a suitable proxy for respiration under normoxia, but scallops were unable to compensate for reduced oxygen availability by increasing heartbeat rates below hypoxic thresholds. Pre-conditioning to diel-cycling hypoxia did not affect survival or cardiac activity in anoxic and severe hypoxic treatments. However, *A. irradians* pre-conditioned to diel-cycling hypoxia were less responsive to normoxia with heightened effort to maintain vital aerobic functions after long-term exposure to severe hypoxia. Survival after 12-14 hours of anoxia and mortality after 23-32 hours of anoxia convey physiological limitations unaffected by acclimation. We speculate that repetitive exposure to periods of DO oscillations with exposure below 5.0 mg L⁻¹ in the field can cause sub-lethal effects to *A. irradians* affecting fitness, growth, and reproductive success.

DEDICATION PAGE

I dedicate this Master's thesis to my mother and father. Their encouragement to pursue my interests and explore the Long Island Sound shoreline throughout my growing years has driven me to become the scientist I am today.

PREVIEW

TABLE OF CONTENTS

ABSTRACT	ii
DEDICATION PAGE.....	iv
LISTS OF TABLES AND FIGURES	vii
ACKNOWLEDGEMENTS	viii
1. INTRODUCTION.....	1
1.1. Hypoxia in coastal environments.....	1
1.2. Effects of hypoxia on benthic invertebrates.....	1
1.3. Cardiac activity as response variable.....	2
1.4. Model organism: <i>Argopecten irradians</i>	3
1.4. Working hypotheses and approaches.....	3
2. METHODS.....	4
2.1. Characterization of diel-cycling hypoxia.....	4
2.2. Heartbeat measurements with infrared sensors.....	5
2.3. Exp 1: Conditioning to diel-cycling hypoxia and effects on the ability to cope with anoxia.....	6
2.4. Exp 2: Conditioning to diel-cycling hypoxia and effects on the ability to cope with severe hypoxic events.....	7
2.5. Respiration measurements.....	8
2.6. Scallop shell growth, biovolume, tissue biomass, and condition index measurements.....	9
2.7. Data Analysis.....	10
3. RESULTS.....	11
3.1. Patterns of diel-cycling hypoxia in bays and harbors around Long Island.....	11
3.1.1 Exp.1: Late-summer deployments a two sites.....	11
3.1.2 Exp. 2: Early-summer deployments at nine sites.....	12
3.2. <i>In-situ</i> heartbeat monitoring.....	13
3.3. Cardiac activity and animal respiration.....	14
3.3.1 Response to temperature.....	14
3.3.2 DO dependence.....	14
3.4. Cardiac activity of acclimated scallops in response to oxygen decline and anoxia.....	15
3.5. Cardiac activity of acclimated scallops in response to severe hypoxia.....	16
3.6. Effects of diel-cycling hypoxia on scallop growth.....	16

4. DISCUSSION	17
4.1. Cardiac activity during diel-cycling hypoxia.....	17
4.2. Multi-stress environments and effects on scallop performance.....	20
4.3. Acclimation to diel-cycling hypoxia and effects on resilience to hypoxia.....	22
4.4. Conclusions and Future Directions.....	23
5. REFERENCES	25

PREVIEW

LIST OF TABLES

1. Descriptors of diel-cycling hypoxia and site characteristics.....	31
--	----

LIST OF FIGURES

1. <i>In-situ</i> cardiac activity at Nicoll Bay, Fire Island, and Seatuck.....	32
2. Percent time of hypoxic periods at nine sites.....	33
3. Box whisker plots of heartbeat rates and DO at Nicoll Bay and Fire Island	34
4. Heartbeat and respiration rate during laboratory temperature stress experiment.....	35
5. Cardiac and respiratory responses to temperature.....	36
6. Heartbeat efficiency and the cardiac and respiratory responses to DO decline.....	37
7. Cardiac activity under lethal duration of exposure to anoxic conditions.....	38
8. Cardiac activity during short-term anoxia and recovery.....	39
9. Cardiac response to an extended period of period of severe hypoxia.....	40
10. Cardiac recovery after an extended period of period of severe hypoxia.....	41
11. Shell growth and condition <i>A. irradians</i> and mean chlorophyll among the nine sites.....	42
12. Cardiac response to diel-cycling dissolved oxygen (CRD _{DO}); Model and <i>in-situ</i> example.....	43
13. CRD _{DO} with Nicoll Bay and Seatuck heartbeat, DO, and temperature data	44

APPENDIX

1. Strongest regression for shell growth and condition index with hypoxia descriptors.....	45
2. Shell growth and percent time of hypoxia under five DO thresholds.....	46
3. Condition index and percent time of hypoxia under five DO thresholds.....	47

ACKNOWLEDGMENTS

This study was partially supported by New York Sea Grant (Project number 1124181). We thank all members of the Long Island Water Quality Index program that provided the temperature, oxygen and *in-situ* site characteristics data for this study. We also thank Steven Tettelbach (LIU Post), Joe Hinton (Marine Field Technician), the Cornell Cooperative Extension Marine Program, Ian Dwyer (SBU), and Andrew Griffith (SBU) for providing specimen for this study. I also thank Robert Cerrato for his intellectual contributions. We also want to extend our gratitude to Fernando Lima for the development and implementation of the infrared heart rate sensor technology. I show the utmost gratitude for my lab partners Ian Dwyer and Molly Graffam; this research would not have been possible without them in addition to the combined assistance, generosity, and innovations from scientists and friends.

1. INTRODUCTION

1.1 Hypoxia in coastal environments

In coastal ecosystems hypoxia is a common physicochemical environmental stressor with an increasing global occurrence in frequency and severity due to nutrient loading and eutrophic conditions (Diaz and Rosenberg 2008). Periods of hypoxia or low dissolved (DO) develop when oxygen demand outweighs supply. Hypoxia intensifies seasonally from the effects of elevated temperature, light intensity, stratification and eutrophication (Diaz 2001; Howarth et al. 2011; Wallace et al. 2014). Large quantities of oxygen are consumed through bacterial degradation of algae resulting in hypoxic conditions within bottom waters that can last weeks to months. In shallow water environments, photosynthetic activity during the day and community respiration during night can lead to extremely variable oxygen availability (D'Avanzo and Kremer 1994). Diel-cycling hypoxia seasonally intensifies in summer and co-occurs with acidification among coastal environments (Wallace et al. 2014). Coupled stressors pose dynamic stress oscillations on seasonal as well as on daily time scales to organisms of commercial and ecological importance (Tyler et al. 2009; Baumann et al. 2014; Gobler and Baumann 2016; Gobler et al. 2017).

1.2 Effects of hypoxia on benthic invertebrates

Ecological issues and alterations of community structure associated with hypoxia are undisputed and result in mass mortalities, metabolic alterations, and escape responses of benthic and pelagic fauna (Howell and Simpson 1994; Diaz and Rosenberg 1995; Burnett and Stickle 2001). Such conditions may constitute a significant burden for benthic organisms affecting physiology, growth, and reproductive success (Baker and Mann 1992; Thomas et al. 2007; Cheung et al. 2008; Wu 2009; Gobler et al. 2014; Steckbauer et al. 2015; Keppel et al. 2016; Gobler et al. 2017), but sub-lethal effects of diel-cycling hypoxia on benthic organisms are not well understood. Importantly, many biological response variables (such as growth rates, development, mortality, etc.) integrate over relatively long time scales that limit the ability to understand and identify effects of dynamic environmental stressors and thresholds at which conditions become harmful.

1.3 Cardiac activity as response variable

Cardiac activity is a widely used proxy of whole animal metabolism (Helm and Trueman 1967) and can be measured with high temporal resolution to reveal immediate metabolic responses to changes in the environment (Burnett et al. 2013; Chapperon et al. 2016). Heartbeat rates of invertebrates exposed to environmental stress have been extensively studied (DeFur and Mangum 1979; Gainey and Shumway 1988; Braby and Somero 2006), but past efforts with minimally invasive techniques were mostly constrained to laboratory use (Aagaard et al. 1991; Rovero et al. 1999; Xing et al. 2016; Seabra et al. 2016). Maximum heartbeat rate at upper critical temperature (T_{cII}) and Arrhenius breakpoint temperature (ABT) identify thermal windows of optimal aerobic performance and acute physiological limitations for marine invertebrates (Fredrich and Pörtner 2000; Pörtner 2012; Xing et al. 2016; Chapperon et al. 2016; Tagliarolo and McQuaid 2016). For example, the European spider crab *Maja squinado* reaches maximum cardiac activity at 31.5°C (T_{cII}) followed by an increase in products of anaerobic metabolism at 33°C when heartbeat rates decline (Fredrich and Pörtner 2000). This provides evidence that heartbeat rate is closely linked with scope of aerobic performance and metabolic transitions. Parallel measurements of heartbeat and respiration rates as a function of temperature (Defur and Magnum 1979) indicate a strong relationship between both physiological parameters, but few studies have addressed the cardiac response of marine invertebrates to dynamically changing oxygen availability (Pörtner and Grieshaber 1993; Grieshaber et al. 1994; Aguirre-Velarde et al. 2016). Some marine invertebrates exhibit a short cardiac upswing under the initial DO decline from full oxygen saturation and drastically decrease heartbeat rate under hypoxia and anoxia exposure (Brand and Roberts 1973; Nicholson 2002). An abnormal decline, elevation, and cease of heartbeat rate are referred to as bradycardia, tachycardia, and acardia, respectively. Slight tachycardia during DO decline followed by bradycardia under severe oxygen limited conditions is evidence of an attempt to maintain aerobic function and transition to anaerobic metabolism (Bayne 1971; DeFur and Pease 1988). Here we have used non-invasive heartbeat sensors in an ecophysiological approach to measure cardiac activity *in-situ* during diel-cycling hypoxia and in the laboratory under hypoxia and anoxic simulations to increase an understanding of immediate physiological responses to stress conditions.