

A FULL-SEQUENCE QUANTITATIVE GAP BETWEEN THE CHROMATIC AND COCHROMATIC NUMBERS OF A RANDOM GRAPH

SAMUIL PETKOV

ABSTRACT. Let $G_n \sim G(n, 1/2)$. We prove, along the full sequence of integers n , that

$$\chi(G_n) - \zeta(G_n) = \Omega\left(\frac{n}{(\log n)^3}\right)$$

with probability tending to one, where χ and ζ denote the chromatic and cochromatic numbers. This resolves the question of Erdős and Gimbel asking whether this difference tends to infinity with high probability. The main difficulty is to obtain estimates that remain uniform when the natural class-size cutoff changes. We use a signed profile supported on four consecutive class sizes, an exact identity for the overlap of two such profiles, and a matching restriction that controls the remaining cycle-space sum. A bounded-differences argument then amplifies the resulting second-moment witness to a high-probability cocoloring.

INTRODUCTION

A *cocoloring* of a graph G is a partition of $V(G)$ into nonempty classes, each inducing either an edgeless graph or a complete graph. The least number of classes in such a partition is the *cochromatic number* $\zeta(G)$. Since an ordinary proper coloring is a cocoloring using only independent-set classes, one always has $\zeta(G) \leq \chi(G)$.

Write $[n] := \{1, \dots, n\}$, and let $G_n \sim G(n, 1/2)$ be the labeled random graph on $[n]$. All logarithms are natural unless a base is displayed. Erdős and Gimbel asked whether

$$\chi(G_n) - \zeta(G_n) \longrightarrow \infty$$

with high probability (Erdős and Gimbel, 1993, p. 263). The question was restated by Gimbel (Gimbel, 2016, Section 7.4) and is cataloged as Erdős Problem 625 (Bloom, 2026). We obtain the following quantitative full-sequence statement.

Main theorem. For $G_n \sim G(n, 1/2)$,

$$\mathbb{P}\left(\chi(G_n) - \zeta(G_n) \geq \frac{(\log 2)^2}{32} \log\left(\frac{200}{153}\right) \frac{n}{(\log n)^3}\right) \longrightarrow 1.$$

In particular, $\chi(G_n) - \zeta(G_n)$ tends to infinity with high probability along the full sequence of integers n .

The phrase *full sequence* is substantive. The natural class-size cutoff jumps whenever the independence-number phase crosses an integer. At such a jump, changing one admissible class size changes the optimizing profile and its first-moment root. An estimate proved only for a density-one set of phase values does not automatically extend to every integer n . All asymptotic estimates in this paper are therefore uniform up to both endpoints of every phase interval.

Unless stated otherwise, every $o(1)$ below is a deterministic error uniform over the complete phase interval. We keep the high-skeleton and residual attachment errors separate until the final second-moment ledger, which makes the uniformity and the absence of double counting explicit.

2020 *Mathematics Subject Classification.* Primary 05C80; Secondary 05C15, 60C05.

Key words and phrases. random graph, chromatic number, cochromatic number, second moment method, configuration model.

Relation to previous work. The first-order asymptotic for the chromatic number of a dense random graph was proved by Bollobás (Bollobás, 1988), following the early work of Grimmett and McDiarmid (Grimmett and McDiarmid, 1975); later refinements include McDiarmid (1990), Panagiotou and Steger (2009), and Heckel (2018). The cochromatic number belongs to the theory of generalized chromatic numbers associated with hereditary graph properties (Scheinerman, 1992; Bollobás and Thomason, 1995).

For the difference $\chi(G_n) - \zeta(G_n)$, Heckel and, independently, Steiner related divergence to the nonconcentration of the chromatic number (Heckel, 2024; Steiner, 2025). Heckel subsequently proved a near-linear lower bound for a phase-dependent set containing approximately 95% of the integers (Heckel, 2025). The unresolved part is precisely the exceptional phase. We retain the signed first-moment idea and rare-seed amplification from that work, but replace the pairwise sign estimate by an exact overlap identity. One four-size profile covers every phase; exact endpoint-table summation and residual cycle-space restriction make its second moment uniform.

Contributions and main ideas. The proof has four reusable components.

- (1) **Uniform root separation.** We compare the ordinary coloring root with the signed cocoloring root supported on four consecutive class sizes. The finite support comparison is exact before passing to the limiting Gaussian dual. We then compare the finite target with the phase-center target through one explicit, phase-uniform error. The resulting separation has order $n/(\log n)^3$.
- (2) **Exact sign compatibility.** For two signed witnesses, the sum over all compatible sign assignments factors into local cell rewards and the size of a binary cycle space. No sign correlation is estimated before this exact identity has been extracted.
- (3) **Large cells without chosen completions.** Cells above half the class-size cap form a matching. We sum the entire physical matching fiber, compare it with full containment, and apply the ambient falling-factorial bound once to the total deficit. The resulting reference weights are regrouped exactly by their endpoint tables.
- (4) **Residual restriction and amplification.** After the large cells have been exposed, deletion outside their matching is injective on even residual edge sets. This converts the residual cycle-space sum into a product bound. The high-skeleton and residual logarithmic costs are kept separate and added only after the exact conditional decomposition. Paley–Zygmund supplies a rare signed witness, and a one-sided bounded-differences argument amplifies it to a high-probability cocoloring.

The four consecutive sizes used here form a small fixed support that spans the entire phase interval and leaves a uniform entropy advantage over ordinary coloring. Here $\delta \in [0, 1]$ is the rounding phase and $D_4(\delta)$ is the finite-support entropy loss defined in Section 5. Lemma 5.1 proves the uniform certificate

$$\log 2 - D_4(\delta) > \log \left(\frac{200}{153} \right).$$

The later factor $1/8$ leaves a factor-four margin over the displayed coefficient $1/32$. Thus the finite support is not merely a technical truncation: it is the device that converts phase dependence into a full-sequence theorem.

Guide to the argument. The *location step* identifies two continuous first-moment roots. The signed witness is placed at a tangent-rounded midpoint, retaining half of their leading separation while satisfying the exact profile conservation laws. The phase, finite-dual, target-displacement, and slope errors are uniform in the complete phase.

The *existence step* studies the normalized second moment of this one signed profile. Partial diagonals are handled first. The remaining overlap tables are divided into canonical high cells and residual cells. Sections 8 and 9 keep these two accounts disjoint: no residual factor is paid in the high-cell estimate, and no high-cell multiplicity is paid again in the residual estimate. The logarithm of the final upper bound is explicitly

$$\Lambda_n = \Gamma_n^{\text{skel}} + \Gamma_n^{\text{att}} = o \left(\frac{n}{(\log n)^4} \right).$$

The *amplification step* converts this second-moment seed into a typical statement. The largest induced subgraph admitting the prescribed number of cocoloring classes is one-Lipschitz under vertex-block exposure. Its concentration, together with a simultaneous coloring bound for every leftover set, makes the amplification uniform in the seed exponent.

Section 1 fixes the phase notation. Sections 2–5 locate and compare the two roots. Section 6 proves the exact signed-overlap identity, and Section 7 treats partial diagonals. Sections 8 and 9 contain the large-cell and residual estimates and the explicit global error ledger. Section 10 proves the amplifier. The final section combines the amplified cocoloring event with the ordinary chromatic lower-tail event and applies the uniform entropy certificate. A concise note after the proof identifies the exact machine-checked theorem and its build.

NOTATION AND PROOF OBJECTS

We collect here only the conventions that are used throughout several later sections. More local notation is introduced where it first becomes necessary. This keeps the main argument self-contained without forcing the reader to retain a large dictionary before the proof begins.

Asymptotic conventions. An event holds *with high probability* if its probability tends to one as $n \rightarrow \infty$. Whenever a phase parameter is present, $o(1)$ denotes one deterministic error sequence that is uniform over the full phase interval, including sequences approaching either endpoint. For integers $x, r \geq 0$, we use the falling factorial

$$(x)_r = x(x-1) \cdots (x-r+1), \quad (x)_0 = 1,$$

and set $(x)_r = 0$ for $r > x$. A quotient involving falling factorials is used only after the denominator has been shown to be nonzero.

Signed witnesses and profiles. A *signed cocoloring witness* is a partition together with an *I*- or *K*-mark on each class. An *I*-marked class must be independent and a *K*-marked class must be complete. The marks are counting data; they do not define a new graph invariant. In the four-size profile every class has size at least two for sufficiently large n , so a realized class cannot satisfy both requirements. Forgetting the marks therefore recovers its cocoloring without multiplicity.

A *profile* is a finitely supported sequence $(k_s)_{s \geq 1}$ of nonnegative integers, where k_s is the number of classes of size s . It is feasible on n vertices when

$$\sum_s k_s = k, \quad \sum_s s k_s = n.$$

The ordinary first moment counts partitions into independent classes. The signed first moment counts the same partitions together with one *I/K* mark per class.

Overlap table and support graph. Let $(A_a)_a$ and $(B_b)_b$ be two ordered profile partitions. Their overlap table is

$$r_{ab} = |A_a \cap B_b|.$$

Its row and column sums are the class sizes of the two partitions. It may also be viewed as the cell-count table of a bipartite configuration model: every underlying vertex pairs one labeled row stub with one labeled column stub.

The *support graph* $H(r)$ is the simple bipartite graph whose vertices are the row and column classes incident to an edge and whose edge set is

$$E(H(r)) = \{(a, b) : r_{ab} \geq 2\}.$$

A subset of its edges is *even* if every vertex has even degree. Writing $c(H)$ for the number of connected components, we put

$$\beta(H) = |E(H)| - |V(H)| + c(H).$$

Then $\beta(H)$ is the dimension of the binary cycle space, so the number of even edge sets is $2^{\beta(H)}$. The threshold two is exact: a cell of size zero or one contains no edge internal to both partition classes, whereas a cell of size at least two forces the marks on its row and column classes to agree.

Three matching objects. The proof uses three distinct objects that should not be conflated.

- (1) The *configuration matching* is the perfect matching of all labeled row and column stubs induced by the common vertex set.
- (2) A *physical partial matching* is a set of selected stub pairs inside specified overlap cells.
- (3) A *block support* is a matching between row-class slots and column-class slots; its edges indicate the cells declared canonical and high.

The product formula for physical cell counts relies on the third object being a matching. Distinct selected cells then use disjoint row and column stub families.

High cells, deficits, and endpoint tables. Let U be the largest class size in the four-size profile and put $R_0 = \lfloor U/2 \rfloor$. A cell is *high* when its multiplicity exceeds R_0 . Two high cells cannot share a row or a column, so the high cells form a canonical block support. For a selected cell e , let s_e, t_e be its endpoint sizes and let j_e be its realized multiplicity. Set

$$m_e = \min\{s_e, t_e\}, \quad h_e = m_e - j_e.$$

Here m_e is the full-containment multiplicity, j_e is the realized multiplicity, and h_e is the deficit. Highness implies $2h_e < m_e$.

Section 8 uses the bijection $i \mapsto i + 2$ from $\{0, 1, 2, 3\}$ to the deficit labels $\{2, 3, 4, 5\}$. For $i \in \{0, 1, 2, 3\}$, define $u_i := \alpha - (i + 2)$ and $\kappa_i := k_{i+2}$. Thus u_i is the endpoint size and κ_i is the number of available blocks corresponding to deficit label $i + 2$. For a block support P , the endpoint table $L(P) = (L_{ij})$ records how many support edges join types i and j . A table is *feasible* when its type- i row sum and type- j column sum are at most κ_i and κ_j , respectively. Every block support has a feasible table, and every feasible table is attained by a block-slot matching.

Separation of accounts. The *bare high-skeleton weight* contains the incidence probability and local rewards of the exposed high cells. It contains neither the rewards of unexposed cells nor the residual binary cycle-space factor. Conditional on a high skeleton, the expectation of those remaining factors is its *residual attachment*. Sections 8 and 9 preserve this separation exactly: a factor paid in one account is never paid again in the other.

Every load-bearing estimate follows the same order. We first state the finite identity, then apply deterministic inequalities, then take phase asymptotics, and only at the end deduce a probability statement. This order is useful both for mathematical auditing and for the theorem-by-theorem formalization.

1. PHASE NOTATION AND ELEMENTARY ESTIMATES

For an integer s , let

$$\mu_s(v) = \binom{v}{s} 2^{-\binom{s}{2}}, \quad \mu_s = \mu_s(n). \quad (1.1)$$

We use the following standard inequalities in their stated forms.

1. For integers $m \geq 1$,

$$\log(m!) = m \log m - m + \frac{1}{2} \log(2\pi m) + O(1/m). \quad (1.2)$$

Consequently, with $0 \log 0 = 0$, one has $\log(m!) = m \log m - m + O(\log(m+1))$ uniformly for $m \in \mathbb{N}_0$, with an absolute implied constant.

2. If a random variable Y is a function of r independent blocks and changing one block changes Y by at most one, then

$$\mathbb{P}(|Y - \mathbb{E}[Y]| \geq t) \leq 2 \exp(-2t^2/r). \quad (1.3)$$

We will use the corresponding one-sided bounds as well.

3. If $Z \geq 0$ and $0 < \mathbb{E}[Z] < \infty$, then

$$\mathbb{P}(Z > 0) \geq \frac{\mathbb{E}[Z]^2}{\mathbb{E}[Z^2]}. \quad (1.4)$$

4. If $X \sim \text{Bin}(m, 1/2)$, then

$$\mathbb{P}(X \leq m/4) \leq e^{-m/16}. \quad (1.5)$$

Indeed, exponential Markov with $t = \log 3$ bounds the probability by $[3^{1/4}(2/3)]^m \leq e^{-m/16}$.

5. For every nonnegative random variable X and $a > 0$, $\mathbb{P}(X \geq a) \leq \mathbb{E}[X] / a$.

The first is Stirling's estimate, the second is McDiarmid's bounded-differences inequality, and the third is the zero-threshold case of Paley–Zygmund. The fourth is the only binomial-tail estimate used below; the fifth is Markov's inequality. The bounded-differences formulation is the one recorded by McDiarmid (1989, Theorem 3.1).

2. THE COMPLETE INDEPENDENCE-NUMBER PHASE

Put

$$q := \log 2, \quad L := \log n, \quad \ell := \log \log n, \quad C := 1 + \log q - q.$$

The usual independence-number center can then be written without a change of logarithm base as

$$\alpha_0 = 2 \log_2 n - 2 \log_2 \log_2 n + 2 \log_2(e/2) + 1 = \frac{2(L - \ell + C)}{q} + 1. \quad (2.1)$$

Let

$$\alpha := \lfloor \alpha_0 \rfloor, \quad \delta := \alpha_0 - \alpha, \quad b := 1 - \delta.$$

Thus $0 \leq \delta < 1$, $0 < b \leq 1$, and

$$\alpha = \frac{2(L - \ell + C)}{q} + b.$$

When emphasizing dependence on n , we write $\delta_n := \delta$. All estimates below are uniform for the closed parameter range $0 \leq \delta \leq 1$; this includes integer sequences approaching either endpoint of the actual half-open phase interval.

Lemma 2.1 (Uniform phase expansion and adjacent-size control). *There is a bounded continuous function $K: [0, 1] \rightarrow \mathbb{R}$ and an absolute constant $C_{\text{ph}} > 0$ such that, for all sufficiently large n , the quantity*

$$\varepsilon_n^{\text{ph}} := C_{\text{ph}} \frac{1 + \ell^2}{L} \longrightarrow 0$$

satisfies the following expansion at the induced phase $\delta = \alpha_0 - \lfloor \alpha_0 \rfloor$:

$$\log \mu_\alpha = \delta L + \left(\frac{2}{q} - \frac{1}{2} - \delta \right) \ell + K(\delta) + E_n(\delta), \quad |E_n(\delta)| \leq \varepsilon_n^{\text{ph}}. \quad (2.2)$$

The error bound is uniform over all induced phase values, and therefore also along sequences approaching either endpoint of $[0, 1]$. Moreover, uniformly over those phase values,

$$\begin{aligned} \log \mu_{\alpha+2} &= (\delta - 2)L + \left(\frac{2}{q} + \frac{3}{2} - \delta \right) \ell + O(1), \\ \mu_{\alpha+2} &\leq \exp\{-L + C_2 \ell\} = o(1) \end{aligned} \quad (2.3)$$

for an absolute C_2 , while

$$\mu_{\alpha-2} \geq c n^2 (\log n)^{2/q-5/2} \quad (2.4)$$

for an absolute $c > 0$ and all sufficiently large n .

Proof. Because $\alpha = O(L)$ uniformly in the phase, one has $\alpha < n/2$ for all sufficiently large n . For $0 \leq x \leq 1/2$, $-2x \leq \log(1-x) \leq -x$. Hence

$$\log(n)_\alpha = \alpha L + \sum_{j=0}^{\alpha-1} \log\left(1 - \frac{j}{n}\right) = \alpha L + R_{n,\alpha}, \quad |R_{n,\alpha}| \leq \frac{\alpha^2}{n}.$$

Using the uniform Stirling remainder

$$\log(\alpha!) = \alpha \log \alpha - \alpha + \frac{1}{2} \log(2\pi\alpha) + O(1/\alpha)$$

in

$$\log \mu_\alpha = \log(n)_\alpha - \log(\alpha!) - q \binom{\alpha}{2}$$

gives

$$\log \mu_\alpha = \alpha \left(L - \log \alpha + 1 - \frac{q}{2}(\alpha - 1) \right) - \frac{1}{2} \log(2\pi\alpha) + O(1/L). \quad (2.5)$$

All implied constants and remainder bounds here are uniform in δ .

Write

$$\alpha = \frac{2L}{q}(1 + x_n), \quad x_n := \frac{-\ell + C + qb/2}{L}.$$

Uniformly for the actual range $0 < b \leq 1$ (and on its closure), one has $|x_n| \leq 1/2$ for all sufficiently large n . The Taylor formula $\log(1 + x) = x + O(x^2)$ therefore gives

$$\log \alpha = \ell + q - \log q + \frac{-\ell + C + qb/2}{L} + O\left(\frac{\ell^2}{L^2}\right).$$

Since

$$\frac{q}{2}(\alpha - 1) = L - \ell + C - \frac{q\delta}{2},$$

we obtain the uniform expansion

$$L - \log \alpha + 1 - \frac{q}{2}(\alpha - 1) = \frac{q\delta}{2} + \frac{\ell - C - qb/2}{L} + O\left(\frac{\ell^2}{L^2}\right). \quad (2.6)$$

Multiplying by $\alpha = 2(L - \ell + C)/q + b$ and collecting terms yields

$$\begin{aligned} \alpha \left(L - \log \alpha + 1 - \frac{q}{2}(\alpha - 1) \right) &= \delta L + \left(\frac{2}{q} - \delta \right) \ell \\ &\quad + \delta C + \frac{q}{2}\delta(1 - \delta) - \frac{2C}{q} - (1 - \delta) + O\left(\frac{1 + \ell^2}{L}\right). \end{aligned}$$

Also

$$-\frac{1}{2} \log(2\pi\alpha) = -\frac{1}{2}\ell - \frac{1}{2} \log(2\pi) - \frac{q}{2} + \frac{1}{2} \log q + O(\ell/L).$$

Thus (2.2) holds with

$$\begin{aligned} K(\delta) &:= \delta C + \frac{q}{2}\delta(1 - \delta) - \frac{2C}{q} - (1 - \delta) \\ &\quad - \frac{1}{2} \log(2\pi) - \frac{q}{2} + \frac{1}{2} \log q. \end{aligned} \quad (2.7)$$

This function is continuous, hence bounded, on $[0, 1]$. The combined remainder in (2.5)–(2.7) has absolute value at most $C'(1 + \ell^2)/L$ for an absolute C' . Taking $C_{\text{ph}} \geq C'$ gives $|E_n(\delta)| \leq \varepsilon_n^{\text{ph}}$ in (2.2).

For integers $1 \leq s < n$, the adjacent-size ratios are exact:

$$\frac{\mu_{s+1}}{\mu_s} = \frac{n-s}{s+1} 2^{-s}, \quad \frac{\mu_{s-1}}{\mu_s} = \frac{s}{n-s+1} 2^{s-1}. \quad (2.8)$$

The displayed formula for α gives the exact phase representation

$$2^\alpha = \exp(q\alpha) = \exp(2C + qb) \frac{n^2}{L^2}.$$

Since $0 < b \leq 1$, the multiplicative factor $\exp(2C + qb)$ is bounded above and below by positive absolute constants. Therefore, for every fixed M , uniformly over integers s with $|s - \alpha| \leq M$,

$$\frac{\mu_{s+1}}{\mu_s} = \Theta(L/n), \quad \frac{\mu_{s-1}}{\mu_s} = \Theta(n/L).$$

In particular,

$$\mu_{\alpha+2} = \mu_\alpha \Theta(L^2/n^2), \quad \mu_{\alpha-2} = \mu_\alpha \Theta(n^2/L^2). \quad (2.8a)$$

Taking logarithms and using (2.2) proves the first line of (2.3). Since $\delta \leq 1$, its leading term is at most $-L$, and the coefficient of ℓ is uniformly bounded; this proves the second line.

For the lower adjacent size, (2.2) and (2.8a) give

$$\log \mu_{\alpha-2} = 2L + \left(\frac{2}{q} - \frac{5}{2}\right)\ell + \delta(L - \ell) + K(\delta) + O(1).$$

For large n , $L - \ell > 0$, while K is bounded below. Exponentiating gives (2.4), with one absolute constant c and one eventuality threshold valid for the complete phase. $\square$

Let $X_{\alpha+2}$ denote the number of independent sets of size $\alpha + 2$. Then $\mathbb{E}X_{\alpha+2} = \mu_{\alpha+2}$, and the event $\alpha(G_n) > \alpha + 1$ implies $X_{\alpha+2} \geq 1$. Consequently Markov's inequality and (2.3) give the deterministic cap error

$$\mathbb{P}(\alpha(G_n) > \alpha + 1) \leq \mu_{\alpha+2} =: \varepsilon_n^{\text{cap}} \longrightarrow 0 \quad (2.9)$$

uniformly along the full sequence of integers.

3. CONTINUOUS PROFILE ROOTS

For an integer class size $u \geq 1$, put

$$d_u := 2^{\binom{u}{2}} u!.$$

Write $u = \alpha - i$, so i is the deficit from the phase center α . We use

$$S_+ := \{-1, 0, 1, 2, \dots\}, \quad S_4 := \{2, 3, 4, 5\}. \quad (3.1)$$

At finite n , the unrestricted support is $S_+^{(n)} := \{-1, 0, \dots, \alpha - 1\}$: the lower endpoint comes from the cap event (2.9), and the upper endpoint from positivity of class sizes. Throughout every finite- n profile, maximization, or sum indexed by S_+ , this cutoff is understood.

For $i \in S_+^{(n)}$, define the curved score

$$h_n(i) := \begin{cases} -\frac{\log 2}{2} i^2 + \sum_{r=0}^{i-1} \log\left(1 - \frac{r}{\alpha}\right), & i \geq 0, \\ -\frac{\log 2}{2} + \log\left(\frac{\alpha}{\alpha+1}\right), & i = -1. \end{cases} \quad (3.1a)$$

For a support $S \in \{S_+^{(n)}, S_4\}$ and a nonnegative real profile $\mathbf{k} = (k_i)_{i \in S}$, put $k := \sum_{i \in S} k_i$. When $k > 0$, define

$$L_{\mathbf{k}} := n \log n - n + k - \sum_{i \in S} k_i \log(k_i d_{\alpha-i}), \quad (3.2)$$

with the convention $0 \log 0 = 0$, and let $L_S(n, k)$ be its maximum over real $k_i \geq 0$ satisfying

$$\sum_{i \in S} k_i = k, \quad \sum_{i \in S} (\alpha - i) k_i = n. \quad (3.3)$$

For k feasible for the support S under consideration, set

$$s := \frac{n}{k}, \quad T := \alpha - s = \alpha - \frac{n}{k}. \quad (3.4)$$

With $p_i := k_i/k$, the constraints give $T = \sum_{i \in S} i p_i$; thus T is the mean deficit of the normalized profile.

Lemma 3.1 (Uniform root, slope, and finite-dual package). *Let*

$$s_0 := \alpha_0 - 1 - \frac{2}{\log 2}, \quad T_0 := \alpha - s_0 = 1 + \frac{2}{\log 2} - \delta. \quad (3.5)$$

There is a fixed $A_ > 0$ such that, for $S \in \{S_+, S_4\}$ and $0 \leq c \leq \log 2$, the equation*

$$L_S(n, k) + ck = 0$$

has, for all sufficiently large n , a unique zero $r_{S,c}$ in the corridor

$$\left| \frac{n}{r_{S,c}} - s_0 \right| \leq A_* \frac{\log \log n}{\log n}.$$

Uniformly in S , c , and the phase,

$$\frac{n}{r_{S,c}} = s_0 + O\left(\frac{\log \log n}{\log n}\right), \quad \alpha - \frac{n}{r_{S,c}} = T_0 + O\left(\frac{\log \log n}{\log n}\right). \quad (3.6)$$

For every fixed $A > 0$, uniformly over $S \in \{S_+, S_4\}$, $0 \leq c \leq \log 2$, and k feasible for S with $|n/k - s_0| \leq A \log \log n / \log n$,

$$\frac{\partial}{\partial k} \{L_S(n, k) + ck\} = \frac{2}{\log 2} (\log n)^2 + O_A((\log n)(\log \log n)). \quad (3.7)$$

In particular, the deterministic normalized slope error

$$\varepsilon_{n,A}^{\text{slope}} := \max_{\substack{S \in \{S_+, S_4\} \\ 0 \leq c \leq \log 2}} \sup_{\substack{k \text{ feasible for } S \\ |n/k - s_0| \leq A \log \log n / \log n}} \left| \frac{1}{(\log n)^2} \frac{\partial}{\partial k} \{L_S(n, k) + ck\} - \frac{2}{\log 2} \right|$$

satisfies $\varepsilon_{n,A}^{\text{slope}} \rightarrow 0$.

For two supports $R, S \in \{S_+, S_4\}$ at a k feasible for both supports,

$$\frac{L_R(n, k) - L_S(n, k)}{k} = \mathcal{F}_{n,R}(T) - \mathcal{F}_{n,S}(T), \quad (3.8)$$

where

$$\mathcal{F}_{n,S}(T) := \max_{\substack{p_i \geq 0 \ (i \in S) \\ \sum_{i \in S} p_i = 1 \\ \sum_{i \in S} i p_i = T}} \left[- \sum_{i \in S} p_i \log p_i + \sum_{i \in S} p_i h_n(i) \right]. \quad (3.8a)$$

Fix the compact target interval

$$K_* := \left[\frac{2}{\log 2} - \frac{1}{10}, 1 + \frac{2}{\log 2} + \frac{1}{10} \right] \subset (2, 5). \quad (3.9)$$

Then, uniformly for $T \in K_$,*

$$\mathcal{F}_{n,S}(T) \longrightarrow \mathcal{F}_S(T) := \max_{\substack{p_i \geq 0 \ (i \in S) \\ \sum_{i \in S} p_i = 1 \\ \sum_{i \in S} i p_i = T}} \left[- \sum_{i \in S} p_i \log p_i - \frac{\log 2}{2} \sum_{i \in S} i^2 p_i \right]. \quad (3.9a)$$

If $\lambda_{n,S}(T)$ and $\lambda_S(T)$ are the finite and limiting effective tilts, then there is a fixed $\Lambda > 0$ such that

$$|\lambda_{n,S}(T)| + |\lambda_S(T)| \leq 2\Lambda$$

for all sufficiently large n , and the deterministic dual error

$$\varepsilon_n^{\text{dual}} := \max_{S \in \{S_+, S_4\}} \sup_{T \in K_*} (|\mathcal{F}_{n,S}(T) - \mathcal{F}_S(T)| + |\lambda_{n,S}(T) - \lambda_S(T)|)$$

satisfies $\varepsilon_n^{\text{dual}} \rightarrow 0$.

For S_4 , if $p_{n,i}(T)$ and $p_i(T)$ denote the finite and limiting optimizers, then

$$\sup_{T \in K_*} \max_{2 \leq i \leq 5} |p_{n,i}(T) - p_i(T)| \longrightarrow 0, \quad \inf_{T \in K_*} \min_{n \text{ large } 2 \leq i \leq 5} p_{n,i}(T) > 0. \quad (3.9b)$$

All error sequences and eventuality thresholds above are independent of δ .

Proof. Put $q := \log 2$, $L := \log n$, and $\ell := \log \log n$. Dividing (3.2) by $k = n/s$ and writing $p_i = k_i/k$ gives

$$\frac{L_S(n, n/s)}{n/s} = (s-1)L - s + 1 + \log s + \max_p \left[- \sum_i p_i \log p_i - \sum_i p_i \log d_{\alpha-i} \right].$$

There is an exact affine-plus-curved decomposition

$$-\log d_{\alpha-i} = A_n + B_n i + h_n(i), \quad (3.10)$$

where

$$A_n := -\log d_\alpha, \quad B_n := q\alpha - \frac{q}{2} + \log \alpha, \quad (3.11)$$

and h_n is the curved score defined in (3.1a). Direct subtraction verifies the displayed identity also at $i = -1$. Thus

$$h_n(i) = -\frac{q}{2}i^2 + O(i^2/\alpha) \quad \text{for every fixed } i, \quad h_n(i) \leq -\frac{q}{2}i^2 \quad (i \geq -1). \quad (3.12)$$

The first estimate is uniform on every fixed finite set of deficits; the second is global on the finite support.

FINITE DUALS AND OPTIMIZING TILTS.

For $S \in \{S_+, S_4\}$, define

$$Z_{n,S}(\lambda) := \sum_{i \in S} \exp\{\lambda i + h_n(i)\}, \quad M_{n,S}(\lambda) := \frac{d}{d\lambda} \log Z_{n,S}(\lambda),$$

with the cutoff $S_+^{(n)}$ in the finite unrestricted sum. Define the limiting functions by

$$Z_S(\lambda) := \sum_{i \in S} \exp\left(\lambda i - \frac{q}{2}i^2\right), \quad M_S(\lambda) := \frac{d}{d\lambda} \log Z_S(\lambda).$$

The target center satisfies

$$\frac{2}{q} \leq T_0 \leq 1 + \frac{2}{q}, \quad (3.13)$$

so it lies in the interior of K_* . The same is true of every target in K_* relative to both supports.

Choose $\Lambda > 0$ so that

$$M_S(-\Lambda) < \min K_* < \max K_* < M_S(\Lambda) \quad (S = S_+, S_4).$$

Such a choice exists because the limiting mean maps are strictly increasing and their endpoint limits bracket K_* . For $m = 0, 1, 2$ and $|\lambda| \leq \Lambda$, (3.12) gives the summable majorant

$$|i|^m \exp\left(\Lambda|i| - \frac{q}{2}i^2\right).$$

On every fixed finite set of indices, the weights converge by (3.12). The majorant makes the remaining Gaussian tail uniformly small, and the omitted tail beyond the finite cutoff $\alpha - 1$ is bounded by the same series. Consequently the zeroth, first, and second tilted moments converge uniformly on $[-\Lambda, \Lambda]$. In particular,

$$Z_{n,S} \rightarrow Z_S, \quad M_{n,S} \rightarrow M_S, \quad M'_{n,S} \rightarrow M'_S$$

uniformly there.

Now $M'_S(\lambda) = \text{Var}_{S,\lambda}(i) > 0$. By continuity and compactness,

$$v_* := \frac{1}{2} \min_{\substack{S \in \{S_+, S_4\} \\ |\lambda| \leq \Lambda}} M'_S(\lambda) > 0.$$

For all sufficiently large n , one has $M'_{n,S}(\lambda) \geq v_*$ on the same interval, and the finite mean maps also bracket K_* . Hence, for every $T \in K_*$, the equations

$$M_{n,S}(\lambda_{n,S}(T)) = T, \quad M_S(\lambda_S(T)) = T$$

have unique solutions in $[-\Lambda, \Lambda]$. The inverse mean maps are uniformly v_*^{-1} -Lipschitz, and therefore

$$\sup_{T \in K_*} |\lambda_{n,S}(T) - \lambda_S(T)| \leq v_*^{-1} \sup_{|\lambda| \leq \Lambda} |M_{n,S}(\lambda) - M_S(\lambda)| \longrightarrow 0.$$

The entropy functional in (3.8a) is strictly concave on its feasible simplex. Its unique optimizer is

$$p_{n,S,T}(i) = \frac{\exp\{\lambda_{n,S}(T)i + h_n(i)\}}{Z_{n,S}(\lambda_{n,S}(T))},$$

and the dual value is exactly

$$\mathcal{F}_{n,S}(T) = \log Z_{n,S}(\lambda_{n,S}(T)) - \lambda_{n,S}(T)T.$$

The corresponding formulas hold in the limit. Uniform convergence of the partition functions and tilts proves (3.9a) and $\varepsilon_n^{\text{dual}} \rightarrow 0$. On the finite support S_4 , the optimizer formula also gives uniform coordinatewise convergence. The limiting coordinates are positive and continuous in T on the compact set K_* ; this proves (3.9b).

EXACT CANCELLATION BETWEEN SUPPORTS.

Substituting (3.10) into the normalized exponent gives the exact identity

$$\begin{aligned} \Psi_{n,S}(s) &:= \frac{L_S(n, n/s)}{n/s} \\ &= (s-1)L - s + 1 + \log s + A_n + B_n(\alpha - s) + \mathcal{F}_{n,S}(\alpha - s). \end{aligned} \quad (3.14)$$

At a fixed feasible k , the values of s and $T = \alpha - s$ are the same for both supports. Every term in (3.14) except the final dual value is therefore common, which proves the exact difference identity (3.8). No limiting replacement is used in this cancellation.

THE VALUE AND DERIVATIVE AT THE PHASE CENTER.

At $s = s_0$, let $T_0 = \alpha - s_0$. Applying Stirling once to A_n in (3.14), and then simplifying the affine terms exactly, gives the scalar expression

$$\begin{aligned} \Psi_{n,S}(s_0) - \mathcal{F}_{n,S}(T_0) &= s_0 \left(L - \log \alpha - \frac{q}{2}s_0 + \frac{q}{2} \right) - L + T_0 + 1 + \log s_0 + \frac{q}{2}T_0^2 \\ &\quad - \frac{1}{2} \log(2\pi\alpha) + O(1/L). \end{aligned} \quad (3.15)$$

Equation (2.1) gives the exact identity

$$\frac{qs_0}{2} = L - \ell + \log q - q.$$

Since $\alpha = s_0 + T_0$ and T_0 is uniformly bounded,

$$\log \alpha = \log s_0 + \frac{T_0}{s_0} + O(L^{-2}), \quad \log s_0 = \ell + q - \log q + O(\ell/L).$$

Consequently

$$\begin{aligned} s_0 \left(L - \log \alpha - \frac{q}{2}s_0 + \frac{q}{2} \right) &= -T_0 + \frac{q}{2}s_0 + O(\ell) \\ &= -T_0 + L - \ell + \log q - q + O(\ell). \end{aligned}$$

Substitution in (3.15) cancels the terms L and T_0 and leaves $O(\ell)$ uniformly in the phase. The dual term in (3.14) is $O(1)$ uniformly on K_* , by the Gaussian majorant above. Therefore

$$\Psi_{n,S}(s_0) = O(\ell) \quad (3.16)$$

uniformly for both supports.

The phase formula for α and the expansion of $\log \alpha$ also give

$$B_n = 2L - \ell + O(1) \quad (3.17)$$

uniformly in δ . Strict concavity and the optimizer formula imply

$$\frac{d}{dT} \mathcal{F}_{n,S}(T) = -\lambda_{n,S}(T).$$

Differentiating (3.14), with $T = \alpha - s$, yields

$$\Psi'_{n,S}(s) = L - 1 + s^{-1} - B_n + \lambda_{n,S}(T) = -L + O_A(\ell) \quad (3.18)$$

uniformly whenever $|s - s_0| \leq A\ell/L$. Indeed, the image of this corridor under $T = \alpha - s$ lies inside K_* for all sufficiently large n , because K_* has a fixed margin around the complete range of T_0 .

EXISTENCE, UNIQUENESS, AND SLOPE OF THE ROOTS.

Set

$$\Psi_{n,S,c}(s) := \Psi_{n,S}(s) + c.$$

By (3.16), there is an absolute C_0 such that $|\Psi_{n,S,c}(s_0)| \leq C_0\ell$ for both supports and every $0 \leq c \leq q$. By (3.18), after one phase-independent eventuality threshold,

$$\Psi'_{n,S,c}(s) \leq -\frac{1}{2}L$$

throughout each fixed corridor. Choose $A_* > 2C_0$. Integration gives

$$\Psi_{n,S,c}\left(s_0 - A_*\frac{\ell}{L}\right) \geq \left(\frac{A_*}{2} - C_0\right)\ell > 0,$$

$$\Psi_{n,S,c}\left(s_0 + A_*\frac{\ell}{L}\right) \leq \left(C_0 - \frac{A_*}{2}\right)\ell < 0.$$

The function is strictly decreasing in s on this interval, so it has one and only one zero there. Since $k = n/s > 0$, this is equivalent to the unique root $r_{S,c}$ of $L_S(n, k) + ck = 0$, and proves (3.6).

Finally,

$$L_S(n, k) + ck = k\Psi_{n,S,c}(s), \quad \frac{ds}{dk} = -\frac{s}{k}.$$

Hence

$$\begin{aligned} \frac{\partial}{\partial k} \{L_S(n, k) + ck\} &= \Psi_{n,S,c}(s) - s\Psi'_{n,S}(s) \\ &= \frac{2}{q}L^2 + O_A(L\ell), \end{aligned} \quad (3.19)$$

because $s = (2/q)L + O_A(\ell)$, (3.18) holds, and $\Psi_{n,S,c}(s) = O_A(\ell)$ throughout the corridor. This proves (3.7) and $\varepsilon_{n,A}^{\text{slope}} \rightarrow 0$ with one eventuality threshold for the complete phase. $\square$

4. A UNIFORM LOWER LOCATION FOR χ

Let $r_+(n) = r_{S_+,0}$ be the unrestricted zero from Lemma 3.1, and put

$$k_\chi^- := \lfloor r_+(n) \rfloor - \lceil \log n \rceil. \quad (4.1)$$

This is a deterministic integer. Since $r_+(n) = \Theta(n/\log n)$ uniformly in the phase, one has $1 \leq k_\chi^- < n$ for all sufficiently large n .

For an integer profile $\mathbf{k} = (k_i)_{-1 \leq i \leq \alpha-1}$, let $X_{\mathbf{k}}$ be the number of unordered proper colorings having exactly k_i classes of size $\alpha - i$. Direct enumeration gives

$$\mathbb{E}[X_{\mathbf{k}}] = \frac{n!}{\prod_i ((\alpha - i)!)^{k_i} k_i!} 2^{-\sum_i k_i \binom{\alpha-i}{2}}. \quad (4.2)$$

For an integer k , let $\mathcal{K}_{n,k,\alpha+1}$ be the set of these profiles satisfying

$$\sum_i k_i = k, \quad \sum_i (\alpha - i) k_i = n,$$

and define

$$E_{n,k,\alpha+1} := \sum_{\mathbf{k} \in \mathcal{K}_{n,k,\alpha+1}} \mathbb{E}[X_{\mathbf{k}}].$$

Thus $E_{n,k,\alpha+1}$ is the expected number of unordered proper colorings with exactly k nonempty parts, each of size at most $\alpha + 1$.

The profile sum. There are $\alpha + 1 = O(\log n)$ profile coordinates, and each coordinate lies between zero and n . Hence

$$|\mathcal{K}_{n,k,\alpha+1}| \leq (n+1)^{\alpha+1} = \exp\{O((\log n)^2)\} \quad (4.3a)$$

uniformly in k and in the phase. Applying the uniform factorial estimate (1.2) to (4.2), with the convention $0 \log 0 = 0$, gives

$$\log \mathbb{E}[X_{\mathbf{k}}] \leq L_{\mathbf{k}} + O((\log n)^2).$$

The error is uniform even when some $k_i = 0$: only $O(\log n)$ factorials occur, and every one contributes $O(\log n)$. Maximizing over the feasible profile and then summing (4.3a) yields

$$\log E_{n,k,\alpha+1} \leq L_+(n, k) + O((\log n)^2). \quad (4.3)$$

Displacement below the root. Set

$$\Delta_n := r_+(n) - k_{\chi}^-.$$

The floor and ceiling in (4.1) give the deterministic bounds

$$\log n \leq \Delta_n < \log n + 2. \quad (4.3b)$$

We next verify that the entire interval $[k_{\chi}^-, r_+(n)]$ lies in the derivative corridor of Lemma 3.1. Write $s(k) = n/k$. Uniformly on this interval, $k = \Theta(n/\log n)$ and $s(k) = \Theta(\log n)$. Therefore

$$|s(k_{\chi}^-) - s(r_+)| \leq \sup_{k \in [k_{\chi}^-, r_+]} \frac{n}{k^2} \Delta_n = O\left(\frac{(\log n)^3}{n}\right) = o\left(\frac{\log \log n}{\log n}\right).$$

Thus, after one phase-independent threshold, every point of the interval lies inside the common corridor used in (3.7).

Equation (3.7), with $S = S_+$ and $c = 0$, consequently gives an absolute $c_* > 0$ such that

$$\frac{d}{dk} L_+(n, k) \geq c_*(\log n)^2 \quad (k_{\chi}^- \leq k \leq r_+) \quad (4.3c)$$

for all sufficiently large n , uniformly in the phase. Since $L_+(n, r_+) = 0$, the mean-value theorem and (4.3b) imply

$$L_+(n, k_{\chi}^-) \leq -c_*(\log n)^2 \Delta_n \leq -c_*(\log n)^3. \quad (4.4)$$

Combining (4.3) and (4.4), and decreasing the constant once, gives a deterministic sequence

$$\varepsilon_n^{\text{prof}} := \exp\{-c_{\chi}(\log n)^3\} \longrightarrow 0$$

such that

$$E_{n,k_{\chi}^-, \alpha+1} \leq \varepsilon_n^{\text{prof}}. \quad (4.4a)$$

Removing the size cap. Let

$$\mathcal{A}_n := \{\alpha(G_n) \leq \alpha + 1\}.$$

Equation (2.9) gives a deterministic phase-uniform sequence $\varepsilon_n^{\text{cap}} \rightarrow 0$ such that

$$\mathbb{P}(\mathcal{A}_n^c) \leq \varepsilon_n^{\text{cap}}.$$

On $\mathcal{A}_n$, every class of every proper coloring has size at most $\alpha + 1$.

Suppose that $\chi(G_n) \leq k_\chi^-$ and choose a proper coloring with $h \leq k_\chi^-$ nonempty classes. If $h < k_\chi^-$, then some class contains at least two vertices: otherwise $h = n$, contradicting $h < k_\chi^- < n$. Splitting such a class into two nonempty subsets preserves independence. Repeating this operation produces a proper coloring with exactly k_χ^- nonempty classes. The size cap is preserved under splitting. Therefore

$$\{\chi(G_n) \leq k_\chi^-\} \cap \mathcal{A}_n \subseteq \{\text{an exactly } k_\chi^-, (\alpha + 1)\text{-bounded coloring exists}\}.$$

Markov's inequality and (4.4a) now give

$$\begin{aligned} \mathbb{P}(\chi(G_n) \leq k_\chi^-) &\leq \mathbb{P}(\mathcal{A}_n^c) + E_{n, k_\chi^-, \alpha+1} \\ &\leq \varepsilon_n^{\text{cap}} + \varepsilon_n^{\text{prof}} \rightarrow 0. \end{aligned} \tag{4.5}$$

Equivalently,

$$\mathbb{P}(\chi(G_n) > k_\chi^-) \rightarrow 1. \tag{4.6}$$

The direction is strict, as required by the final event intersection.

Finally, (4.3b) also records the location precision:

$$|k_\chi^- - r_+(n)| < \log n + 2 = o\left(\frac{n}{(\log n)^3}\right). \tag{4.7}$$

Thus both the probability statement and the location error hold with one set of phase-independent thresholds along the full sequence of integers.

5. THE FOUR-SIZE SIGNED FIRST-MOMENT ADVANTAGE

Recall that a signed cocoloring witness is a profile partition with one independent-or-complete declaration on each class. It is realized when each class induces the declared graph. These declarations are auxiliary counting data, not a new graph invariant.

The four-size comparison has two steps. Restricting the deficits to $S_4 = \{2, 3, 4, 5\}$ must cost strictly less than $\log 2$ per part. The 2^k sign choices then convert this strict entropy margin into a macroscopic root separation. Lemma 5.1 proves the margin, and the slope estimate in Lemma 3.1 converts it into displacement.

For $S \in \{S_+, S_4\}$, introduce the tilted weights, partition function, and mean map

$$\begin{aligned} w_i(\lambda) &:= \exp\left(\lambda i - \frac{\log 2}{2} i^2\right), & Z_S(\lambda) &:= \sum_{i \in S} w_i(\lambda), \\ M_S(\lambda) &:= \frac{\sum_{i \in S} i w_i(\lambda)}{Z_S(\lambda)} = \frac{d}{d\lambda} \log Z_S(\lambda). \end{aligned}$$

The Gaussian factor makes both sums finite. Moreover, $M'_S(\lambda) = \text{Var}_{S, \lambda}(i) > 0$, so for every target mean T in the interior of the convex hull of S there is a unique tilt $\lambda_S(T)$ with $M_S(\lambda_S(T)) = T$. The optimizer in (3.9a) is therefore

$$p_i = \frac{w_i(\lambda_S(T))}{Z_S(\lambda_S(T))} = \frac{e^{\lambda_S(T)i - \log 2 i^2 / 2}}{\sum_{j \in S} e^{\lambda_S(T)j - \log 2 j^2 / 2}}. \tag{5.1}$$

The same calculation gives the dual representation

$$\mathcal{F}_S(T) = \inf_{\lambda \in \mathbb{R}} \{\log Z_S(\lambda) - \lambda T\} = \log Z_S(\lambda_S(T)) - \lambda_S(T)T.$$

Define

$$D_4(\delta) = \mathcal{F}_{S_+}(T_0) - \mathcal{F}_{S_4}(T_0), \quad \gamma_4 = \log \frac{200}{153}. \quad (5.2)$$

This certificate is sufficient both for the root separation and for the displayed numerical constant in the final theorem.

Lemma 5.1 (Uniform entropy certificate). *For every $0 \leq \delta \leq 1$,*

$$0 \leq D_4(\delta) < \log \frac{153}{100}, \quad \log 2 - D_4(\delta) > \gamma_4. \quad (5.3)$$

Proof. Fix $\delta \in [0, 1]$, put $T_0 = 1 + 2/\log 2 - \delta$, and abbreviate

$$\lambda_4 := \lambda_{S_4}(T_0).$$

We begin by bracketing this tilt. At $\lambda = 2\log 2$, set $j = i - 2$, a bijection from S_4 onto $\{0, 1, 2, 3\}$. Substituting $i = j + 2$ gives

$$\begin{aligned} w_{j+2}(2\log 2) &= \exp\left(2\log 2(j+2) - \frac{\log 2}{2}(j+2)^2\right) \\ &= \exp\left(2\log 2 - \frac{\log 2}{2}j^2\right) = 4 \cdot 2^{-j^2/2}. \end{aligned}$$

Consequently, reindexing both sums in the definition of the mean—and only reindexing them, without discarding any term—gives

$$\begin{aligned} Z_{S_4}(2\log 2) &= \sum_{i=2}^5 w_i(2\log 2) = 4 \sum_{j=0}^3 2^{-j^2/2}, \\ \sum_{i=2}^5 i w_i(2\log 2) &= 4 \sum_{j=0}^3 (j+2) 2^{-j^2/2} \\ &= 8 \sum_{j=0}^3 2^{-j^2/2} + 4 \sum_{j=0}^3 j 2^{-j^2/2}. \end{aligned}$$

Dividing the second equality by the first explains the additive constant 2 in the following formula:

$$\begin{aligned} M_{S_4}(2\log 2) &= 2 + \frac{\sum_{j=0}^3 j 2^{-j^2/2}}{\sum_{j=0}^3 2^{-j^2/2}} \\ &< 2 + \frac{\sum_{j \geq 0} j 2^{-j^2/2}}{\sum_{j \geq 0} 2^{-j^2/2}} < 2 + \frac{4}{5} < \frac{2}{\log 2} \leq T_0. \end{aligned} \quad (5.4)$$

We verify the first strict inequality in (5.4) directly. Put

$$A = \sum_{j=0}^3 2^{-j^2/2}, \quad B = \sum_{j=0}^3 j 2^{-j^2/2}, \quad C = \sum_{j=4}^{\infty} 2^{-j^2/2}, \quad D = \sum_{j=4}^{\infty} j 2^{-j^2/2}.$$

Here $B/A \leq 3$, while $D/C \geq 4$. Since $A, C > 0$,

$$\frac{B+D}{A+C} - \frac{B}{A} = \frac{AD-BC}{A(A+C)} > 0.$$

Thus adjoining the terms whose new indices satisfy $j \geq 4$ strictly increases the weighted mean. For the next inequality, put $a_j = j2^{-j^2/2}$. For $j \geq 4$,

$$\frac{a_{j+1}}{a_j} = \frac{j+1}{j} 2^{-(2j+1)/2} \leq \frac{1}{16},$$

and $a_4 = 1/64$. Hence

$$\sum_{j \geq 4} j2^{-j^2/2} \leq \frac{1/64}{1 - 1/16} = \frac{1}{60}.$$

Writing $b = 2^{-1/2}$, retaining $j = 0, 1, 2$ in the denominator and $j = 0, 1, 2, 3$ in the numerator gives

$$\frac{\sum_{j \geq 0} j2^{-j^2/2}}{\sum_{j \geq 0} 2^{-j^2/2}} < \frac{19b/16 + 1/2 + 1/60}{5/4 + b} < \frac{4}{5},$$

where the last inequality follows from $b < 71/100$. Finally, $2/3 < \log 2 < 5/7$ gives $2 + 4/5 < 2/\log 2$.

At $\lambda = 9\log 2/2$, put $t = 2^{-1/8}$. The identity

$$\frac{9\log 2}{2}i - \frac{\log 2}{2}i^2 = \frac{\log 2}{8}\{81 - (2i - 9)^2\}$$

gives

$$w_i(9\log 2/2) = 2^{81/8} t^{(2i-9)^2}.$$

Thus the four weights, in the order $i = 2, 3, 4, 5$, are proportional to

$$t^{25}, \quad t^9, \quad t, \quad t.$$

The numerator of $M_{S_4}(9\log 2/2) - 4$ is therefore

$$t - t^9 - 2t^{25} = t(1 - t^8 - 2t^{24}) = t/4 > 0, \quad (5.5)$$

and the denominator is positive. Hence this mean is greater than four, whereas $T_0 \leq 1 + 2/\log 2 < 4$. Since M_{S_4} is strictly increasing, the two mean comparisons give

$$2\log 2 < \lambda_4 < 9\log 2/2. \quad (5.6)$$

For a tilt λ , define the omitted low- and high-deficit ratios

$$L(\lambda) := \frac{\sum_{i=-1}^1 w_i(\lambda)}{Z_{S_4}(\lambda)}, \quad H(\lambda) := \frac{\sum_{i \geq 6} w_i(\lambda)}{Z_{S_4}(\lambda)}.$$

Let $M_A(\lambda)$ denote the mean of i under the weights $w_i(\lambda)$ restricted to an index set A . Differentiation gives

$$\frac{d}{d\lambda} \log L(\lambda) = M_{\{-1,0,1\}}(\lambda) - M_{S_4}(\lambda) \leq 1 - 2 = -1,$$

and

$$\frac{d}{d\lambda} \log H(\lambda) = M_{\{6,7,\dots\}}(\lambda) - M_{S_4}(\lambda) \geq 6 - 5 = 1.$$

Thus L is decreasing and H is increasing on $\mathbb{R}$. Direct summation and Gaussian-tail bounds give

$$L(2\log 2) < \frac{51}{100}, \quad H(3\log 2) < \frac{1}{50}, \quad L(3\log 2) < \frac{3}{25}, \quad H(9\log 2/2) < \frac{1}{4}. \quad (5.7)$$

Here are explicit checks. At $2\log 2$, with $b = 2^{-1/2}$,

$$L(2\log 2) = \frac{b + 1/4 + b/16}{1 + b + 1/4 + b/16} < 51/100.$$

At $\lambda = 3 \log 2$,

$$w_i(3 \log 2) = \exp\left((\log 2) \left(3i - \frac{i^2}{2}\right)\right) = 2^{9/2} 2^{-(i-3)^2/2}.$$

Set $b = 2^{-1/2}$. After canceling the common factor $2^{9/2}$, the weight at index i is $b^{(i-3)^2}$. Thus the four S_4 -weights, for $i = 2, 3, 4, 5$, are

$$b, \quad 1, \quad b, \quad b^4,$$

and

$$2^{-9/2} Z_{S_4}(3 \log 2) = 1 + 2b + b^4 = \frac{5}{4} + 2b.$$

For the omitted low indices $i = -1, 0, 1$, the weights are

$$b^{16}, \quad b^9, \quad b^4 = \frac{1}{256}, \quad \frac{b}{16}, \quad \frac{1}{4}.$$

For the omitted high indices $i \geq 6$, they are

$$b^9 + b^{16} + \sum_{q \geq 5} b^{q^2}.$$

The first term in the remaining sum is $b^{25} = b/4096$, and the ratio between successive terms is at most $b^{11} < 1/32$. Therefore

$$\sum_{q \geq 5} b^{q^2} \leq \frac{b^{25}}{1 - 1/32} < \frac{1}{3968}.$$

It follows that

$$L(3 \log 2) = \frac{1/256 + b/16 + 1/4}{5/4 + 2b} < \frac{3}{25},$$

and

$$H(3 \log 2) \leq \frac{b/16 + 1/256 + 1/3968}{5/4 + 2b} < \frac{1}{50},$$

where the final inequalities follow from $7/10 < b < 71/100$.

Finally, at $9 \log 2/2$ the normalized exponent is $(2i - 9)^2$. Dividing all weights by the common S_4 weight t , the high indices $i = 6, 7, 8$ contribute t^8, t^{24}, t^{48} . From $i = 8$ onward the ratio between successive terms is at most $1/16$, so the tail beginning with t^{48} is less than $1/60$. The normalized S_4 denominator is $2 + t^8 + t^{24}$. Hence $H(9 \log 2/2) < 1/4$ follows from

$$3(t^8 + t^{24}) + 4/60 < 2, \tag{5.8}$$

because $t^8 = 1/2$ and $t^{24} = 1/8$.

If $\lambda_4 \leq 3 \log 2$, monotonicity and (5.7) give $L(\lambda_4) + H(\lambda_4) < 53/100$; if $\lambda_4 \geq 3 \log 2$, they give $L(\lambda_4) + H(\lambda_4) < 37/100$. By the dual representation above,

$$\mathcal{F}_{S_4}(T_0) = \log Z_{S_4}(\lambda_4) - \lambda_4 T_0,$$

whereas evaluating the S_+ dual function at the same parameter gives

$$\mathcal{F}_{S_+}(T_0) \leq \log Z_{S_+}(\lambda_4) - \lambda_4 T_0.$$

Subtracting these two relations yields

$$\begin{aligned} D_4(\delta) &\leq \log \frac{Z_{S_+}(\lambda_4)}{Z_{S_4}(\lambda_4)} \\ &= \log(1 + L(\lambda_4) + H(\lambda_4)) < \log(153/100). \end{aligned}$$

Since S_4 is a subset of S_+ , the loss is nonnegative. Subtracting from $\log 2$ completes the proof. $\square$

The following 2^k gain is the one-partition identity isolated by Heckel (2025, Proposition 6). We repeat its short argument because it is an input to the new root calculation. For a fixed partition into k classes, assigning each class the declaration “independent” or “complete” multiplies its first moment by 2^k : every declaration prescribes all internal edge bits, and all 2^k declarations have probability $2^{-\sum \binom{u}{2}}$. The declarations are disjoint here because all four allowed class sizes tend to infinity (in particular, there are no singleton classes). Let $r_4^{\text{co}} = r_{S_4, \log 2}$ be the unique corridor zero of

$$L_{S_4}(n, k) + \log 2k. \quad (5.9)$$

Put

$$\Phi_n(k) := L_{S_4}(n, k) + (\log 2)k.$$

Then $\Phi_n(r_4^{\text{co}}) = 0$ by definition. To evaluate the same function at the unrestricted root, let

$$T_+(n) := \alpha - \frac{n}{r_+(n)}$$

and introduce the finite- n support loss

$$D_{4,n}(T) := \mathcal{F}_{n,S_+}(T) - \mathcal{F}_{n,S_4}(T).$$

Since $L_{S_+}(n, r_+) = 0$, the exact support-comparison identity (3.8) gives

$$\begin{aligned} \Phi_n(r_+) &= L_{S_4}(n, r_+) - L_{S_+}(n, r_+) + (\log 2)r_+ \\ &= r_+ \{\log 2 - D_{4,n}(T_+(n))\}. \end{aligned} \quad (5.9a)$$

This equality is finite and contains no limiting substitution.

We next compare the finite support loss at $T_+(n)$ with the limiting support loss at the phase-center target T_0 . Define the deterministic target displacement

$$\varepsilon_n^{\text{target}} := |T_+(n) - T_0|.$$

Equation (3.6) gives the equivalent phase-uniform statements

$$T_+(n) = T_0 + O\left(\frac{\log \log n}{\log n}\right), \quad \varepsilon_n^{\text{target}} = O\left(\frac{\log \log n}{\log n}\right) \longrightarrow 0. \quad (5.9b)$$

The common corridor lies inside the fixed compact interval K_* for all sufficiently large n .

The limiting dual value is differentiable, with

$$\mathcal{F}'_S(T) = -\lambda_S(T),$$

because the envelope theorem cancels the derivative of the optimizing tilt. Section 3 bounds the limiting tilts by one constant on K_* , so the two limiting dual values have a common Lipschitz constant C_{Lip} . The deterministic finite-dual error from Section 3 satisfies

$$\sup_{T \in K_*} |\mathcal{F}_{n,S}(T) - \mathcal{F}_S(T)| \leq \varepsilon_n^{\text{dual}} \quad (S = S_+, S_4).$$

Consequently the single theorem-facing root error

$$\omega_n^{\text{root}} := 2\varepsilon_n^{\text{dual}} + 2C_{\text{Lip}}\varepsilon_n^{\text{target}} \longrightarrow 0 \quad (5.9c)$$

is deterministic and phase-uniform. Indeed,

$$\begin{aligned} |D_{4,n}(T_+(n)) - D_4(\delta)| &\leq |\mathcal{F}_{n,S_+}(T_+) - \mathcal{F}_{S_+}(T_+)| \\ &\quad + |\mathcal{F}_{n,S_4}(T_+) - \mathcal{F}_{S_4}(T_+)| \\ &\quad + |\mathcal{F}_{S_+}(T_+) - \mathcal{F}_{S_+}(T_0)| \\ &\quad + |\mathcal{F}_{S_4}(T_+) - \mathcal{F}_{S_4}(T_0)| \\ &\leq \omega_n^{\text{root}}. \end{aligned}$$

Thus ω_n^{root} is one deterministic error sequence valid across the complete phase, including integer sequences approaching either endpoint.

Substitution into (5.9a) yields the theorem-facing estimate

$$L_{S_4}(n, r_+) + (\log 2)r_+ = r_+\{\log 2 - D_4(\delta) + O(\omega_n^{\text{root}})\} = r_+\{\log 2 - D_4(\delta) + o(1)\}. \quad (5.10)$$

Thus the $o(1)$ in (5.10) is not an independent asymptotic convention: it is controlled by the explicit sum of the finite-dual and target-displacement errors.

Finally, Lemma 5.1 gives $\log 2 - D_4(\delta) \geq \gamma_4 > 0$. Hence (5.10) implies $\Phi_n(r_+) > 0$ for all sufficiently large n , uniformly in the phase. The normalized slope error in (3.7) tends to zero, so the derivative of Φ_n is positive throughout the common root corridor after one phase-independent threshold. Since $\Phi_n(r_4^{\text{co}}) = 0$, this proves

$$r_4^{\text{co}} < r_+.$$

The interval used in the subsequent mean-value argument is therefore nonempty, with its orientation fixed. Both roots lie in the corridor (3.6), so the whole interval between them is inside a fixed corridor. The mean-value theorem therefore gives a point $\xi_n \in (r_4^{\text{co}}, r_+)$ such that

$$\Phi_n(r_+) - \Phi_n(r_4^{\text{co}}) = \Phi'_n(\xi_n)(r_+ - r_4^{\text{co}}).$$

Now $r_+ = (\log 2/2 + o(1))n/\log n$, and (3.7), uniformly at ξ_n , gives

$$\Phi'_n(\xi_n) = \frac{2}{\log 2}(\log n)^2 + O((\log n)(\log \log n)).$$

Substituting this and (5.10) into the preceding identity yields

$$r_+ - r_4^{\text{co}} = \left(\frac{(\log 2)^2}{4} \{\log 2 - D_4(\delta)\} + o(1) \right) \frac{n}{(\log n)^3}. \quad (5.11)$$

In particular, for all sufficiently large n ,

$$r_+ - r_4^{\text{co}} \geq \frac{(\log 2)^2 \gamma_4}{8} \frac{n}{(\log n)^3}. \quad (5.12)$$

Choose the midpoint integer

$$k_{\text{co}} = \left\lceil \frac{r_4^{\text{co}} + r_+}{2} \right\rceil. \quad (5.13)$$

We call the resulting profile the *exact midpoint profile*: its total number of parts is the rounded midpoint between the signed four-size root r_4^{co} and the unrestricted chromatic root r_+ . The type counts are chosen next so that both finite- n conservation constraints hold exactly.

At this exact k , let $p_i^{(n)}$ be the exact finite- n maximizer of L_{S_4} . Let $\lambda_{n,\text{def}}$ denote the Lagrange multiplier for the deficit-mean constraint, with the sign convention that its contribution is $e^{\lambda_{n,\text{def}} i}$. The Lagrange equations and the decomposition (3.10) then give

$$p_i^{(n)} \propto d_{\alpha-i}^{-1} e^{\lambda_{n,\text{def}} i} \propto e^{\widehat{\lambda}_n i + h_n(i)}, \quad \widehat{\lambda}_n := B_n + \lambda_{n,\text{def}}. \quad (5.14)$$

Since $e^{B_n i} e^{\lambda_{n,\text{def}} i} = e^{(B_n + \lambda_{n,\text{def}}) i} = e^{\widehat{\lambda}_n i}$, the quantity $\lambda_{n,\text{def}}$ is the multiplier in the exact finite- n coordinates, whereas $\widehat{\lambda}_n$ is the tilt in the limiting Gaussian coordinates. (If a size-coordinate multiplier τ_n is used instead, then $\lambda_{n,\text{def}} = -\tau_n$.) The target deficit mean is $\alpha - n/k_{\text{co}}$. Lemma 3.1 makes these proportions converge uniformly to (5.1), with $\widehat{\lambda}_n \rightarrow \lambda_{S_4}(T_0)$. Equation (5.6) also shows directly that the four limiting weights differ by at most a fixed factor. Thus

$$\min_{2 \leq i \leq 5} p_i^{(n)} \geq c_p > 0 \quad (5.15)$$

uniformly in the phase.

Round $k_{\text{co}} p_i^{(n)}$ to preliminary integers $\tilde{k}_i$. Define the signed constraint errors

$$e_0 = \sum_i \tilde{k}_i - k_{\text{co}}, \quad e_1 = \sum_i i \tilde{k}_i - (\alpha k_{\text{co}} - n),$$

so $e_0, e_1 = O(1)$, and add

$$\Delta k_2 = e_1 - 3e_0, \quad \Delta k_3 = 2e_0 - e_1. \quad (5.16)$$

Set $\Delta k_4 = \Delta k_5 = 0$, $k_i := \tilde{k}_i + \Delta k_i$ for $2 \leq i \leq 5$, and $\mathbf{k} := (k_2, k_3, k_4, k_5)$.

Indeed, $\Delta k_2 + \Delta k_3 = -e_0$ and $2\Delta k_2 + 3\Delta k_3 = -e_1$. This enforces both constraints exactly, changes only $O(1)$ counts, and preserves positivity by (5.15). Put $\mathbf{k}^* = (k_{\text{co}} p_i^{(n)})_{i=2}^5$ and let $\Delta^{\text{tot}} = \mathbf{k} - \mathbf{k}^*$ be the total rounding and correction displacement. It has bounded norm and satisfies both homogeneous constraint equations, because $\mathbf{k}^*$ and $\mathbf{k}$ obey the same two constraints. Thus it is tangent to the feasible affine plane. At the exact optimizer the linear term vanishes, while the Hessian of the entropy term is $-\text{diag}(1/k_i)$. Hence

$$L_{\mathbf{k}^* + \Delta^{\text{tot}}} - L_{\mathbf{k}^*} = O\left(\sum_i \frac{(\Delta_i^{\text{tot}})^2}{k_i^*}\right) = O(1/k_{\text{co}}).$$

The resulting exact integer vector satisfies

$$u_i = \alpha - i, \quad k_i = \Theta(n/\log n) \quad (2 \leq i \leq 5). \quad (5.17)$$

Applying (1.2) to the factorials in the exact signed first moment of this vector gives

$$\mathbb{E}[Z_{\mathbf{k}}^{\text{sgn}}] = 2^{k_{\text{co}}} \frac{n!}{\prod_i (u_i!)^{k_i} k_i!} 2^{-\sum_i k_i \binom{u_i}{2}}, \quad (5.18)$$

and

$$\log \mathbb{E}[Z_{\mathbf{k}}^{\text{sgn}}] = L_{S_4}(n, k_{\text{co}}) + \log 2k_{\text{co}} + O(\log n) \geq c_Z k_{\text{co}} \quad (5.19)$$

for a phase-independent $c_Z > 0$. Indeed, the midpoint is at distance at least $(r_+ - r_4^{\text{co}})/2 - O(1)$ to the right of r_4^{co} . Integrating the positive derivative (3.7) over that interval and using (5.12) gives the scale calculation

$$\frac{n}{(\log n)^3} \cdot (\log n)^2 \asymp \frac{n}{\log n} \asymp k_{\text{co}}.$$

Consequently,

$$L_{S_4}(n, k_{\text{co}}) + \log 2k_{\text{co}} \geq c_Z k_{\text{co}},$$

after decreasing c_Z if necessary. Every u_i tends to infinity, so the independent and complete declarations of one part are disjoint. Thus $Z_{\mathbf{k}}^{\text{sgn}} > 0$ implies an actual cocoloring with k_{co} parts.

Finally, (4.1), (5.12), and (5.13) give

$$k_{\chi}^- - k_{\text{co}} \geq \left(\frac{(\log 2)^2 \gamma_4}{16} - o(1) \right) \frac{n}{(\log n)^3}. \quad (5.20)$$

6. EXACT SIGNED SECOND-MOMENT REPRESENTATION

For two partitions sharing exactly ℓ whole classes, Heckel (2025, Proposition 6) bounded the number of joint sign declarations by $2^{2k-\ell}$, thereby transferring selected ordinary-coloring overlap bounds to cocolorings. Lemma 6.1 below does not import that inequality. It keeps the entire overlap matrix and counts the compatible signs exactly: compatibility is controlled by the components of the graph of cells of size at least two. Whole common classes appear as one special case of this component formula.

Temporarily label all slots within each of the four types. This multiplies the unordered witness by the deterministic factor $\prod_i k_i!$ and hence does not change its normalized second moment. Choose two independent uniform ordered partitions of the profile. If their slots are indexed by a and b , put

$$r_{ab} = |V_a \cap W_b|. \quad (6.1)$$

The overlap matrix has the exact law

$$p(r) = \frac{\prod_a s_a! \prod_b t_b!}{n! \prod_{a,b} r_{ab}!}, \quad (6.2)$$

where here both degree lists are the multiset given by (5.17). Equivalently, place s_a stubs at each row slot and t_b stubs at each column slot and take a uniform perfect matching of the two n -stub sets.

Let $H(r)$ be the simple bipartite graph whose edges are cells with $r_{ab} \geq 2$, omitting isolated slot vertices. Write

$$W = \sum_{a,b} \binom{r_{ab}}{2}, \quad \beta(H) = |E(H)| - |V(H)| + c(H). \quad (6.3)$$

Lemma 6.1 (Exact sign sum). *The normalized local factor of the pair of partitions is*

$$A_\zeta(r) = 2^{W+c(H)-|V(H)|} = \left(\prod_{a,b} g(r_{ab}) \right) 2^{\beta(H)}, \quad (6.4)$$

where

$$g(0) = g(1) = g(2) = 1, \quad g(x) = 2^{\binom{x}{2}-1} \quad (x \geq 3). \quad (6.5)$$

Consequently

$$\frac{\mathbb{E}[(Z_{\mathbf{k}}^{\text{sgn}})^2]}{\mathbb{E}[Z_{\mathbf{k}}^{\text{sgn}}]^2} = \sum_{r \in \mathcal{R}(\mathbf{s}, \mathbf{t})} p(r) A_\zeta(r). \quad (6.6)$$

where $\mathcal{R}(\mathbf{s}, \mathbf{t})$ is the finite set of nonnegative integer matrices with the prescribed row margins $\mathbf{s}$ and column margins $\mathbf{t}$.

Proof. Let $B = \sum_a \binom{s_a}{2}$, which is the number of internal edge bits prescribed by one partition. A row sign and a column sign are compatible on a cell of size at least two exactly when they agree. Therefore signs must be constant on every component of H . There are

$$2^{2k_{\text{co}} - |V(H)| + c(H)}$$

compatible pairs of sign assignments, including the free signs on slots isolated from H . For each compatible pair, W edge bits are prescribed twice, so the probability of all constraints is $2^{-(2B-W)}$. Division by the square of the one-partition signed probability $(2^{k_{\text{co}}} 2^{-B})^2$ gives

$$\frac{2^{2k_{\text{co}} - |V(H)| + c(H)} 2^{-(2B-W)}}{(2^{k_{\text{co}}} 2^{-B})^2} = 2^{W+c(H)-|V(H)|},$$

which is the first expression in (6.4).

Formula (6.4) separates two effects that will be estimated by different methods. The product of the $g(r_{ab})$ is purely local, with $g(0) = g(1) = g(2) = 1$; the factor $2^{\beta(H)}$ is topological and counts the independent cycle choices of the support graph.

For every support edge, split $\binom{r}{2}$ as $(\binom{r}{2} - 1) + 1$. Since $|E| - |V| + c = \beta$,

$$\prod_{a,b} g(r_{ab}) = 2^{W - |E(H)|},$$

and multiplying by $2^{\beta(H)}$ gives the second expression. Averaging over (6.2) proves (6.6). $\square$

The identity

$$2^{\beta(H)} = \#\{F \subseteq E(H) : \deg_F(v) \text{ is even for every } v\} \quad (6.7)$$

will be used repeatedly. It is the elementary fact that the even subgraphs form the binary cycle space of dimension $\beta(H)$.

We also need one exact configuration-model estimate.

Lemma 6.2 (Joint prescribed-cell bound). *In a bipartite configuration model of total degree m_0 , with row degrees d_a and column degrees d'_b , prescribe distinct cells D and demands $x_{ab} \geq 1$. Put*

$$x = \sum_D x_{ab}, \quad D_a = \sum_b x_{ab}, \quad D'_b = \sum_a x_{ab}.$$

In the feasible case

$$x \leq m_0, \quad D_a \leq d_a \quad (\forall a), \quad D'_b \leq d'_b \quad (\forall b),$$

we have

$$\mathbb{P}(r_{ab} \geq x_{ab} \text{ for every } ab \in D) \leq \frac{\prod_a (d_a)_{D_a} \prod_b (d'_b)_{D'_b}}{(m_0)_x \prod_{ab \in D} x_{ab}!}. \quad (6.8)$$

If any feasibility condition fails, the prescribed event is empty and is handled separately. In particular, in all cases with $m_0 > 0$, with $e := \exp(1)$ and $\theta_{ab} = e d_a d'_b / m_0$, its probability is at most

$$\prod_{ab \in D} \frac{\theta_{ab}^{x_{ab}}}{x_{ab}!}. \quad (6.9)$$

Proof. If a feasibility condition fails, no matching realizes all demands. In the feasible case, assigning disjoint row stubs to the demanded cells gives

$$\prod_a \frac{(d_a)_{D_a}}{\prod_b x_{ab}!}$$

choices. Independently selecting the column stubs gives

$$\prod_b \frac{(d'_b)_{D'_b}}{\prod_a x_{ab}!}$$

choices. For each cell ab , the selected row and column stubs can be paired in $x_{ab}!$ ways. Hence the product over all cells cancels one of the two copies of $\prod_{ab \in D} x_{ab}!$, and the total number of demand witnesses is

$$\frac{\prod_a (d_a)_{D_a} \prod_b (d'_b)_{D'_b}}{\prod_{ab \in D} x_{ab}!}.$$

Any fixed witness occurs with probability $1/(m_0)_x$, because after its x prescribed pairs are exposed the remaining matching is uniform on the unused stubs. A union bound proves (6.8).

It remains to simplify the global denominator. For $1 \leq x \leq m_0$,

$$\binom{m_0}{x} = \prod_{j=0}^{x-1} \frac{m_0 - j}{x - j} \geq \left(\frac{m_0}{x}\right)^x, \quad x! \geq \left(\frac{x}{e}\right)^x.$$

The factorial inequality follows, for example, from $\sum_{j=1}^x \log j \geq \int_1^x \log t \, dt = x \log x - x + 1$. Consequently

$$(m_0)_x = x! \binom{m_0}{x} \geq (m_0/e)^x, \quad (6.10)$$

while $(m_0)_0 = 1 = (m_0/e)^0$ covers $x = 0$. Combining (6.10) with $(d)_r \leq d^r$ proves (6.9). Notice that (6.8) retains the single global falling factorial before (6.9) is taken. $\square$

7. EXACT PARTIAL DIAGONALS

Purpose of this section. The exact overlap representation of Section 6 still permits two partitions to share whole classes. The definitions below assign a nonnegative marked weight to each such common subprofile. We prove that the total marked contribution is $1 + o(1)$, uniformly through the phase. This partial-diagonal estimate is the input used in Section 8 to control the endpoint-table sum. The counting identities in (7.1)–(7.4) are finite- n identities; asymptotic estimates are explicitly labeled below.

A *partial diagonal* is a specified collection of whole classes that the two partitions have in common. It is marked: one overlap may contain several such collections and then contributes to every corresponding nonnegative marked term. This intentional overcount is what makes the subsequent summation an upper-bound device.

For a selected common subprofile $\ell = (\ell_i)$, put

$$\ell_\bullet = \sum_i \ell_i, \quad m = \sum_i u_i \ell_i, \quad b_\ell = \sum_i \binom{u_i}{2} \ell_i. \quad (7.1)$$

Here $0 \leq \ell_i \leq k_i$ coordinatewise. Since the full profile has $\sum_i u_i k_i = n$, this automatically gives $m \leq n$; thus every factorial in the counting formula below has its ordinary combinatorial meaning.

Exact marked decomposition. The signed first moment of such partial partitions occupying any m vertices of $[n]$ is

$$Y_\ell^{\text{sgn}} = 2^{\ell_\bullet} \frac{n!}{(n-m)! \prod_i (u_i!)^{\ell_i} \ell_i!} 2^{-b_\ell}. \quad (7.2)$$

For each type i , choose the ℓ_i marked slots in the first partition and the ℓ_i marked slots in the second, giving $\binom{k_i}{\ell_i}^2$ choices. The common blocks themselves have the signed first-moment weight Y_ℓ^{sgn} . In the normalized second moment they are counted once rather than independently in each marginal, so their contribution is the reciprocal of this weight. Therefore the exact *marked* common-subprofile weight is

$$D(\ell) = \frac{\prod_i \binom{k_i}{\ell_i}^2}{Y_\ell^{\text{sgn}}}. \quad (7.3)$$

Different marked choices need not be disjoint: an overlap containing several common blocks contributes to each corresponding marked choice. The sums below are sums of these nonnegative marked weights, so no disjointness is used. In particular, $D(0) = 1$ and $D(k) = 1/\mathbb{E}[Z_k^{\text{sgn}}]$. To add one common block of type i , first note that

$$\frac{\binom{k_i}{\ell_i+1}^2}{\binom{k_i}{\ell_i}^2} = \frac{(k_i - \ell_i)^2}{(\ell_i + 1)^2},$$

while (7.2) gives

$$\frac{Y_{\ell+e_i}^{\text{sgn}}}{Y_\ell^{\text{sgn}}} = \frac{2(n-m)u_i}{u_i! (\ell_i + 1) 2^{\binom{u_i}{2}}} = \frac{2\mu_{u_i}(n-m)}{\ell_i + 1}.$$

Dividing these two ratios gives

$$\frac{D(\ell + e_i)}{D(\ell)} = \frac{(k_i - \ell_i)^2}{2(\ell_i + 1)\mu_{u_i}(n-m)}. \quad (7.4)$$

At the opposite endpoint, put $h = k - \ell$, $v = \sum_i u_i h_i$, and let $Z_h^{\text{sgn}}(v)$ be the signed first moment of a complete profile h on v vertices. Exact factorial cancellation gives

$$D(k - h) = \frac{B(h)}{\mathbb{E}[Z_k^{\text{sgn}}]}, \quad B(h) = \left(\prod_i \binom{k_i}{h_i} \right) Z_h^{\text{sgn}}(v), \quad (7.5)$$

and

$$\frac{\binom{k_i}{h_i+1}}{\binom{k_i}{h_i}} = \frac{k_i - h_i}{h_i + 1}, \quad \frac{Z_{h+e_i}^{\text{sgn}}(v + u_i)}{Z_h^{\text{sgn}}(v)} = \frac{2\mu_{u_i}(v + u_i)}{h_i + 1}.$$

Therefore

$$\frac{B(h + e_i)}{B(h)} = \frac{2(k_i - h_i)\mu_{u_i}(v + u_i)}{(h_i + 1)^2}. \quad (7.6)$$

Equations (7.2)–(7.6) are exact finite- n counting identities. We now use them only through uniform asymptotic bounds on the three ranges of selected mass.

In Heckel's admissible-phase argument, the corresponding control of partial profiles enters through the tame-coloring estimates of Heckel and Panagiotou (2023, Lemmas 5.1–5.3, 6.3–6.5, and 7.20), as used in Heckel (2025, Proposition 5); their hypotheses are the source of the lower μ_α -window in that theorem. The next lemma is the replacement needed here. It is proved directly for the four-size signed profile, including both corners and every intermediate mass, and no tame-profile theorem is invoked.

Lemma 7.1 (All common subprofiles). *For the exact midpoint profile,*

$$\sum_{0 \leq \ell_i \leq k_i} D(\ell) = 1 + o(1). \quad (7.7)$$

We split the common-subprofile sum according to the vertex mass occupied by the marked classes. The recurrence (7.4) controls the empty corner, the continuous rate function controls the central range, and the reverse recurrence (7.6) controls the full corner. For all sufficiently large n , when $\eta < 31/32$, the three ranges are the disjoint sets $m \leq \eta n$; $m > \eta n$ and $n - m > n/32$; and $n - m \leq n/32$. They are exhaustive, so no inclusion–exclusion or boundary convention is hidden in the assembly.

Proof. EMPTY CORNER.

Let

$$\eta = \frac{\log \log n}{32 \log n}, \quad \xi_i = \frac{k_i^2}{2\mu_{u_i}(n)}, \quad \Xi_{\text{empty}} = \sum_i \xi_i. \quad (7.8)$$

Equations (2.4), (2.8), and $k_i = \Theta(n/\log n)$ imply, with one phase-independent constant,

$$\Xi_{\text{empty}} = O\left((\log n)^{-(2/\log 2 - 1/2)}\right). \quad (7.9)$$

We now state explicitly how this activity at ambient size n controls every intermediate denominator in the selected-mass range.

Fix a final subprofile ℓ with selected mass $m \leq \eta n$, and expose its blocks in any order. At an intermediate selected mass $m' \leq m$, the exact ratio of first moments is

$$\frac{\mu_{u_i}(n)}{\mu_{u_i}(n - m')} = \frac{(n)_{u_i}}{(n - m')_{u_i}} = \prod_{r=0}^{u_i-1} \frac{n - r}{n - m' - r}.$$

Uniformly in the phase, $u_i = O(\log n)$ whereas $\eta n \gg \log n$. Hence, for all sufficiently large n , $m' + r \leq 2\eta n$ throughout this product, and therefore

$$\frac{\mu_{u_i}(n)}{\mu_{u_i}(n - m')} \leq (1 - 2\eta)^{-u_i}.$$

The phase expansion and $\log 2 > 2/3$ give $u_i \leq 3 \log n$ eventually. Since $\eta \leq 1/4$ and $-\log(1 - 2\eta) \leq 4\eta$,

$$(1 - 2\eta)^{-u_i} \leq \exp(4\eta u_i) \leq \exp\left(\frac{3}{8} \log \log n\right) = (\log n)^{3/8}. \quad (7.10)$$

Every threshold in this estimate is independent of the phase.

Suppose the current partial profile is a and the next selected block has type i . The exact recurrence (7.4), the inequality $(k_i - a_i)^2 \leq k_i^2$, and (7.10) give

$$\frac{D(a + e_i)}{D(a)} \leq \frac{(\log n)^{3/8} \xi_i}{a_i + 1}.$$

Multiplication along any exposure order gives the order-independent bound

$$D(\ell) \leq \prod_i \frac{((\log n)^{3/8} \xi_i)^{\ell_i}}{\ell_i!}.$$

All summands are nonnegative, so enlarging from the selected-mass range to the whole nonnegative four-dimensional lattice and expanding the exponential series yields

$$1 \leq \sum_{m \leq \eta n} D(\ell) \leq \exp\left((\log n)^{3/8} \Xi_{\text{empty}}\right). \quad (7.11)$$

The lower bound is the empty marked subprofile.

For clarity, the exponent tends to zero with a fixed power margin. Indeed, put $q = \log 2$. By (7.9),

$$(\log n)^{3/8} \Xi_{\text{empty}} = O\left((\log n)^{-(2/q - 7/8)}\right),$$

and $q < 7/10$ gives

$$\frac{2}{q} - \frac{7}{8} > \frac{20}{7} - \frac{7}{8} = \frac{111}{56} > 0.$$

Thus, for one deterministic sequence $\varepsilon_n^{\text{empty}} \rightarrow 0$ independent of the phase,

$$1 \leq \sum_{m \leq \eta n} D(\ell) \leq e^{\varepsilon_n^{\text{empty}}} = 1 + o(1).$$

CENTRAL RANGE.

Write

$$\begin{aligned} p_i &= k_i/k_{\text{co}}, \quad y_i = \ell_i/k_{\text{co}}, \quad Y = \sum_i y_i, \quad I = \sum_i i y_i, \\ z_i &= p_i - y_i, \quad R = \sum_i z_i = 1 - Y, \quad T := \sum_i i p_i = \alpha - \frac{n}{k_{\text{co}}}, \quad I_r = \sum_i i z_i = T - I. \end{aligned} \quad (7.12)$$

The residual vertex fraction is

$$\rho = \frac{n - m}{n} = R + \frac{I - TY}{\alpha - T}. \quad (7.13)$$

UNIFORM STIRLING EXTRACTION.

Before approximating any factorial, equations (7.2) and (7.3) give the exact identity

$$\begin{aligned} \log D(\ell) &= 2 \sum_i \{\log(k_i!) - \log(\ell_i!) - \log((k_i - \ell_i)!)\} \\ &\quad - \ell_{\bullet} \log 2 - \log(n!) + \log((n - m)!) + \sum_i \ell_i \log(u_i!) + \sum_i \log(\ell_i!) + (\log 2) b_{\ell}. \end{aligned}$$

We use the uniform form of Stirling's estimate

$$\log(t!) = t \log t - t + \sigma(t), \quad |\sigma(t)| \leq C_S \log(t + 2) \quad (t \in \mathbb{N}_0),$$

with the convention $0 \log 0 = 0$. Only a fixed number of factorials occur in the preceding identity, and every argument is at most n . Therefore the total remainder is $O(\log n)$ uniformly, including when some ℓ_i or $k_i - \ell_i$ is zero. Using $n = k_{\text{co}}(\alpha - T)$ gives

$$\begin{aligned} \log D(\ell) &= n\rho \log \rho \\ &+ k_{\text{co}} \sum_i \{2p_i \log p_i - 2(p_i - y_i) \log(p_i - y_i) - y_i \log y_i - y_i + y_i E_i\} + O(\log n), \end{aligned} \quad (7.14)$$

where

$$E_i = \log k_{\text{co}} + \log(u_i!) + u_i - u_i \log n + (\log 2) \binom{u_i}{2} - \log 2. \quad (7.15)$$

The entropy term in (7.14) is uniform even at the boundary of the coordinate box. By (5.15) and the bounded tangent-rounding displacement, the exact integer profile satisfies $p_i \geq c_p > 0$ after decreasing c_p once. Hence there is an absolute C_H such that, for $c_p \leq p \leq 1$ and $0 \leq y \leq p$,

$$2p \log p - 2(p - y) \log(p - y) - y \log y - y \leq C_H y \log\left(\frac{e}{y}\right). \quad (7.14a)$$

Indeed, when $y \leq p/2$, the mean-value theorem bounds $p \log p - (p - y) \log(p - y)$ by Cy , while $-y \log y$ supplies the only singular term. When $y \geq p/2$, one has $y \geq c_p/2$, and the assertion follows by compactness. Finally,

$$\sum_i y_i \log\left(\frac{e}{y_i}\right) \leq Y \log\left(\frac{4e}{Y}\right),$$

by the entropy bound on four coordinates. This is $O(Y \log(e/Y))$ uniformly.

Since $u_{i+1} = u_i - 1$, exact subtraction in (7.15) gives

$$\begin{aligned} E_{i+1} - E_i &= \log n - \log(\alpha - i) - (\log 2)(\alpha - i) + \log 2 - 1 \\ &= -\frac{\log 2}{2} \alpha + O(1), \end{aligned} \quad (7.16)$$

uniformly for $2 \leq i \leq 4$. For the second line, use $(\log 2)\alpha/2 = \log n - \log \log n + O(1)$ and $\log(\alpha - i) = \log \log n + O(1)$.

Let $\bar{E} = \sum_i p_i E_i$. Applying Stirling once to the complete signed first moment gives

$$-\frac{1}{k_{\text{co}}} \log \mathbb{E}[Z_{\mathbf{k}}^{\text{sgn}}] = \sum_i p_i \log p_i - 1 + \bar{E} + o(1). \quad (7.17)$$

The phase-uniform margin (5.19) implies $\bar{E} \leq C_E$. To expose the affine structure in (7.16), set

$$F_i := E_i + \frac{(\log 2)\alpha}{2} i.$$

Then $F_{i+1} - F_i = O(1)$. Because the support has only four consecutive indices,

$$F_i = \sum_j p_j F_j + O(1) = \bar{E} + \frac{(\log 2)\alpha}{2} T + O(1).$$

Consequently

$$E_i = \bar{E} - \frac{(\log 2)\alpha}{2} (i - T) + O(1),$$

and therefore

$$\sum_i y_i E_i \leq \frac{(\log 2)\alpha}{2} (TY - I) + O(Y). \quad (7.18)$$

Only the upper bound on $\bar{E}$ is used here.

It remains to compare the vertex fraction ρ with the profile fraction R . Since $2 \leq i, T \leq 5$,

$$|I - TY| = \left| \sum_i (i - T)y_i \right| \leq 3Y,$$

so (7.13) gives

$$|\rho - R| \leq \frac{3Y}{\alpha - T} = O(Y/\alpha).$$

If $\rho \geq 1/32$, then $Y \leq 1$ and the preceding bound implies $R \geq 1/64$ for all sufficiently large n . Hence $x \mapsto x \log x$ is uniformly Lipschitz on the interval between R and ρ . Moreover, $-R \log R \leq 1 - R = Y$. Thus

$$\begin{aligned} |n\rho \log \rho - k_{\text{co}}\alpha R \log R| \\ \leq k_{\text{co}}(\alpha - T)C|\rho - R| + k_{\text{co}}T|R \log R| \leq Ck_{\text{co}}Y, \end{aligned}$$

which proves

$$n\rho \log \rho = k_{\text{co}}\alpha R \log R + O(k_{\text{co}}Y). \quad (7.19)$$

Combining (7.14), (7.14a), (7.18), and (7.19) yields one absolute constant C such that

$$\log D(\ell) \leq k_{\text{co}}\alpha \Phi_T(z) + Ck_{\text{co}}Y \log\left(\frac{e}{Y}\right) + C \log n, \quad (7.20)$$

where

$$\Phi_T(z) = R \log R + \frac{\log 2}{2}(I_r - TR). \quad (7.21)$$

All constants and eventuality thresholds in this extraction are independent of the phase.

UNIFORM RATE NEGATIVITY.

The four-deficit geometry gives

$$I_r - TR \leq (5 - T)R, \quad I_r - TR = \sum_i (T - i)y_i \leq (T - 2)(1 - R). \quad (7.22)$$

Put $q = \log 2$ and recall that $Y = 1 - R$. Multiplying the first inequality in (7.22) by Y , the second by R , and adding gives

$$I_r - TR \leq 3RY. \quad (7.22a)$$

For $0 < R \leq 1$,

$$\log R \leq \frac{2(R - 1)}{R + 1}.$$

Indeed, for $x = (1 - R)/R \geq 0$, the function $\log(1 + x) - 2x/(2 + x)$ has derivative $x^2/((1 + x)(2 + x)^2) \geq 0$ and vanishes at zero. The positive expansion for $\log 2$ gives

$$\frac{2}{3} < q < \frac{7}{10}.$$

If $1/64 \leq R \leq 3/4$, then

$$\begin{aligned} \Phi_T &\leq -\frac{2R}{1 + R}Y + \frac{21}{20}RY \\ &= -\left(\frac{2}{1 + R} - \frac{21}{20}\right)RY \leq -\frac{13}{8960}Y. \end{aligned}$$

Here $2/(1 + R) \geq 8/7$ and $R \geq 1/64$. If $3/4 \leq R \leq 1$, then the phase corridor gives $T < 4$, so the second inequality in (7.22) implies $I_r - TR \leq 2Y$. Since $\log R \leq R - 1 = -Y$,

$$\Phi_T \leq -RY + qY = (q - R)Y \leq -\frac{1}{20}Y.$$

Thus, throughout the scalar range,

$$\Phi_T(z) \leq -\frac{Y}{5000}. \quad (7.23)$$

All rational inequalities in this two-range ledger are explicit; the logarithmic inequality and four-deficit reduction are proved above.

UNIFORM SUMMATION OVER THE CENTRAL RANGE.

Assume

$$m > \eta n, \quad n - m > n/32. \quad (7.24)$$

Then $\rho > 1/32$, and

$$\frac{m}{n} = \frac{\alpha Y - I}{\alpha - T} \leq \frac{\alpha Y}{\alpha - T}$$

implies

$$Y > \eta \frac{\alpha - T}{\alpha} \geq \frac{\eta}{2}$$

for all sufficiently large n . Define the deterministic error ratio

$$\varepsilon_n^{\text{diag}} := \frac{C \log(2e/\eta)}{\alpha} + \frac{2C \log n}{k_{\text{co}} \alpha \eta}. \quad (7.24a)$$

Since

$$\alpha = \Theta(\log n), \quad k_{\text{co}} = \Theta(n/\log n), \quad \eta = \Theta(\log \log n / \log n),$$

one has $\varepsilon_n^{\text{diag}} \rightarrow 0$, uniformly in the phase. Because $Y \geq \eta/2$, equations (7.20) and (7.23) give

$$\log D(\ell) \leq -k_{\text{co}} \alpha Y \left(\frac{1}{5000} - \varepsilon_n^{\text{diag}} \right).$$

For all sufficiently large n , the parenthesis is at least $1/10000$. Moreover, $\alpha \eta = \Theta(\log \log n)$. Hence there is a phase-independent $c > 0$ such that

$$D(\ell) \leq \exp\{-ck_{\text{co}} \log \log n\}. \quad (7.25)$$

Define

$$\varepsilon_n^{\text{central}} := (k_{\text{co}} + 1)^4 \exp\{-ck_{\text{co}} \log \log n\}.$$

There are at most $(k_{\text{co}} + 1)^4$ subprofiles, so the total central-range contribution is at most $\varepsilon_n^{\text{central}}$. Since $k_{\text{co}} \log \log n \gg \log n$, this deterministic sequence tends to zero, uniformly in the phase and with one eventuality threshold for the complete phase.

FULL CORNER.

Put $h = k - \ell$ and write

$$v(h) := \sum_i u_i h_i = n - m.$$

The full corner is the range

$$v(h) \leq \frac{n}{32}. \quad (7.26a)$$

We use the exact reverse recurrence (7.6) without replacing any ambient falling factorial.

First make the uniform smallness of the residual first moments explicit. For any $0 \leq w \leq n$ and every one of the four block sizes,

$$\frac{\mu_{u_i}(w)}{\mu_{u_i}(n)} = \frac{(w)_{u_i}}{(n)_{u_i}} \leq \left(\frac{w}{n}\right)^{u_i}.$$

Equations (2.2) and (2.8) give $\mu_{u_i}(n) \leq n^{6+o(1)}$ uniformly in the phase. Moreover, $u_i = (2/\log 2 + o(1)) \log n$, and $\log 32 = 5 \log 2$. Consequently, uniformly for $w \leq n/32$,

$$\mu_{u_i}(w) \leq n^{6+o(1)} 32^{-u_i} = n^{-4+o(1)}. \quad (7.26)$$

In particular, there is one phase-independent threshold after which

$$\mu_{u_i}(w) \leq n^{-3} \quad (w \leq n/32, \ 2 \leq i \leq 5). \quad (7.26b)$$

The relaxed exponent -3 leaves a fixed margin and is all that the recurrence requires.

Fix a final residual profile h satisfying (7.26a), and expose its blocks in any order. Let a be the current residual profile and $v(a) = \sum_i u_i a_i$. If the next block has type i , then $a_i < h_i$ and therefore

$$v(a) + u_i \leq v(h) \leq n/32.$$

The exact recurrence (7.6), $k_i - a_i \leq k_i \leq n$, and (7.26b) now give

$$\frac{B(a + e_i)}{B(a)} = \frac{2(k_i - a_i)\mu_{u_i}(v(a) + u_i)}{(a_i + 1)^2} \leq 2n^{-2} < 1$$

for all sufficiently large n . Hence B never increases along the path from 0 to h , and

$$B(h) \leq B(0) = 1. \quad (7.27a)$$

By the exact complementary identity (7.5) and the phase-uniform first-moment margin (5.19),

$$D(k - h) = \frac{B(h)}{\mathbb{E}[Z_k^{\text{sgn}}]} \leq e^{-c_Z k_{\text{co}}}. \quad (7.27)$$

There are at most $(k_{\text{co}} + 1)^4$ residual profiles. Therefore

$$\sum_{v(h) \leq n/32} D(k - h) \leq \exp\{4 \log(k_{\text{co}} + 1) - c_Z k_{\text{co}}\} = o(1)$$

uniformly in the phase. Equivalently, this sum is bounded by one deterministic sequence $\varepsilon_n^{\text{full}} \rightarrow 0$.

DISJOINT THREE-RANGE ASSEMBLY.

For sufficiently large n , one has $\eta < 31/32$. The coordinate box of all partial subprofiles is then the disjoint union of

$$\begin{aligned} \mathcal{E}_n &:= \{\ell : m \leq \eta n\}, \\ C_n &:= \{\ell : m > \eta n, n - m > n/32\}, \\ \mathcal{F}_n &:= \{\ell : n - m \leq n/32\}. \end{aligned}$$

Indeed, if $\ell \in \mathcal{F}_n$, then $m \geq 31n/32 > \eta n$, so the full corner cannot meet the empty corner; after excluding those two cases, the two strict inequalities defining C_n are automatic. Thus no boundary term is counted twice and no subprofile is omitted.

The three estimates proved above have the form

$$\begin{aligned} 1 &\leq \sum_{\ell \in \mathcal{E}_n} D(\ell) \leq e^{\varepsilon_n^{\text{empty}}}, \\ 0 &\leq \sum_{\ell \in C_n} D(\ell) \leq \varepsilon_n^{\text{central}}, \\ 0 &\leq \sum_{\ell \in \mathcal{F}_n} D(\ell) \leq \varepsilon_n^{\text{full}}, \end{aligned}$$

where every error sequence is deterministic, phase-independent, and tends to zero. The lower bound in the first line is the term $D(0) = 1$. Adding the three disjoint sums gives

$$1 \leq \sum_{0 \leq \ell_i \leq k_i} D(\ell) \leq e^{\varepsilon_n^{\text{empty}}} + \varepsilon_n^{\text{central}} + \varepsilon_n^{\text{full}} = 1 + o(1),$$

which proves (7.7) with one eventuality threshold for the complete phase. $\square$

8. CANONICAL HIGH CELLS AND ENDPOINT-TABLE COMPARISON

This section estimates the contribution of the large overlap cells. The finite combinatorics and the phase asymptotics are kept separate. We first sum the set of all labeled stub matchings that realize a fixed high-cell demand. We then compare realized multiplicities with full containment, sum all positive deficits, and regroup the full-containment references by endpoint table. Only in the final subsection do we insert the four-size phase estimates. No residual local reward or cycle-space factor is charged here; those factors belong to Section 9.

Section 7 has not subtracted or conditioned away overlaps containing common whole classes. Instead, its weights provide a normalized one-sided reference measure for the endpoint-table comparison below. Every overlap matrix remains present and is represented exactly once by its canonical high-cell skeleton.

8.1. All labeled realizations of a high skeleton. Let U be the largest class size in the fixed four-size profile and put

$$R_0 := \left\lfloor \frac{U}{2} \right\rfloor.$$

A cell is *high* if its multiplicity exceeds R_0 . Two high cells cannot share a row or a column: otherwise that row or column would contain more than U stubs. Hence all high cells form a canonical bipartite matching, denoted by P .

For $e \in P$, let s_e and t_e be its endpoint block sizes and set

$$m_e := \min\{s_e, t_e\}, \quad d_e := |s_e - t_e|, \quad h_e := m_e - j_e,$$

where j_e is the realized multiplicity. Thus m_e is the full-containment multiplicity, h_e is the deficit, and $j_e = m_e - h_e$. Since $j_e > U/2$ and $m_e \leq U$,

$$2h_e < m_e. \tag{1}$$

Put $J := \sum_{e \in P} j_e$.

For fixed (P, j) , let $\mathfrak{F}(P, j)$ be the finite set of labeled partial stub matchings that place exactly j_e pairs in every cell $e \in P$; no other stub pair is prescribed. We call $\mathfrak{F}(P, j)$ the *physical fiber* of the high skeleton. Thus the word “fiber” refers to the inverse image of (P, j) under the map that records only the selected cells and their multiplicities.

A size- j partial matching between labeled blocks of sizes s and t can be chosen in

$$\binom{s}{j} \binom{t}{j} j! = \frac{(s)_j (t)_j}{j!}$$

ways: choose its two endpoint sets and then the bijection between them.

Proposition 8.1 (Aggregate weight of a fixed high skeleton). *For a fixed matching support P and multiplicity vector $j = (j_e)_{e \in P}$, the sum of the bare weights over $\mathfrak{F}(P, j)$ is*

$$w(P, j) = \frac{\prod_{e \in P} (s_e)_{j_e} (t_e)_{j_e}}{(n)_J \prod_{e \in P} j_e!} \prod_{e \in P} g(j_e). \tag{2}$$

Proof. Because P is a matching, distinct selected cells use disjoint row and column stub sets. Therefore

$$|\mathfrak{F}(P, j)| = \prod_{e \in P} \frac{(s_e)_{j_e} (t_e)_{j_e}}{j_e!}.$$

A prescribed set of J stub pairs occurs with probability $(n)_J^{-1}$ in the uniform bipartite configuration matching. Multiplying this probability by the displayed cardinality and by the exposed reward $\prod_{e \in P} g(j_e)$ gives (2). $\square$

From now on we write

$$w_{\text{hi}}(P, j) := w(P, j).$$

This is the same bare high-skeleton weight that appears in the exact conditional decomposition of Section 9.

Remark 8.2. The matching hypothesis is load-bearing. If two selected cells shared a row or a column, their endpoint choices would compete for the same stubs and the one-cell fiber counts would not multiply.

8.2. Deficits and the single ambient loss. For endpoint sizes m and $m + d$, define

$$R_{m,d}(h) := \frac{\binom{m}{h}}{(d+1)(d+2)\cdots(d+h)} 2^{-hm+h(h+1)/2}, \quad R_{m,d}(0) := 1. \quad (3)$$

The product in the denominator is empty when $h = 0$.

Lemma 8.3 (Exact one-cell deficit ratio). *For a selected high cell with $h \geq 0$ and $2h < m$, the ratio of the aggregate one-cell factor at multiplicity $m - h$ to its full-containment value at multiplicity m is $R_{m,d}(h)$.*

Proof. The ratio of the physical matching counts is

$$\frac{\binom{m}{m-h} \binom{m+d}{m-h}}{(m-h)!} \frac{m!}{(m)_m (m+d)_m} = \frac{\binom{m}{h}}{(d+1)(d+2)\cdots(d+h)}.$$

The local signed rewards satisfy

$$\frac{g(m-h)}{g(m)} = 2^{-hm+h(h+1)/2}.$$

Multiplying the two identities proves the claim. $\square$

Write

$$w_{\text{full}}(P) := w(P, (m_e)_{e \in P}), \quad J_* := \sum_{e \in P} m_e, \quad H := J_* - J = \sum_{e \in P} h_e.$$

After the one-cell ratios have been extracted, the only factor that still couples different cells is the ambient denominator:

$$\frac{(n)_{J_*}}{(n)_J} = (n - J)_H \leq n^H. \quad (4)$$

Lemma 8.4 (Aggregate deficit comparison). *For every admissible deficit vector $h = (h_e)_{e \in P}$,*

$$w(P, m - h) \leq w_{\text{full}}(P) \prod_{e \in P} n^{h_e} R_{m_e, d_e}(h_e). \quad (5)$$

Proof. Apply Lemma 8.3 in each selected cell, apply (4) once, and use $n^H = \prod_{e \in P} n^{h_e}$. $\square$

The order of operations matters: first sum the physical fiber, and then apply the single aggregate bound $(n - J)_H \leq n^H$ from (4).

8.3. Summing all positive deficits. For a selected cell e , enlarge the exact positive-deficit range to

$$A_e := \{h \in \mathbb{N} : h \geq 1, 2h < m_e\}.$$

All weights are nonnegative, so the enlargement preserves the direction of the upper bound.

Lemma 8.5 (Optional-choice product). *Let $u_e(h) \geq 0$ for $e \in P$ and $h \in A_e$. Then*

$$\sum_{\omega} \prod_{e \in P} u_e(\omega_e) = \prod_{e \in P} \left(1 + \sum_{h \in A_e} u_e(h) \right),$$

where each ω_e is either the zero-deficit choice, of weight one, or one positive deficit $h \in A_e$.

Proof. Expand the finite product. Each monomial records exactly one independent choice in every selected cell, and every optional-choice vector occurs once. $\square$

For $2h < m$, integrality gives $2h + 1 \leq m$. Hence

$$\frac{h+1}{2} \leq \frac{m+1}{4} \leq m - \left\lfloor \frac{3m-1}{4} \right\rfloor.$$

Multiplying by h yields

$$h \left\lfloor \frac{3m-1}{4} \right\rfloor \leq hm - \frac{h(h+1)}{2}. \quad (6)$$

Together with $\binom{m}{h} \leq m^h$ and (3), this gives

$$n^h R_{m,d}(h) \leq \left(\frac{nm}{2^{\lfloor (3m-1)/4 \rfloor}} \right)^h. \quad (7)$$

For the remainder of this section, let $q = i + 2$. The map $i \mapsto q$ is a bijection from $\{0, 1, 2, 3\}$ to the deficit labels $\{2, 3, 4, 5\}$ used in Sections 5–7, with inverse $q \mapsto q - 2$. Define

$$\kappa_i := k_{i+2}, \quad u_i := \alpha - 2 - i, \quad 0 \leq i \leq 3.$$

Thus κ_i is the number of endpoint blocks of size u_i . For $i, j \in \{0, 1, 2, 3\}$ let $m_{ij} := \min\{u_i, u_j\}$ and put

$$\rho_{ij} := \frac{nm_{ij}}{2^{\lfloor (3m_{ij}-1)/4 \rfloor}}, \quad \rho_{16} := \sum_{i,j=0}^3 \rho_{ij}.$$

Every local base in (7) is at most ρ_{16} .

Lemma 8.6 (Coarse local deficit sum). *Suppose $\rho_{16} \leq 1$. Then, for every selected cell e ,*

$$\sum_{h \in A_e} n^h R_{m_e, d_e}(h) \leq (\alpha + 1) \rho_{16}.$$

Proof. If e has endpoint type (i, j) , then (7) bounds its term of deficit h by ρ_{ij}^h . Since $h \geq 1$ and $\rho_{ij} \leq \rho_{16} \leq 1$, this is at most ρ_{16} . Moreover, $|A_e| \leq m_e \leq \alpha + 1$. Summation proves the bound. $\square$

Proposition 8.7 (Fixed-support all-deficit bound). *Write $\mathbf{m} = (m_e)_{e \in P}$. If $\rho_{16} \leq 1$, then*

$$\sum_{\substack{h_e \in \mathbb{N}_0 \ (e \in P) \\ m_e - h_e > R_0 \text{ for every } e}} w(P, \mathbf{m} - h) \leq w_{\text{full}}(P) (1 + (\alpha + 1) \rho_{16})^{|P|}. \quad (8)$$

Proof. Use Lemma 8.4, enlarge the deficit range to A_e , and apply Lemma 8.5 and Lemma 8.6. $\square$

Projection onto the row endpoint is injective on a block matching. Therefore, if $K = \sum_{i=0}^3 \kappa_i$ is the number of row-class slots, then

$$|P| \leq K. \quad (9)$$

8.4. Regrouping by endpoint table. For $L = (\ell_{ij}) \in \mathbb{N}^{4 \times 4}$, write

$$r_i = \sum_j \ell_{ij}, \quad c_j = \sum_i \ell_{ij}.$$

Call L *feasible* if $r_i \leq \kappa_i$ and $c_j \leq \kappa_j$ for every i, j . Such a table is realized by choosing the indicated row and column block slots and pairing them cell by cell; conversely, every block matching has a feasible table. For a feasible endpoint table L , define

$$W(L) := \sum_{P: L(P)=L} w_{\text{full}}(P).$$

Equivalently, $W(L)$ is the sum of the full-containment atoms over all block-slot matching supports with table L .

Proposition 8.8 (Reference grouping). *The full-containment references satisfy*

$$\sum_P w_{\text{full}}(P) = \sum_{L \text{ feasible}} W(L). \quad (10)$$

More generally, for every nonnegative function F of the endpoint table,

$$\sum_P w_{\text{full}}(P) F(L(P)) = \sum_{L \text{ feasible}} W(L) F(L). \quad (11)$$

Proof. Partition the finite set of block-slot matching supports by the value of $L(P)$. On each fiber the factor $F(L(P))$ is constant. Summing first inside each fiber gives $W(L)$, and then summing over the feasible tables gives the two identities. $\square$

Whenever $\rho_{16} \leq 1$, combining Proposition 8.7, (9), and (10) yields

$$\sum_{\text{high skeletons}} w_{\text{hi}} \leq \left(\sum_{L \text{ feasible}} W(L) \right) (1 + (\alpha + 1)\rho_{16})^K. \quad (12)$$

Remark 8.9 (Sharper table-preserving form). If every $\rho_{ij} \leq 1/2$, then $\sum_{h \geq 1} \rho_{ij}^h \leq 2\rho_{ij}$. Applying (11) with $F(L) = \prod_{i,j} (1 + 2\rho_{ij})^{L_{ij}}$ gives

$$\sum_{\text{high skeletons}} w_{\text{hi}} \leq \sum_{L \text{ feasible}} W(L) \prod_{i,j} (1 + 2\rho_{ij})^{L_{ij}}.$$

The proof below uses the coarser form (12), which requires only the single condition $\rho_{16} \leq 1$ and cleanly separates the deficit sum from endpoint tables.

Remark 8.10 (Reusable finite core). The aggregate-fiber formula, the single ambient-denominator comparison, the optional-choice identity, and the endpoint-table regrouping are not specific to four endpoint sizes. They apply to any finite endpoint alphabet for which the selected cells form a matching. The four consecutive sizes enter only in the numerical deficit charge and the endpoint-table estimate below.

8.5. Endpoint-table comparison. Recall that $u_i = a - i$ and $\kappa_i = k_{i+2}$ for $0 \leq i \leq 3$, where $a = \alpha - 2$ and the corresponding deficit label is $i + 2$. For a full-containment table $L = (\ell_{ij})$, set

$$r_i = \sum_j \ell_{ij}, \quad c_j = \sum_i \ell_{ij}, \quad x_{ij} = \min\{u_i, u_j\}, \quad J(L) = \sum_{i,j} x_{ij} \ell_{ij}.$$

Its exact reference weight is

$$W(L) = \frac{\prod_i (\kappa_i)_{r_i} \prod_j (\kappa_j)_{c_j}}{\prod_{i,j} \ell_{ij}!} \frac{1}{(n)_{J(L)}} \prod_{i,j} \left[\frac{(u_i)_{x_{ij}} (u_j)_{x_{ij}}}{x_{ij}!} g(x_{ij}) \right]^{\ell_{ij}}. \quad (13)$$

For a one-sided margin vector v satisfying $0 \leq v_i \leq \kappa_i$, define

$$D(v) = \frac{\prod_i (\kappa_i)_{v_i}^2 \prod_i [u_i! g(u_i)]^{v_i}}{\prod_i v_i! (n)_{m(v)}}, \quad m(v) = \sum_i u_i v_i. \quad (14)$$

For comparison with Section 7, denote its common-subprofile vector by $\ell^{\text{pd}} = (\ell_q^{\text{pd}})_{q=2}^5$. Under the bijection $\ell_{i+2}^{\text{pd}} = v_i$, the multiplicity and block size are $k_{i+2} = \kappa_i$ and $\alpha - (i + 2) = u_i$; hence (7.3) gives $D(\ell^{\text{pd}}) = D(v)$. Below, $D(r)$ and $D(c)$ evaluate (14) on the margins in $0 \leq v_i \leq \kappa_i$.

For the endpoint pair (i, j) , put

$$s_{ij} := \min\{u_i, u_j\}, \quad t_{ij} := \max\{u_i, u_j\}, \quad d_{ij} := t_{ij} - s_{ij},$$

and define

$$Q_{ij} := (n + 1)^{d_{ij}/2} \frac{\sqrt{(t_{ij})^{d_{ij}}}}{d_{ij}!} 2^{-\{d_{ij}s_{ij} + \binom{d_{ij}}{2}\}/2}.$$

Thus $Q_{ii} = 1$. Since $t_{ij} \leq a$, $s_{ij} \geq a - 3$, and $d_{ij} \leq 3$,

$$Q_{ij} \leq \frac{(\tau_n^{\text{end}})^{d_{ij}}}{d_{ij}!}, \quad \tau_n^{\text{end}} = 2^{3/2} \sqrt{\frac{(n+1)a}{2a}} = O\left(\frac{(\log n)^{3/2}}{\sqrt{n}}\right). \quad (15)$$

Let

$$A_L = \frac{\prod_i r_i!}{\prod_{i,j} \ell_{ij}!}, \quad C_L = \frac{\prod_j c_j!}{\prod_{i,j} \ell_{ij}!}, \quad Q^L = \prod_{i,j} Q_{ij}^{\ell_{ij}}.$$

Lemma 8.11 (Endpoint-table product comparison). *For every feasible endpoint table L ,*

$$W(L)^2 \leq (D(r)A_L Q^L)(D(c)C_L Q^L). \quad (16)$$

Proof. The profile factors and the table factorials cancel exactly in the quotient of $W(L)^2$ by $(D(r)A_L)(D(c)C_L)$. It remains to compare the ambient falling factorials and the local full-containment atoms.

Fix one endpoint pair and abbreviate $s = s_{ij}$, $t = t_{ij}$, $d = d_{ij} = t - s$, and $x_{ij} = s$. Its local atom in (13) is

$$\frac{(s)_s (t)_s}{s!} g(s) = \frac{t!}{d!} g(s).$$

Consequently

$$\frac{\left[\frac{(s)_s (t)_s}{s!} g(s)\right]^2}{s! g(s) t! g(t)} = \frac{(t)_d}{(d!)^2} 2^{-ds - \binom{d}{2}} = \frac{Q_{ij}^2}{(n+1)^d}.$$

For all sufficiently large n , one has $s \geq \alpha - 5 \geq 2$. Hence we may use $g(t)/g(s) = 2^{ds + \binom{d}{2}}$; the identity is also valid for $d = 0$.

Moreover, $m(r), m(c) \geq J(L)$ and

$$m(r) + m(c) - 2J(L) = \sum_{i,j} d_{ij} \ell_{ij}.$$

Therefore

$$\frac{(n)_{m(r)} (n)_{m(c)}}{(n)_{J(L)}^2} \leq (n+1)^{m(r)+m(c)-2J(L)}.$$

Combining this inequality with the local ratio cell by cell gives

$$\begin{aligned} \frac{W(L)^2}{(D(r)A_L)(D(c)C_L)} &\leq (n+1)^{m(r)+m(c)-2J(L)} \prod_{i,j} \left(\frac{Q_{ij}^2}{(n+1)^{d_{ij}}} \right)^{\ell_{ij}} \\ &= \prod_{i,j} Q_{ij}^{2\ell_{ij}} = (Q^L)^2, \end{aligned}$$

where the equality uses $m(r) + m(c) - 2J(L) = \sum_{i,j} d_{ij} \ell_{ij}$. This is (16). $\square$

Proposition 8.12 (Endpoint-table sum). *For all sufficiently large n , uniformly in the phase,*

$$\sum_{L \text{ feasible}} W(L) \leq \exp\{O(\tau_n^{\text{end}} K)\} \sum_r D(r) \leq \exp\{O(\sqrt{n \log n})\}. \quad (17)$$

All sums in this proposition are finite.

Proof. By the arithmetic–geometric mean inequality and (16),

$$2W(L) \leq D(r)A_L Q^L + D(c)C_L Q^L.$$

Consider the first term. For a fixed row margin r , feasible tables form a subset of all nonnegative integer tables with row margin r . Since $A_L Q^L \geq 0$, the multinomial theorem gives

$$\sum_{\substack{L \text{ feasible} \\ \text{row}(L)=r}} A_L Q^L \leq \sum_{\substack{L \in \mathbb{N}_0^{3 \times 4} \\ \text{row}(L)=r}} A_L Q^L = \prod_i \left(\sum_j Q_{ij} \right)^{r_i} \leq (1 + C\tau_n^{\text{end}})^K.$$

After multiplication by $D(r)$ and summation over r , the first half is at most $(1 + C\tau_n^{\text{end}})^K \sum_r D(r)$. The second half is identical after exchanging rows and columns. Section 7 proves $\sum_r D(r) = 1 + o(1)$ uniformly in the phase. This is a domination by a reference sum, not a deletion of common-class overlaps. Finally, $K = \Theta(n/\log n)$ and (15) imply $\tau_n^{\text{end}} K = O(\sqrt{n \log n})$. $\square$

8.6. Insertion of the phase estimates. The phase satisfies

$$\alpha = \left(\frac{2}{\log 2} + o(1) \right) \log n.$$

In particular, for all sufficiently large n ,

$$\frac{5}{2} \log n \leq \alpha. \quad (18)$$

Every endpoint minimum satisfies $m_{ij} \geq \alpha - 5$, and elementary floor arithmetic gives

$$3\alpha - 19 \leq 4 \left\lfloor \frac{3m_{ij} - 1}{4} \right\rfloor.$$

Using $\log 2 > 2/3$ and (18), we obtain

$$\frac{5}{4} \log n - \frac{19}{6} \leq (\log 2) \left\lfloor \frac{3m_{ij} - 1}{4} \right\rfloor. \quad (19)$$

Hence

$$2^{\lfloor (3m_{ij}-1)/4 \rfloor} \geq e^{-19/6} n^{5/4}.$$

Since every $m_{ij} = O(\log n)$,

$$\rho_{16} = O\left(\frac{\log n}{n^{1/4}}\right) \rightarrow 0. \quad (20)$$

Thus $\rho_{16} \leq 1$ for all sufficiently large n .

Define the bare high-skeleton sum by

$$\text{BareSkeletonSum}_n := \sum_{(P,j) \text{ feasible}} w_{\text{hi}}(P, j),$$

where the sum ranges over the canonical high-cell matching supports and their admissible high multiplicities defined above.

Proposition 8.13 (Bare high-skeleton estimate). *There is a deterministic sequence $\varepsilon_n \rightarrow 0$ such that*

$$\text{BareSkeletonSum}_n \leq \exp \left\{ \varepsilon_n \frac{n}{(\log n)^4} \right\}. \quad (21)$$

Proof. The deficit factor in (12) satisfies

$$\log(1 + (\alpha + 1)\rho_{16})^K \leq K(\alpha + 1)\rho_{16} = O(n^{3/4} \log n).$$

Proposition 8.12 contributes $O(\sqrt{n \log n})$ to the logarithm. Both quantities are $o(n/(\log n)^4)$. Combining these estimates with (12) proves the proposition. $\square$

No residual local reward or binary cycle-space factor has been included in BareSkeletonSum_n . Section 9 treats exactly those remaining factors. All overlap matrices, including those containing whole common classes, remain part of the exact conditional decomposition there.

9. RESIDUAL ATTACHMENTS BY MATCHING RESTRICTION

Section 8 bounded the sum of the bare high-skeleton weights. It did not include the rewards of unexposed cells or the binary cycle-space factor of the residual support. We now condition on one feasible high skeleton and estimate precisely those remaining factors. Section 7 has not removed any overlap from the exact sum: its diagonal weights enter Section 8 only as a comparison measure for the endpoint-table sum.

Throughout this section, write $Z := Z_{\mathbf{k}}^{\text{sgn}}$ for the selected signed-profile witness count.

9.1. Conditional decomposition. Fix a canonical high skeleton with exposed block matching M , exposed multiplicities j , and total exposed mass J . Put

$$m_0 := n - J.$$

Let $(d_a)_a$ and $(d'_b)_b$ be the residual row and column degrees. Their two sums equal m_0 , and every degree is at most the phase cap U .

The unexposed pairs form a uniform bipartite configuration matching with these residual degrees; when $m_0 = 0$, this means the unique empty matching. Let r'_{ab} be its cell counts and let H_{res} be the simple support graph of cells with $r'_{ab} \geq 2$. The residual event $\mathcal{E}(M, j)$ imposes the cap $r'_{ab} \leq \lfloor U/2 \rfloor$ and forbids any further pair in a cell of M . The latter no-return condition makes the canonical exposure unique. More explicitly, for any fixed set S of J exposed stub pairs and any event $\mathcal{B}$ of the residual matching,

$$\mathbb{P}\{S \subseteq \Pi, \Pi \setminus S \in \mathcal{B}\} = \frac{1}{(n)_J} \mathbb{P}_{\text{res}}(\mathcal{B}). \quad (22)$$

After the physical fiber is summed, the residual attachment depends on the block matching M and the residual degree lists, but not on which labeled stubs realize the exposed pairs. Define

$$\mathcal{A}(M, j) := \mathbb{E}_{\text{res}} \left[\left(\prod_{a,b} g(r'_{ab}) \right) 2^{\beta(M \cup H_{\text{res}})} \mathbf{1}_{\mathcal{E}(M, j)} \right]. \quad (23)$$

Only the residual configuration matching is random in this expectation.

The exact overlap decomposition is

$$\frac{\mathbb{E}Z^2}{(\mathbb{E}Z)^2} = \sum_{(M, j)} w_{\text{hi}}(M, j) \mathcal{A}(M, j), \quad (24)$$

where the sum ranges over the finite family of feasible canonical high skeletons. To justify the word *exact*, start with an overlap matrix r . Its cells above $\lfloor U/2 \rfloor$ determine one and only one matching M and multiplicity vector j . Removing the exposed pairs leaves one residual matrix r' satisfying $\mathcal{E}(M, j)$. Conversely, a feasible skeleton and a residual matching satisfying $\mathcal{E}(M, j)$ reconstruct one overlap matrix. The configuration-model cancellation proved in Section 8 factors its probability and exposed local reward into $w_{\text{hi}}(M, j)$ times the residual law in (23); the remaining sign factor is exactly $2^{\beta(M \cup H_{\text{res}})}$. Thus no overlap is omitted and no overlap is assigned to two skeletons.

The case $m_0 = 0$ is complete without introducing cell activities. The residual matching is empty, every residual local factor is one, and $H_{\text{res}} = \emptyset$. Since M is a matching, $\beta(M) = 0$, so

$$\mathcal{A}(M, j) = 1 \quad (m_0 = 0). \quad (25)$$

In all subsequent subsections we assume $m_0 > 0$. In view of Proposition 8.13, it remains to bound $\mathcal{A}(M, j)$ uniformly in the positive-residual case.

9.2. Threshold expansion and cell activities. For every row–column pair (a, b) , put

$$\theta_{ab} := \frac{e d_a d'_b}{m_0}. \quad (26)$$

The intensity is defined on cells of M as well as outside M ; only the activities will be suppressed on M . Let

$$\Delta_x := g(x) - g(x-1), \quad R := \left\lfloor \frac{U}{2} \right\rfloor,$$

and, for $(a, b) \notin M$, define

$$\lambda_{ab} := \sum_{x=3}^R \Delta_x \frac{\theta_{ab}^x}{x!}, \quad q_{ab} := \frac{\theta_{ab}^2}{2} + \lambda_{ab}. \quad (27)$$

On M , set $\lambda_{ab} = q_{ab} = 0$.

We first record the configuration estimate used in the expansion. For a finite array of nonnegative demands $x = (x_{ab})$ supported outside M , write $x_a = \sum_b x_{ab}$, $x'_b = \sum_a x_{ab}$, and $X = \sum_{a,b} x_{ab}$. Markov's inequality applied to the product of binomial cell counts gives

$$\mathbb{P}_{\text{res}}(r'_{ab} \geq x_{ab} \text{ for all } a, b) \leq \frac{\prod_a (d_a)_{x_a} \prod_b (d'_b)_{x'_b}}{(m_0)_X \prod_{a,b} x_{ab}!}.$$

If $X \leq m_0$, then $(m_0)_X \geq (m_0/e)^X$; if $X > m_0$, the event is empty. Hence

$$\mathbb{P}_{\text{res}}(r'_{ab} \geq x_{ab} \text{ for all } a, b) \leq \prod_{a,b} \frac{\theta_{ab}^{x_{ab}}}{x_{ab}!}. \quad (28)$$

This joint estimate is the reason that cells sharing a row or a column may be expanded simultaneously; no independence between cells is asserted.

For $0 \leq r \leq R$,

$$g(r) = 1 + \sum_{x=3}^R \Delta_x \mathbf{1}_{\{r \geq x\}}.$$

Let E_0 be the set of all residual cell positions outside M , and let $C(M)$ be the family of even subsets of $M \cup E_0$. For $F \in C(M)$, define

$$\begin{aligned} \Phi_F(r') &:= \prod_{e \in F \setminus M} \left(\mathbf{1}_{\{r'_e \geq 2\}} + \sum_{x=3}^R \Delta_x \mathbf{1}_{\{r'_e \geq x\}} \right) \\ &\quad \times \prod_{e \in E_0 \setminus F} \left(1 + \sum_{x=3}^R \Delta_x \mathbf{1}_{\{r'_e \geq x\}} \right). \end{aligned}$$

The first product contains the support threshold required when e belongs to the even set. On the event $\mathcal{E}(M, j)$, expansion of the cycle-space cardinality and of all local rewards gives the exact identity

$$\left(\prod_{a,b} g(r'_{ab}) \right) 2^{\beta(M \cup H_{\text{res}})} = \sum_{F \in C(M)} \Phi_F(r').$$

Lemma 9.1 (Fixed even-set expansion). *For every $F \in C(M)$,*

$$\mathbb{E}_{\text{res}}[\Phi_F(r') \mathbf{1}_{\mathcal{E}(M, j)}] \leq \prod_{e \in F \setminus M} q_e \prod_{e \in E_0 \setminus F} (1 + \lambda_e).$$

Proof. Expand Φ_F into its nonnegative threshold monomials. In a cell of $F \setminus M$, a monomial chooses either the threshold-two term or one higher increment; it never chooses both. In a cell outside $F \cup M$, it chooses nothing or one higher increment. Apply (28) once to the complete demand of each monomial. The threshold-two contribution is $\theta_e^2/2$, the higher increments sum to λ_e , and the empty choice contributes one. Every monomial is nonnegative and $\mathbf{1}_{\mathcal{E}(M, j)} \leq 1$, so omitting the cap and no-return indicator can only increase the expectation. $\square$

Summing over F and then inserting the missing factors $1 + \lambda_e \geq 1$ gives

$$\mathcal{A}(M, j) \leq \left(\prod_{e \in E_0} (1 + \lambda_e) \right) \sum_{F \in C(M)} \prod_{e \in F \setminus M} q_e. \quad (29)$$

9.3. Restriction outside the exposed matching. The next finite lemma is independent of random graphs.

Lemma 9.2 (Restriction-product bound). *Let E be a finite set, let C be a finite family of subsets of E , and let $I \subseteq E$. Suppose that*

$$A \mapsto A \setminus I$$

is injective on C . For nonnegative activities $(q_e)_{e \in E}$,

$$\sum_{A \in C} \prod_{e \in A \setminus I} q_e \leq \prod_{e \in E \setminus I} (1 + q_e).$$

Proof. The restrictions $A \setminus I$ form a subfamily of the power set of $E \setminus I$ and occur without repetition. Enlarge the sum to the full power set and expand the finite product. $\square$

Apply the lemma with $E = M \cup E_0$, $I = M$, and $C = C(M)$. To verify injectivity, suppose that two even sets have the same restriction outside M . Their symmetric difference is an even subset of the matching M . Every nonempty subset of a matching has a vertex of degree one, so this symmetric difference must be empty. Therefore

$$\sum_{F \in C(M)} \prod_{e \in F \setminus M} q_e \leq \prod_{e \in E_0} (1 + q_e). \quad (30)$$

Since $0 \leq \lambda_e \leq q_e$, equations (29) and (30), together with $1 + x \leq e^x$, imply

$$\mathcal{A}(M, j) \leq \exp \left(2 \sum_{e \in E_0} q_e \right). \quad (31)$$

Thus both the local rewards and the cycle-space cardinality are controlled by one total residual activity.

Remark 9.3. Lemma 9.2 applies to any weighted set family whose restriction map is injective. Here the family happens to be a binary cycle space and injectivity follows solely from the fact that the deleted edge set is a matching.

9.4. The intrinsic residual regime. Assume

$$2^U \leq m_0^3. \quad (32)$$

Then $m_0 \geq 2^{U/3}$ and, by the degree cap,

$$\theta_{ab} \leq \epsilon U^2 2^{-U/3}.$$

Lemma 9.4 (Quadratic activity bound). *There is an absolute constant C_0 such that, under (32),*

$$q_{ab} \leq C_0 \theta_{ab}^2 \quad (33)$$

for every residual cell outside M .

Proof. Fix a cell and write $\theta = \theta_{ab}$. If $\theta = 0$, then $q_{ab} = 0$, so the claim holds. Assume $\theta > 0$. Since $0 \leq \Delta_x \leq g(x)$, it is enough to bound

$$a_x := g(x) \frac{\theta^x}{x!}, \quad 3 \leq x \leq R.$$

For $x \geq 3$,

$$\frac{a_{x+1}}{a_x} = \frac{2^x \theta}{x+1}, \quad \frac{a_{x+2}/a_{x+1}}{a_{x+1}/a_x} = 2 \frac{x+1}{x+2} > 1.$$

Thus (a_x) is log-convex, and its maximum on $[3, R]$ occurs at an endpoint. Consequently

$$\lambda_{ab} \leq R(a_3 + a_R).$$

Now $a_3 = (2/3)\theta^3$. Since $R\theta = O(U^3 2^{-U/3})$, one has $Ra_3 \leq \theta^2$ for all sufficiently large U .

For the other endpoint, using $R = \lfloor U/2 \rfloor$ and $\theta \leq eU^2 2^{-U/3}$ gives

$$\log_2 \theta \leq \log_2 e + 2 \log_2 U - \frac{U}{3}.$$

For sufficiently large U , one has $R \geq 3$ and $g(R) = 2^{\binom{R}{2}-1}$. Hence

$$\begin{aligned} \log_2 \left(\frac{a_R}{\theta^2} \right) &= \binom{R}{2} - 1 + (R-2) \log_2 \theta - \log_2(R!) \\ &\leq \left(\frac{U^2}{8} + O(U) \right) - \left(\frac{U^2}{6} + O(U) \right) + O(U \log U) \\ &= -\frac{U^2}{24} + O(U \log U). \end{aligned}$$

The two quadratic terms come respectively from the local reward $g(R)$ and from the factor θ^{R-2} ; the factorial term only improves the upper bound. Hence $Ra_R \leq \theta^2$ for all sufficiently large U . It follows that $\lambda_{ab} \leq 2\theta^2$ in that range. For the finitely many remaining values of U , the quotient q_{ab}/θ^2 is bounded on the compact interval $0 \leq \theta \leq eU^2 2^{-U/3}$, with limit $1/2$ at $\theta = 0$. Enlarging the constant proves the lemma. $\square$

Because θ_{ab} was defined for every pair, the cell intensities satisfy the exact identity

$$\sum_{a,b} \theta_{ab}^2 = \frac{e^2}{m_0^2} \left(\sum_a d_a^2 \right) \left(\sum_b (d'_b)^2 \right).$$

Since every degree is at most U and both degree sums equal m_0 ,

$$\sum_a d_a^2 \leq Um_0, \quad \sum_b (d'_b)^2 \leq Um_0.$$

On M the activities are zero, while outside M the preceding lemma applies. Therefore

$$\sum_{e \in E_0} q_e \leq C_1 U^2. \quad (34)$$

Together with (31), this proves

$$\mathcal{A}(M, j) \leq \exp(CU^2) \quad \text{if } 2^U \leq m_0^3. \quad (35)$$

9.5. The complementary residual regime. Suppose instead that $2^U > m_0^3$. Then

$$m_0 < 2^{U/3} \leq 2^{\lceil U/3 \rceil}. \quad (36)$$

We first bound the integrand pointwise. Since the integrand is nonnegative and $0 \leq \mathbf{1}_{\mathcal{E}(M,j)} \leq 1$, equation (23) remains bounded above after the indicator is omitted. Each residual cell count is at most U , so

$$\prod_{a,b} g(r'_{ab}) \leq 2^{\sum_{a,b} \binom{r'_{ab}}{2}} \leq 2^{(U-1)m_0/2}.$$

Moreover, every edge of H_{res} uses at least two residual pairs, so $|E(H_{\text{res}})| \leq m_0/2$. The restriction map from even subsets of $M \cup H_{\text{res}}$ to subsets of $E(H_{\text{res}})$ is injective by the matching argument in Lemma 9.2. Hence

$$2^{\beta(M \cup H_{\text{res}})} \leq 2^{|E(H_{\text{res}})|} \leq 2^{m_0/2}.$$

Multiplication yields

$$\mathcal{A}(M, j) \leq 2^{Um_0/2} \quad \text{if } 2^U > m_0^3. \quad (37)$$

Proposition 9.5 (Uniform residual attachment). *There is an absolute constant C such that every feasible canonical high skeleton satisfies*

$$\mathcal{A}(M, j) \leq \begin{cases} 1, & m_0 = 0, \\ \exp(CU^2), & m_0 > 0 \text{ and } 2^U \leq m_0^3, \\ 2^{Um_0/2}, & m_0 > 0 \text{ and } 2^U > m_0^3. \end{cases}$$

9.6. Explicit global logarithmic ledger. The preceding propositions already prove the required normalized second-moment bound. We now record their error terms in a form that can be consumed directly by the amplification theorem and by a theorem-facing formalization.

First combine the three deterministic partial-diagonal errors from Section 7:

$$\varepsilon_n^{\text{pd}} := e^{\varepsilon_n^{\text{empty}}} - 1 + \varepsilon_n^{\text{central}} + \varepsilon_n^{\text{full}}. \quad (9.31)$$

Then $\varepsilon_n^{\text{pd}} \rightarrow 0$ uniformly in the phase, and the disjoint three-range assembly gives

$$\sum_r D(r) \leq 1 + \varepsilon_n^{\text{pd}}. \quad (9.32)$$

Here and below every sum has the finite combinatorial domain specified in its original definition.

Let

$$\tau_n^{\text{end}} := 2^{3/2} \sqrt{\frac{(n+1)a}{2a}}, \quad a = \alpha - 2.$$

This is the same τ_n^{end} as in (15). For all sufficiently large n , $\tau_n^{\text{end}} \leq 1$. Since $Q_{ii} = 1$ and each row has only three off-diagonal positions, (15) gives the explicit row bound

$$\sum_j Q_{ij} \leq 1 + 3\tau_n^{\text{end}}. \quad (9.33)$$

Indeed, for $i \neq j$ one has $Q_{ij} \leq (\tau_n^{\text{end}})^{d_{ij}} / d_{ij}! \leq \tau_n^{\text{end}}$.

Repeating the two arithmetic–geometric-mean sums in the proof of Proposition 8.12, now with the explicit constant in (9.33), gives

$$\sum_L W(L) \leq (1 + 3\tau_n^{\text{end}})^K (1 + \varepsilon_n^{\text{pd}}). \quad (9.34)$$

Combining this with the one-charge deficit reduction (12), define

$$\begin{aligned} \Gamma_n^{\text{skel}} &:= K \log(1 + (\alpha + 1)\rho_{16}) \\ &\quad + K \log(1 + 3\tau_n^{\text{end}}) + \log(1 + \varepsilon_n^{\text{pd}}). \end{aligned} \quad (9.35)$$

With the notation $M = P$, the skeleton sum in (24) is precisely the BareSkeletonSum $_n$ defined in Section 8. For all sufficiently large n ,

$$\text{BareSkeletonSum}_n \leq e^{\Gamma_n^{\text{skel}}}. \quad (9.36)$$

This inequality displays separately the unique ambient deficit charge, the off-diagonal endpoint-table charge, and the partial-diagonal reference mass.

The phase estimates from Section 8 imply

$$\begin{aligned} K \log(1 + (\alpha + 1)\rho_{16}) &\leq K(\alpha + 1)\rho_{16} = O(n^{3/4} \log n), \\ K \log(1 + 3\tau_n^{\text{end}}) &\leq 3K\tau_n^{\text{end}} = O(\sqrt{n \log n}), \\ \log(1 + \varepsilon_n^{\text{pd}}) &= o(1). \end{aligned}$$

Consequently the deterministic normalized skeleton error

$$\varepsilon_n^{\text{skel}} := \frac{(\log n)^4}{n} \Gamma_n^{\text{skel}} \longrightarrow 0. \quad (9.37)$$

Let C_{att} be an absolute constant for which (35) holds, and put $q := \log 2$. Define

$$\Gamma_n^{\text{att}} := \max \left\{ C_{\text{att}} U^2, \frac{q}{2} U 2^{U/3} \right\}. \quad (9.38)$$

Proposition 9.5 implies the uniform bound

$$\mathcal{A}(M, j) \leq e^{\Gamma_n^{\text{att}}} \quad (9.39)$$

for every feasible high skeleton. In the intrinsic regime this is precisely (35). In the complementary regime, $m_0 < 2^{U/3}$ and

$$2^{Um_0/2} = \exp\left(\frac{q}{2} U m_0\right) \leq \exp\left(\frac{q}{2} U 2^{U/3}\right).$$

The zero-residual case contributes one.

Since

$$U = O(\log n), \quad 2^U = \Theta\left(\frac{n^2}{(\log n)^2}\right),$$

one has

$$\Gamma_n^{\text{att}} = O\left((\log n)^2 + n^{2/3}(\log n)^{1/3}\right) = o\left(\frac{n}{(\log n)^4}\right).$$

Thus the deterministic normalized attachment error

$$\varepsilon_n^{\text{att}} := \frac{(\log n)^4}{n} \Gamma_n^{\text{att}} \longrightarrow 0. \quad (9.40)$$

Proposition 9.6 (Explicit normalized second-moment ledger). *Define*

$$\Lambda_n := \Gamma_n^{\text{skel}} + \Gamma_n^{\text{att}} = \left(\varepsilon_n^{\text{skel}} + \varepsilon_n^{\text{att}} \right) \frac{n}{(\log n)^4}. \quad (9.41)$$

Then

$$\Lambda_n = o\left(\frac{n}{(\log n)^4}\right)$$

and

$$1 \leq \frac{\mathbb{E}Z^2}{(\mathbb{E}Z)^2} \leq e^{\Lambda_n}. \quad (9.42)$$

Proof. Insert (9.39) into the exact decomposition (24). Because all summands are nonnegative, the uniform attachment factor may be taken outside the finite skeleton sum. Equation (9.36) then gives

$$\frac{\mathbb{E}Z^2}{(\mathbb{E}Z)^2} \leq e^{\Gamma_n^{\text{att}}} \text{BareSkeletonSum}_n \leq e^{\Gamma_n^{\text{att}} + \Gamma_n^{\text{skel}}} = e^{\Lambda_n}.$$

The lower bound is the nonnegativity-of-variance inequality. Equations (9.37) and (9.40) give the stated order of Λ_n . $\square$

Thus the global second-moment exponent is not an unspecified new error. It is the sum of two deterministic ledgers with disjoint responsibilities:

partial diagonals, deficits, and endpoint-table comparison + residual local rewards and cycle space.

No factor is charged in both ledgers.

10. RARE-EVENT AMPLIFICATION

Proposition 9.6 and (1.4) give the seed

$$\mathbb{P}(Z_{\mathbf{k}}^{\text{sgn}} > 0) \geq e^{-\Lambda_n}. \quad (10.1)$$

A positive signed witness is a cocoloring with k_{co} classes, so the seed implies

$$\mathbb{P}(\zeta(G_n) \leq k_{\text{co}}) \geq e^{-\Lambda_n}. \quad (10.2)$$

This event may still be rare. We turn it into a high-probability cocoloring event by adding the quantity displayed in (10.5), which is later shown to be $o(n/(\log n)^3)$ and hence smaller than the root separation. The argument follows the seed-to-typical principle of Heckel (2025, Theorem 1), using vertex-exposure concentration as in Scott (2017, Theorem 1). The first lemma controls every possible leftover vertex set simultaneously; the second amplifies an arbitrary seed exponent Λ_n .

Lemma 10.1 (Simultaneous leftover coloring). *There is an absolute C_0 such that, with probability $1 - o(1)$, every $S \subseteq [n]$ satisfies*

$$\chi(G_n[S]) \leq C_0 \frac{|S|}{\log n} + n^{1/3}. \quad (10.3)$$

Proof. Let H be the complement of G_n . Put $u_0 = \lceil n^{1/4} \rceil$. For any fixed u_0 -set, (1.5) gives probability $\exp(-\Omega(u_0^2))$ that its H -edge density is below $1/4$. There are at most $\binom{n}{u_0}$ such sets, and for an absolute $c > 0$,

$$\binom{n}{u_0} e^{-cu_0^2} \leq \exp\{u_0 \log(en/u_0) - cu_0^2\} = o(1).$$

Thus a union bound shows that, with probability $1 - o(1)$, every u_0 -set has density at least $1/4$. If S has size $s \geq u_0$, double-counting each edge of $H[S]$ over the u_0 -subsets containing it gives

$$\frac{1}{\binom{s}{u_0}} \sum_{\substack{T \subseteq S \\ |T|=u_0}} \frac{e_H(T)}{\binom{u_0}{2}} = \frac{e_H(S)}{\binom{s}{2}}.$$

Therefore every larger set also has density at least $1/4$.

Let S_0 be any vertex set of size at least $n^{1/3}$. While $|S_t| \geq u_0$, choose $v_t \in S_t$ of maximum degree in $H[S_t]$, define $S_{t+1} := N_H(v_t) \cap S_t$, and put $s_t := |S_t|$. Density at least $1/4$ implies $|N_H(v_t) \cap S_t| \geq (s_t - 1)/4$. Hence

$$s_{t+1} \geq (s_t - 1)/4, \quad s_t \geq 4^{-t} s_0 - 1/3. \quad (10.3a)$$

The sets are nested and $S_{t+1} \subseteq N_H(v_t)$, so every later choice is adjacent in H to every earlier one. Thus the chosen vertices form a clique in H . Since $s_0 \geq \lceil n^{1/3} \rceil$, for every $0 \leq t \leq \lfloor \log n / (13 \log 4) \rfloor$, equation (10.3a) gives

$$s_t \geq n^{1/3-1/13} - \frac{1}{3} = n^{10/39} - \frac{1}{3} \geq n^{1/4}$$

for all sufficiently large n . Thus the procedure constructs at least $c \log n$ vertices for an absolute $c > 0$, and these vertices form an independent set in G_n .

For an arbitrary S , repeatedly remove such independent sets and give each a new color until fewer than $n^{1/3}$ vertices remain; color the rest singly. Every removed set has at least $c \log n$ vertices, so the total number of colors is at most

$$\left\lceil \frac{|S|}{c \log n} \right\rceil + n^{1/3}.$$

After enlarging the absolute constant C_0 , this is (10.3), simultaneously for every S . □

Lemma 10.2 (Amplification from a seed). *There is an absolute constant $C > 0$ and a deterministic $\varepsilon_n = o(1)$ with the following property. Suppose deterministic integers $k_n \geq 0$ and deterministic reals $\Lambda_n \geq 0$ satisfy*

$$\mathbb{P}(\zeta(G_n) \leq k_n) \geq e^{-\Lambda_n}. \quad (10.4)$$

The same C and ε_n , independent of k_n, Λ_n, r , satisfy, uniformly for every deterministic choice $r = r(n) > 0$,

$$\begin{aligned} \mathbb{P}\left(\zeta(G_n) > k_n + C\left(\frac{\sqrt{n\Lambda_n} + \sqrt{nr}}{\log n} + n^{1/3} + 1\right)\right) \\ \leq e^{-r} + \varepsilon_n. \end{aligned} \quad (10.5)$$

Proof. Let $\mathcal{E}_n$ be the simultaneous event in Lemma 10.1, and set $\varepsilon_n := \mathbb{P}(\mathcal{E}_n^c) = o(1)$. This sequence is independent of k_n, Λ_n , and r .

For an induced set W , let $\zeta(W) = \zeta(G_n[W])$, and define

$$S_k = \max\{|W| : \zeta(W) \leq k_n\}. \quad (10.6)$$

Expose the random graph in $n - 1$ independent vertex blocks, where block v contains the edges from v to earlier vertices. Changing one block changes S_k by at most one. Indeed, after deleting the affected vertex, every feasible induced set loses at most one vertex, and the two graph configurations agree on all remaining edges. Thus a maximizer in either configuration yields a feasible set of size at least one less in the other configuration. Therefore (1.3) applies.

Since $S_k = n$ exactly when $\zeta(G_n) \leq k_n$, (10.4) and the upper one-sided bounded-differences tail give

$$e^{-\Lambda_n} \leq \mathbb{P}(S_k - \mathbb{E}[S_k] \geq n - \mathbb{E}[S_k]) \leq \exp\left\{-\frac{2(n - \mathbb{E}[S_k])^2}{n - 1}\right\}.$$

Taking logarithms and rearranging gives

$$n - \mathbb{E}[S_k] \leq \sqrt{(n - 1)\Lambda_n/2}. \quad (10.7)$$

The lower tail with radius $\sqrt{(n - 1)r/2}$ gives, outside an event of probability at most e^{-r} ,

$$n - S_k \leq \sqrt{(n - 1)\Lambda_n/2} + \sqrt{(n - 1)r/2}. \quad (10.8)$$

Choose a maximizing set W and put $V_{\text{left}} = [n] \setminus W$. Concatenating a k_n -part cocoloring of $G_n[W]$ with an ordinary coloring of $G_n[V_{\text{left}}]$ gives

$$\zeta(G_n) \leq k_n + \chi(G_n[V_{\text{left}}]). \quad (10.9)$$

On $\mathcal{E}_n$ and the event in (10.8), apply (10.3) to V_{left} . Since $|V_{\text{left}}| = n - S_k$, substitution into (10.9), followed by an enlargement of the absolute constant C , gives the number of additional parts displayed in (10.5). A union bound gives the failure probability $e^{-r} + \varepsilon_n$, proving (10.5) uniformly in the three deterministic parameters. $\square$

Fix the absolute constant C and the sequence ε_n from Lemma 10.2, and apply that lemma to (10.2). Put

$$r_n = \frac{\sqrt{n}}{(\log n)^2}. \quad (10.10)$$

Proposition 9.6 gives the first estimate below. The definition of r_n and elementary asymptotics give the other two; in particular, $r_n \rightarrow \infty$:

$$\frac{\sqrt{n\Lambda_n}}{\log n} = o(n/(\log n)^3), \quad \frac{\sqrt{nr_n}}{\log n} = o(n/(\log n)^3), \quad n^{1/3} = o(n/(\log n)^3). \quad (10.11)$$

Define the deterministic sequence

$$a_n = C \left(\frac{\sqrt{n\Lambda_n} + \sqrt{nr_n}}{\log n} + n^{1/3} + 1 \right). \quad (10.12)$$

Then (10.11) and Lemma 10.2 give

$$a_n = o(n/(\log n)^3), \quad \mathbb{P}(\zeta(G_n) > k_{\text{co}} + a_n) \leq e^{-r_n} + \varepsilon_n \longrightarrow 0. \quad (10.13)$$

11. FINAL ASSEMBLY AND THE QUANTITATIVE CONSTANT

We now combine the chromatic lower location, the signed cocoloring upper location, and the amplification estimate. The two final events are intersected by a union bound; no independence is used.

11.1. The phase-resolved gap. Let $r_+(n)$ be the ordinary first-moment root and let $r_4^{\text{co}}(n)$ be the signed four-size root. Section 5 gives

$$r_+(n) - r_4^{\text{co}}(n) = \left\lceil \frac{(\log 2)^2}{4} A_4(\delta_n) + o(1) \right\rceil \frac{n}{(\log n)^3}, \quad A_4(\delta) := \log 2 - D_4(\delta). \quad (38)$$

The signed profile is placed at the rounded midpoint

$$k_{\text{co}} = \left\lceil \frac{r_4^{\text{co}}(n) + r_+(n)}{2} \right\rceil.$$

At this fixed value of k_{co} , the tangent correction in Section 5 changes only the four type multiplicities by $O(1)$. Here $i \in \{2, 3, 4, 5\}$, $u_i = \alpha - i$, and k_i is the number of classes of size u_i . The correction preserves exactly both

$$\sum_i k_i = k_{\text{co}} \quad \text{and} \quad \sum_i u_i k_i = n.$$

Thus there is no additional correction to the total number of classes. Hence

$$r_+(n) - k_{\text{co}} = \left\lceil \frac{(\log 2)^2}{8} A_4(\delta_n) + o(1) \right\rceil \frac{n}{(\log n)^3}. \quad (39)$$

Indeed, (5.13) gives $r_+(n) - k_{\text{co}} = \frac{1}{2}(r_+(n) - r_4^{\text{co}}) + O(1)$. The coefficient $1/4$ in (38) is therefore multiplied by $1/2$, giving $1/8$. The ceiling contributes the displayed $O(1)$; the tangent correction preserves k_{co} itself.

Section 4 constructs a deterministic integer k_χ^- such that

$$\mathbb{P}(\chi(G_n) > k_\chi^-) \longrightarrow 1$$

and $k_\chi^- = r_+(n) + O(\log n)$. Combining this with (39) gives

$$k_\chi^- - k_{\text{co}} = \left\lceil \frac{(\log 2)^2}{8} A_4(\delta_n) + o(1) \right\rceil \frac{n}{(\log n)^3}. \quad (40)$$

The amplification result (10.13) gives a deterministic sequence $a_n = o(n/(\log n)^3)$ such that

$$\mathbb{P}(\zeta(G_n) \leq k_{\text{co}} + a_n) \longrightarrow 1.$$

On this event and the chromatic lower event,

$$\chi(G_n) - \zeta(G_n) > k_\chi^- - k_{\text{co}} - a_n.$$

A union bound and (40) therefore give a nonnegative deterministic sequence $\varepsilon_n^{\text{gap}} \rightarrow 0$ for which

$$\mathbb{P}\left(\chi(G_n) - \zeta(G_n) \geq \left\lceil \frac{(\log 2)^2}{8} A_4(\delta_n) - \varepsilon_n^{\text{gap}} \right\rceil \frac{n}{(\log n)^3}\right) \longrightarrow 1. \quad (41)$$

Put $q := \log 2$ and $\gamma_4 := \log(200/153) > 0$. Lemma 5.1 gives $A_4(\delta) > \gamma_4$ throughout the phase. Since $\varepsilon_n^{\text{gap}} \rightarrow 0$, for all sufficiently large n ,

$$\varepsilon_n^{\text{gap}} < \frac{3q^2\gamma_4}{32}.$$

Therefore the coefficient in (41) is strictly larger than

$$\frac{q^2\gamma_4}{8} - \frac{3q^2\gamma_4}{32} = \frac{q^2\gamma_4}{32}.$$

It follows that

$$\mathbb{P}\left(\chi(G_n) - \zeta(G_n) \geq \frac{(\log 2)^2}{32} \log\left(\frac{200}{153}\right) \frac{n}{(\log n)^3}\right) \rightarrow 1.$$

The displayed coefficient equals

$$\frac{(\log 2)^2}{32} \log\left(\frac{200}{153}\right) = 0.004021983962242 \dots$$

This proves the main theorem. Since $n/(\log n)^3 \rightarrow \infty$, the chromatic–cochromatic difference tends to infinity with high probability along the full sequence of integers.

FORMAL VERIFICATION

The exact theorem stated above is proved in Lean 4 by `Erdos625.erdos625 : Erdos625Statement`. The formalization uses the displayed constant $(\log 2)^2 \log(200/153)/32$, the full sequence of natural numbers, and the finite uniform model of $G(n, 1/2)$. The modular project and a generated single-file proof are publicly available at <https://github.com/SamPetkov/Erdos/tree/main/625/formalization>; both use the pinned Lean and Mathlib version `v4.31.0`. The final axiom audit reports only `propext`, `Classical.choice`, and `Quot.sound`.

REFERENCES

- Thomas F. Bloom. Erdős problem #625. <https://www.erdosproblems.com/625>, 2026. Accessed 16 July 2026.
- Béla Bollobás. The chromatic number of random graphs. *Combinatorica*, 8(1):49–55, 1988. doi: 10.1007/BF02122551. URL <https://doi.org/10.1007/BF02122551>.
- Béla Bollobás and Andrew Thomason. Generalized chromatic numbers of random graphs. *Random Structures & Algorithms*, 6(2–3):353–356, 1995. doi: 10.1002/rsa.3240060222. URL <https://doi.org/10.1002/rsa.3240060222>.
- Paul Erdős and John Gimbel. Some problems and results in cochromatic theory. In John Gimbel, John W. Kennedy, and Louis V. Quintas, editors, *Quo Vadis, Graph Theory?*, volume 55 of *Annals of Discrete Mathematics*, pages 261–264. North-Holland/Elsevier, 1993. doi: 10.1016/S0167-5060(08)70393-5. URL [https://doi.org/10.1016/S0167-5060\(08\)70393-5](https://doi.org/10.1016/S0167-5060(08)70393-5).
- John Gimbel. Some of my favorite coloring problems for graphs and digraphs. In Raluca Gera, Stephen T. Hedetniemi, and Craig Larson, editors, *Graph Theory: Favorite Conjectures and Open Problems – 1*, Problem Books in Mathematics, pages 95–108. Springer, Cham, 2016. ISBN 978-3-319-31940-7. doi: 10.1007/978-3-319-31940-7_7. URL https://doi.org/10.1007/978-3-319-31940-7_7.
- Geoffrey R. Grimmett and Colin J. H. McDiarmid. On colouring random graphs. *Mathematical Proceedings of the Cambridge Philosophical Society*, 77(2):313–324, 1975. doi: 10.1017/S0305004100051124. URL <https://doi.org/10.1017/S0305004100051124>.
- Annika Heckel. The chromatic number of dense random graphs. *Random Structures & Algorithms*, 53(1):140–182, 2018. doi: 10.1002/rsa.20757. URL <https://doi.org/10.1002/rsa.20757>.
- Annika Heckel. On a question of Erdős and Gimbel on the cochromatic number. *The Electronic Journal of Combinatorics*, 31(4):P4.72, 2024. doi: 10.37236/13346. URL <https://doi.org/10.37236/13346>.
- Annika Heckel. The difference between the chromatic and the cochromatic number of a random graph, 2025. URL <https://arxiv.org/abs/2409.17614>. Version 2, revised 19 February 2025; first submitted in 2024.
- Annika Heckel and Konstantinos Panagiotou. Colouring random graphs: Tame colourings, 2023. URL <https://arxiv.org/abs/2306.07253>. arXiv:2306.07253.
- Colin McDiarmid. On the method of bounded differences. In Johannes Siemons, editor, *Surveys in Combinatorics, 1989*, volume 141 of *London Mathematical Society Lecture Note Series*, pages 148–188. Cambridge University Press, 1989. doi: 10.1017/CBO9781107359949.008. URL <https://doi.org/10.1017/CBO9781107359949.008>.

- Colin McDiarmid. On the chromatic number of random graphs. *Random Structures & Algorithms*, 1(4):435–442, 1990. doi: 10.1002/rsa.3240010404. URL <https://doi.org/10.1002/rsa.3240010404>.
- Konstantinos Panagiotou and Angelika Steger. A note on the chromatic number of a dense random graph. *Discrete Mathematics*, 309(10):3420–3423, 2009. doi: 10.1016/j.disc.2008.09.019. URL <https://doi.org/10.1016/j.disc.2008.09.019>.
- Edward R. Scheinerman. Generalized chromatic numbers of random graphs. *SIAM Journal on Discrete Mathematics*, 5(1):74–80, 1992. doi: 10.1137/0405006. URL <https://doi.org/10.1137/0405006>.
- Alex Scott. On the concentration of the chromatic number of random graphs, 2017. URL <https://arxiv.org/abs/0806.0178>. Version 2; first submitted in 2008.
- Raphael Steiner. On the difference between the chromatic and cochromatic number. *SIAM Journal on Discrete Mathematics*, 39(4):2268–2274, 2025. doi: 10.1137/24M1715180. URL <https://doi.org/10.1137/24M1715180>. Published online 17 November 2025; arXiv:2408.02400.

ÉCOLE NORMALE SUPÉRIEURE, UNIVERSITÉ PSL, PARIS, FRANCE
 Email address: samuil.petkov@ens.psl.eu